

United States Department of the Interior Fish and Wildlife Service <u>Fishery Leaflet 12</u>	
Chicago, Ill.	January 1943

FERTILIZATION OF FISH PONDS

Prepared in the Division of Fishery Biology

The number of ponds constructed on farms throughout the country has increased amazingly during the past three or four years. Although complete figures are not obtainable, these farm fish ponds undoubtedly are numbered by tens of thousands and represent a potential source of a significant quantity of food.

The poundage of fish yielded by an individual pond depends largely upon how much fish food is available in its waters. Most pondfish eat smaller fishes, water insects, and small crustacea; these, in turn, are dependent upon minute water plants for foods. These plants, most of which are of microscopic size, are unable to grow unless certain chemicals, especially nitrogen, phosphorous, and potash are present in the water in sufficient quantities. If the pond receives water that drains a rich and fertile soil, it may contain an adequate supply of these necessary chemicals and yield as much as 200 pounds of fish per acre. Ordinarily, however, it is desirable, or even essential, to improve upon natural conditions by the use of fertilizers, thus insuring favorable conditions for rapid growth. By proper fertilization the productive capacity of ponds may be enormously increased, the gain in fish production ranging from 50 percent in naturally poorer waters to a threefold increase in ponds draining rich soils.

The general principles of pond fertilization here outlined are applicable to the numerous ponds maintained by State and Federal fish hatcheries, as well as to privately owned and operated farm ponds.

In principle, pond fertilization is similar to the fertilization of land, but the procedure to be followed differs in several details. In land fertilization the soil acts as a storehouse in which the fertilizer is held temporarily until utilized by plants. In ponds, however, the fertilizing elements go into solution in a short time and if not utilized immediately may be carried away and lost.

It must be emphasized that just as various soils need different fertilizers for best results so diverse bodies of water may vary in their requirements. Unfortunately, however, we have little factual information on the particular combination of fertilizers that will give the best results in various types of waters. Consequently, the only safe course is to use

a so-called "complete" fertilizer. The process of pond fertilization is complicated by the evidence that not all fish respond equally well to the same fertilizer. Even if fertilizers that contain all of the essential elements in proper proportion are employed, the kind of fertilizer to be used may depend, to some extent at least, on the species and size of fish present.

ORGANIC AND INORGANIC FERTILIZERS

Both organic and inorganic fertilizers have been used in ponds with satisfactory results. Of the organic fertilizers, stable manure is probably best known, but has two very serious disadvantages. In the first place, it varies so widely in composition that it is impossible to make definite recommendations regarding the amount to be used. Secondly, unless great care is exercised in its use, serious depletion of the dissolved oxygen in the pond is likely to result, with a consequent loss of fish by asphyxiation. These objections do not apply to dried manures which can be obtained in any quantity desired and have been used with satisfactory results.

Cottonseed, soybean, and peanut meals are probably the most satisfactory organic fertilizers. Formerly, it was recommended that these meals be used alone in fertilizing fish ponds, but recent studies indicate that production can be increased by the addition of inorganic fertilizers such as superphosphate and lime.

The inorganic fertilizers are similar to those applied to field crops. They may be purchased in the form of a complete fertilizer or may be mixed by the user.

CHOICE OF FERTILIZER

In the Service's earlier experiments a combination of dried sheep manure and superphosphate was used with excellent results. The most satisfactory mixture consisted of equal parts sheep manure and superphosphate. The superphosphate should contain 16 percent of available phosphoric acid. Later it was found that an equal amount of cottonseed meal, soybean meal, or peanut meal could be substituted for sheep manure with little change in the results. Since the cost of sheep manure, cottonseed meal, soybean meal, and peanut meal is approximately the same, it is largely a matter of personal preference which product is used.

There is increasing evidence that inorganic fertilizers are superior to organic fertilizers. The nutrient elements are more readily dissolved in the water and produce their effects almost at once. Furthermore, the use of inorganic fertilizers apparently offers greater opportunities for the control of the larger submerged vegetation. Experiments at Leesport, W. Va., during the past 3 years have revealed combinations which will prohibit or actually eliminate the growth of Chara and Potamogetons by stimulating the growth of forms of algae which smother them out or otherwise prevent them from growing.

Recently Swingle and Smith at the Alabama Agricultural Experiment Station have been conducting extensive experiments with inorganic fertilizers.

They originally recommended the following 7-8-2^{1/} mixture as giving most satisfactory results:

40 pounds sulphate of ammonia
60 pounds superphosphate (16 percent)
5 pounds muriate of potash
30 pounds basic slag, or
15 pounds finely ground limestone

The basic slag or limestone is added to neutralize the acid formed when the ammonium sulphate is broken down and probably will be unnecessary in waters containing considerable quantities of lime.

More recently Swingle and Smith have recommended a ready-mixed 6-3-4 fertilizer to which 10 pounds of nitrate of soda (in soft waters) or 10 pounds of ammonium sulphate (in hard waters) are to be added to each 100 pounds of the fertilizer. The above amounts are sufficient for one application to an acre of water. With the use of this fertilizer Swingle and Smith reported annual yields up to 400 pounds of fish per acre. This is a remarkably high production and probably cannot be realized under ordinary conditions.

The advisability of adding larger amounts of nitrogen and potash to the inorganic fertilizer combinations, particularly in hard waters, is indicated by recent experiments at Leetown where a 12-5-5 combination set a record production of 262 pounds of fingerling bass per acre and by the recent publications of Swingle and Smith in which nitrogen and potash ratios have been increased.

AMOUNT OF FERTILIZER REQUIRED

The amount of fertilizer required for best results will, of course, depend on local conditions. A pond that is naturally productive will require less fertilizer than one that is relatively unproductive. Furthermore, ponds with considerable overflow will require more fertilizer than those in which there is little loss of water. When using a dry organic or inorganic fertilizer from 500 to 1,000 pounds per acre for the season ordinarily will be sufficient. In ponds notably lacking in fertility larger amounts may be used to advantage, but as a rule this is not necessary and simply results in waste.

COMPOSITION AND APPLICATION OF FERTILIZER

Tables 1 and 2 list single fertilizers and combinations of fertilizers which should give satisfactory results when used as recommended. At present, it is impossible to make definite recommendations as to which fertilizer to use and the fish-culturist should select the one that seems best suited to his particular needs. Only by experimentation can he determine which will consistently give best results under his conditions.

^{1/} The active elements in a fertilizer are nitrogen, phosphorus, and potash. The characteristics of an individual fertilizer depend on the proportions of these substances, which are indicated by numerals. Thus a "7-8-2 mixture" contains 7 parts nitrogen to 8 parts phosphorus to 2 parts of potash.

Table 1.--Application of Organic Fertilizer

Kind of fertilizer	Pounds per acre per application	Frequency of applications (nursery ponds)	Approximate number of applications during a season (nursery ponds)	Kind of water
Cottonseed meal	50	Every 7 days for first 7 weeks; 14 days thereafter	14	Soft or moderately hard
Soybean meal	50	do	14	Do
Peanut meal	50	do	14	Do
Cottonseed meal (2 parts): 16 percent superphosphate (1 part)	50	do	14	Do
Cottonseed meal (1 part): 16 percent superphosphate (1 part)	50	do	14	Do
Dried sheep manure (1 part): 16 percent superphosphate (1 part)	50	do	14	Do
Horse or fresh cow manure	1,000 first application; 500 thereafter	Every three weeks	6	Do

Table 2.--Application of Inorganic Fertilizer

Kind and composition of fertilizer	Pounds per acre per application	Frequency of application	Kind of water
Inorganic 12-5-5, per application: 60 lbs. ammonium sulphate (20.57 percent N) 30 lbs. superphosphate (16 percent P ₂ O ₅) 10 lbs. muriate of potash (50 percent K ₂ O)	100	Every 10 days for first 3 applications; monthly thereafter or as often as necessary	Hard
Inorganic 8-7-2, ^{1/} per application: 40 lbs. ammonium sulphate (20.57 percent N) 60 lbs. superphosphate (16 percent) 5 lbs. muriate of potash (50 percent K ₂ O) ^{2/} 15 lbs. finely ground limestone 10 lbs. nitrate of soda (16 percent N)	130	Weekly for first 3 weeks. As often as necessary thereafter. ^{1/}	Soft
Inorganic 9-8-2, ^{1/} per application: 50 lbs. ammonium sulphate (20.57 percent N) 60 lbs. superphosphate (16 percent P ₂ O ₅) 5 lbs. muriate of potash (50 percent K ₂ O)	115	Weekly for first 3 weeks. As often as necessary thereafter. ^{1/}	Hard

^{1/}Recommended by Swingle and Smith.

^{2/}Swingle and Smith do not show potash content of the muriate of potash used in their work. Five pounds of 50 percent K₂O in muriate of potash would yield only 2 percent potash in the combination. Commercial muriate of potash contains from 48-62 per cent K₂O, but is usually offered at 50 percent. Swingle and Smith offer as an alternate to this 100 pounds of 6-8-4 inorganic fertilizer plus 10 pounds of nitrate of soda for use in farm ponds.

^{3/}Swingle and Smith recommended 3 or 4 applications of inorganic fertilizer at weekly intervals or until a "bloom" appears in the water. At this time the water becomes murky and green or brown due to the growth of the plankton. Later applications should be made when the bottom becomes visible at 12-18 inches (approximately every 4 weeks.) In farm ponds the last application should be made in September.

LITERATURE CITED

- DAVIS, H. S., and A. H. WIEBE.
1931. Experiments in the culture of black bass and other pondfish. Report, U. S. Commissioner of Fisheries, fiscal year 1930. Appendix IX, pp. 177-203. (Out of print)
- HOGAN, JOE.
1933. Experiments with commercial fertilizers in rearing largemouth black bass fingerlings. Transactions, American Fisheries Society, Vol. 63, pp. 110-119.
- MEEHEAN, O. LLOYD.
1933. The role of fertilizers in pondfish. Transactions, American Fisheries Society, Vol. 63, pp. 103-109.
1935. The dispersal of fertilizing substances in ponds. Transactions, American Fisheries Society, Vol. 65, pp. 184-188.
- NELSON, T. F.
1941. Fertilizing bass ponds. The Progressive Fish Culturist No. 56, pp. 28-29.
- SMITH, E. V., and H. S. SWINGLE.
1939. The relationship between plankton production and fish production in ponds. Transactions, American Fisheries Society, Vol. 68, 1938, pp. 309-315.
1940. Effect of organic and inorganic fertilizers on plankton-carrying capacity and bluegill-bream carrying capacity of ponds. Transactions, American Fisheries Society, Vol. 69, 1939, pp. 257-262.
1941. The use of fertilizer for controlling several submerged aquatic plants in ponds. Transactions, American Fisheries Society, Vol. 71, pp. 94-101.
- SWINGLE, H. S., and E. V. SMITH.
1939. Fertilizers for increasing the natural food of fish in ponds. Transactions, American Fisheries Society, Vol. 68, 1938, pp. 126-135.
1939. Increasing fish production in ponds. Transactions, Fourth North American Wildlife Conference, 1939, pp. 332-338.
1940. Fish production in terrace-water ponds in Alabama. Transactions, American Fisheries Society, Vol. 69, 1939, pp. 101-105.
1942. Management of Farm Fish Ponds. Bulletin 254. Agr. Exp. Station, Alabama Polytechnic Institute.