

Supplemental Table 2. Pigmentation of dorsal and ventral caudal patches in larval *Sebastes moseri* and *S. rufinanus*. BL = body length; Start = myomere at which pigment patch begins; End = myomere at which pigment patch ends; # = number of melanophores in pigment patch; opaque = myomeres not countable.

BL (mm), <i>n</i>	Dorsal pigment			Ventral pigment		
	Start	End	#	Start	End	#
	mode/ range	mode/ range	mode/ range	mode/ range	mode/ range	mode/ range
<i>Sebastes moseri</i>						
2.4-2.9 7	20, 21 20-22	25 24-25	4 4-6	22, 23 21-23	25 24-26	3 2-5
3.0-3.9 40	22 20-23	26 24-26	7 3-8	24 22-25	25 24-26	4 2-6
4.0-4.9 28	21 19-23	25, 26 24-26	7 5-9	22 20-24	25 23-26	4 3-6
5.0-5.5 8	20 19-22	26 25-26	7 5-9	21 20-23	26 24-26	4, 5 4-7
6.2-6.8 3	19 19	26 25-26	7 7-9	20 20-21	25 24-25	7 4-7
7.1 2	19-20	26	10, 12	22	25-26	8-9
8.8-8.9 2	opaque	opaque	14	opaque	opaque	5, 10
<i>Sebastes rufinanus</i>						
2.6-2.9 6	22 21-23	26 24-26	4 4-5	23 23-25	26 24-26	3 3-4
3.0-3.9 20	21 20-22	25 24-26	6 4-8	22 21-23	25 24-26	4 3-7
4.0-4.8 15	21 18-22	26 25-26	7 5-10	23 20-24	25 24-26	5 3-7
5.0-5.8 7	19, 20 19-23	26 25-26	8 5-9	20 20-24	25 24-26	4 4-7