


**SARDINE EGGS AND LARVAE
AND OTHER FISH LARVAE
PACIFIC COAST, 1957**



SPECIAL SCIENTIFIC REPORT-FISHERIES No. 328

**UNITED STATES DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE**

Explanatory Note

The Special Scientific Report series comprises results of investigations, usually of restricted scope, intended to aid or direct management or utilization practices and as guides for administrative or legislative action. Reports in this series are issued in limited quantities for the official use of Federal, State, or other cooperating agencies, and in processed form for economy and to avoid delay in publication.

United States Department of the Interior, Fred A. Seaton, Secretary
Fish and Wildlife Service, Arnie J. Suomela, Commissioner

SARDINE EGGS AND LARVAE AND OTHER FISH LARVAE
PACIFIC COAST, 1957

by

Elbert H. Ahlstrom
Fishery Research Biologist
Bureau of Commercial Fisheries
La Jolla, California



United States Fish and Wildlife Service
Special Scientific Report--Fisheries No. 328

Washington, D. C.
December 1959

TABLE OF CONTENTS

| | <u>Page</u> |
|---|-------------|
| Area covered, 1957 | 3 |
| Methods of sampling | 5 |
| Abundance of fish larvae in 1957 | 5 |
| Record of standardized haul factors for oblique hauls made with plankton nets during cruises 5701-5712 | 8 |
| Record of sardine eggs, 1957 | 18 |
| Record of sardine larvae, 1957 | 27 |
| Record of anchovy larvae, 1957 | 37 |
| Record of the larvae of the jack mackerel (<u>Trachurus symmetri-</u> <u>cus</u>), 1957 | 59 |
| Record of the larvae of the Pacific mackerel (<u>Pneumatophorus</u> <u>diego</u>), 1957 | 73 |
| Record of the larvae of the hake (<u>Merluccius productus</u>), 1957 | 79 |
| Record of the larvae of rockfish (<u>Sebastes</u> spp.), 1957 | 89 |
| Literature cited | 98 |

FIGURES

Nos.

| | |
|--|----|
| 1. Station plan, 1957, of the California Cooperative Oceanic Fisheries Investigations | 2 |
| 2. Sardine eggs, 1957: Distribution and relative abundance | 20 |
| 3. Sardine larvae, 1957: Distribution and relative abundance | 26 |
| 4. Anchovy larvae, 1957: Distribution and relative abundance | 36 |
| 5. Jack mackerel larvae, 1957: Distribution and relative abundance | 60 |
| 6. Pacific mackerel larvae, 1957: Distribution and relative abundance | 72 |
| 7. Hake larvae, 1957: Distribution and relative abundance | 78 |

TEXT TABLES

| Nos. | Page |
|--|------|
| 1. Coverage during 1957 | 4 |
| 2. Abundance (standard haul totals) of fish larvae in 1957, summarized by month | 6 |
| 3. Temperature ranges and means for stations containing sardine eggs in 1957, summarized by month and area (temperatures at 10 M) | 21 |
| 4. Occurrence and abundance (standard haul totals) of sardine eggs by month and area, in hauls made during 1957 | 22 |
| 5. Occurrence and abundance (standard haul totals) of sardine larvae, by month and area, in hauls made during 1957 | 29 |
| 6. Occurrence and abundance (standard haul totals) of anchovy larvae (<u>Engraulis mordax</u>), by month and area, in hauls made during 1957 | 39 |
| 7. Monthly abundance of anchovy larvae, 1953-1957, based on standard haul summations | 40 |
| 8. Abundance of anchovy larvae by size categories, 1953-1957, based on standard haul summations | 40 |
| 9. Occurrence and abundance (standard haul totals) of jack mackerel larvae (<u>Trachurus symmetricus</u>), by month and area, in hauls made during 1957. | 61 |
| 10. Occurrence and abundance (standard haul totals) of Pacific mackerel larvae (<u>Pneumatophorus diego</u>), by month and area, in hauls made during 1957 | 74 |
| 11. Occurrence and abundance (standard haul totals) of hake larvae (<u>Merluccius productus</u>), by month and area, in hauls made during 1957 | 80 |
| 12. Occurrence and abundance (standard haul totals) of rockfish larvae (<u>Sebastes</u> spp.), by month and area, in hauls made during 1957 | 90 |

TABLES

| Nos. | | Page |
|-------|---|------|
| I. | Record of standardized haul factors for oblique hauls made with plankton nets during cruises 5701-5712, 1957 | 9 |
| II. | Record of sardine eggs, 1957 | 23 |
| III. | Record of sardine larvae, 1957 | 30 |
| IV. | Record of anchovy larvae, 1957 | 41 |
| V. | Record of the larvae of jack mackerel (<u>Trachurus</u> <u>symmetricus</u>), 1957 | 63 |
| VI. | Record of the larvae of Pacific mackerel (<u>Pneumatophorus</u> <u>diego</u>), 1957 | 75 |
| VII. | Record of the larvae of hake (<u>Merluccius productus</u>), 1957) . . . | 81 |
| VIII. | Record of the larvae of rockfish (<u>Sebastes</u> spp.), 1957. | 91 |

ABSTRACT

©

This report contains the results of quantitative sampling of fish eggs and larvae off the coasts of California and Baja California during 1957 on cruises of the California Cooperative Oceanic Fisheries Investigations. It is the eighth in a continuing yearly series.

In addition to eggs and larvae of the Pacific sardine (Sardinops caerulea), larvae of the following are dealt with: northern anchovy (Engraulis mordax), jack mackerel (Trachurus symmetricus), Pacific mackerel (Pneumatophorus diego), hake (Merluccius productus), and rockfish (Sebastes spp.). Larvae of all species except hake and rockfish are reported by size. The report includes charts showing distribution and relative abundance of the above species (except rockfish) and a brief discussion of each.

SARDINE EGGS AND LARVAE AND OTHER FISH LARVAE
PACIFIC COAST, 1957

The first report in this series, which appeared as Special Scientific Report: Fisheries No. 80, contained a record of Pacific sardine or pilchard (Sardinops caerulea) eggs and larvae for 1950, and of larvae of the northern anchovy (Engraulis mordax), jack mackerel (Trachurus symmetricus), hake (Merluccius productus) and rockfish (Sebastes spp.). Larvae of sardine and anchovy were reported by size; those of the other species by total number per station. The initial paper contained a more detailed account of methods of sampling and standardization of hauls than subsequent ones. The report has gradually evolved into its present format. A record of total number of larvae of Pacific mackerel (Pneumatophorus diego) per station was added to the report dealing with 1951 collections (Spec. Sci. Rept: Fisheries No. 102). Jack mackerel larvae were reported by size in the report for 1952 (Spec. Sci. Rept.: Fisheries No. 123) and subsequently. Pacific mackerel larvae were reported by size since 1953 (Spec. Sci. Rept.: Fisheries No. 155). Beginning with the report for 1955, yearly distribution charts were included for sardine eggs, and for larvae of sardine, anchovy, jack mackerel, Pacific mackerel, and hake; a series of summary tables (text tables) were added, along with a partial analysis of the data on distribution and abundance of each category. The present report is the eighth in a continuing series.

As in previous reports, the basic data are presented in eight tables, designated by Roman numerals:

- I. Record of standardized haul factors for oblique hauls made with plankton nets during cruises 5701-5712, 1957.
- II. Record of sardine eggs, reported by age in days.
- III. Record of all hauls containing sardine larvae, reported by size groups.
- IV. Record of all hauls containing anchovy larvae, reported by size groups.
- V. Record of all hauls containing jack mackerel larvae, reported by size groups.
- VI. Record of all hauls containing Pacific mackerel larvae, reported by size groups.
- VII. Hake larvae, reported by number per station.
- VIII. Rockfish larvae, reported by number per station.

The fish eggs and larvae included in this report were obtained in quantitative plankton hauls on cruises of the California Cooperative Oceanic Fisheries Investigations. The investigations are sponsored by the California Marine Research Committee and are carried out cooperatively by the Biological Laboratory, La Jolla, of the U. S. Bureau of Commercial Fisheries, the Scripps Institution of Oceanography




Figure 1.--Station plan, 1957, of the California Cooperative Oceanic Fisheries Investigations.

of the University of California, the Hopkins Marine Station of Stanford University, the California Department of Fish and Game, and the California Academy of Sciences. The biological-oceanographic cruises are a primary responsibility of the Scripps Institution of Oceanography and the Bureau of Commercial Fisheries.

It is with deep pleasure that I acknowledge the cooperation given by the Scripps Institution of Oceanography in the collection of data at sea. Most personnel of the Biological Laboratory, La Jolla, contributed to this project, many devoting their full time to it. Lois Hunter did most of the work of identifying, enumerating and measuring fish eggs and larvae in the 1957 material; James Thrailkill supervised the separation of fish eggs and larvae from plankton collections and also prepared the charts included in this report; standardization of egg and larval counts and preparation of the basic tables were done by Margaret Ahlstrom.

AREA COVERED, 1957

The area surveyed during 1957 is shown in figure 1. Stations occupied on three or more cruises are shown as black dots, those occupied on 1 or 2 cruises as open circles. The most extensive coverage was obtained during June, when the area surveyed extended from off San Francisco, California, (line 60) to below Cape San Lucas, Baja California, (line 157) and offshore to station 90 on many lines. The January cruise was an abbreviated one, because one of the two vessels scheduled to make the cruise was laid up for repairs. Extensive coverage was obtained on seven cruises - February through July and October (text table 1). The area off central Baja California only was surveyed during August and September, while the coverage during November and December was mostly confined to waters off southern California.

The Gulf of California was surveyed on four cruises during 1957 - in February, April, June, and August. Data from the Gulf cruises are not included in this report.

One to four vessels participated in each cruise. The "Black Douglas" made eight survey cruises (February through September). Five vessels of the Scripps Institution of Oceanography participated in one or more cruises, as follows: "Horizon", four cruises (February, April-June); "Spencer F. Baird", one cruise (February); "Stranger", seven cruises (April-August, October, December); "Orca", six cruises (January, March, May-July, November); "Paolina T", three cruises (July, October, December). Vessels making the Gulf of California cruises are given in the above listing, since these vessels occupied stations on lines 153 and 157 of the regular pattern in addition to the Gulf stations. The vessels used for Gulf of California cruises were as follows: February - "Spencer F. Baird", April - "Black Douglas", June - "Stranger", August - "Stranger".

Text table 1. --Coverage during 1957

| Month | Cruise number | Number vessels participating | Area covered | Number of stations occupied in each subarea | | | | | | | | | | Total stations |
|-----------|---------------|------------------------------|--------------|---|-------|--------|---------|---------|---------|-------|-------|-------|-------|----------------|
| | | | | 60-77 | 80-93 | 97-107 | 110-120 | 123-137 | 140-157 | Lines | Lines | Lines | Lines | |
| January | 5701 | 1 | 100-137 | - | - | 12 | 25 | 20 | - | - | - | 57 | | |
| February | 5702 | 3 | 80-157 | - | 35 | 27 | 31 | 25 | 24 | - | - | 142 | | |
| March | 5703 | 2 | 80-150 | - | 34 | 19 | 28 | 26 | 17 | - | - | 124 | | |
| April | 5704 | 3 | 80-157 | - | 41 | 38 | 39 | 43 | 46 | - | - | 207 | | |
| May | 5705 | 4 | 60-137 | 32 | 54 | 52 | 50 | 17 | - | - | - | 205 | | |
| June | 5706 | 4 | 60-157 | 36 | 57 | 51 | 56 | 30 | 23 | - | - | 253 | | |
| July | 5707 | 4 | 60-137 | 26 | 54 | 45 | 50 | 42 | - | - | - | 217 | | |
| August | 5708 | 2 | 110-137 | - | - | - | 23 | 14 | - | - | - | 37 | | |
| | | | 153-157 | - | - | - | - | - | 4 | - | - | 4 | | |
| September | 5709 | 1 | 110-137 | - | - | - | 23 | 14 | - | - | - | 37 | | |
| October | 5710 | 2 | 80-137 | - | 41 | 32 | 34 | 36 | - | - | - | 143 | | |
| November | 5711 | 1 | 70-97 | 7 | 30 | 7 | - | - | - | - | - | 44 | | |
| December | 5712 | 2 | 80-90(110) | - | 18 | 4 | 1 | - | - | - | - | 23 | | |
| Total | | | 60-157 | 101 | 364 | 287 | 360 | 267 | 114 | - | - | 1,493 | | |

METHODS OF SAMPLING

Plankton nets used during 1957 were either constructed of No. 30xxx grit gauze, a heavy grade of silk bolting cloth, or No. 471 "Nitex", a nylon monofilament screen cloth. The mesh opening of Nitex cloth measures approximately 0.47 mm. The mesh opening of No. 30xxx grit gauze measures approximately 0.70 mm. before use, but soon shrinks to approximately 0.55 mm. between threads. Mesh openings in nets constructed of Nitex tend to enlarge slightly, hence the mesh openings of the two kinds of cloth, after use, are not too dissimilar.

The rapidity with which nylon cloth dries has proved to be a liability in plankton sampling. Unless a nylon net is very thoroughly washed after each use, any adhering plankton dries against and clings to the nylon threads, soon causing the nets to clog badly. Plankton nets are hosed down after each haul, but only scrubbed at intervals during a cruise. Silk nets, which dry more slowly and consequently remain damp between stations, do not develop this type of clogging problem.

Plankton hauls are made obliquely from approximately 140 meters deep to the surface (200 meters of wire out at greatest depth) where depth of water permits; at lesser depths in shallow water. The research vessel moves slowly ahead during a plankton haul, usually at a speed of between 1 and 2 knots. The amount of water strained during a haul is determined from the revolutions registered by a current meter fastened in the center of the mouth of the net.

ABUNDANCE OF FISH LARVAE IN 1957

Fish larvae were more abundant in 1957 than in the preceding two years, as is shown by the average number of larvae obtained per station (standardized counts):

| Year | Number of stations occupied | Total number of larvae | Average per station |
|------|-----------------------------|------------------------|---------------------|
| 1955 | 1375 | 359,155 | 261 |
| 1956 | 1397 | 408,140 | 292 |
| 1957 | 1493 | 493,549 | 331 |

Text table 2.--Abundance (standard haul totals)
of fish larvae in 1957, summarized by month

| | Sardine | Anchovy | Jack | | Pacific | | Hake | Rockfish | All other | |
|-----------|---------|---------|----------|----------|----------|-------------|---------|----------|-----------|--|
| | | | mackerel | mackerel | mackerel | fish larvae | | | Total | |
| January | 759 | 10,499 | 2 | 6 | 1,278 | 1,010 | 3,591 | 17,145 | | |
| February | 945 | 48,261 | 565 | 34 | 37,998 | 11,001 | 17,662 | 116,466 | | |
| March | 1,164 | 14,328 | 331 | 10 | 34,589 | 6,585 | 30,889 | 87,896 | | |
| April | 423 | 20,232 | 1,302 | 44 | 4,047 | 7,225 | 25,372 | 58,645 | | |
| May | 1,730 | 20,796 | 10,850 | 75 | 260 | 4,819 | 29,300 | 67,830 | | |
| June | 284 | 18,406 | 6,188 | 211 | 88 | 2,742 | 27,954 | 55,873 | | |
| July | 1,446 | 8,886 | 752 | 191 | 16 | 1,383 | 25,029 | 37,703 | | |
| August | 1,513 | 1,446 | 8 | 663 | 3 | 47 | 12,853 | 16,533 | | |
| September | 1,036 | 35 | 0 | 334 | 0 | 28 | 9,553 | 10,986 | | |
| October | 315 | 862 | 8 | 298 | 0 | 231 | 17,158 | 18,872 | | |
| November | 218 | 2,372 | 0 | 0 | 4 | 586 | 581 | 3,761 | | |
| December | 0 | 508 | 0 | 0 | 0 | 816 | 515 | 1,839 | | |
| Total | 9,833 | 146,631 | 20,006 | 1,866 | 78,283 | 36,473 | 200,457 | 493,549 | | |
| Percent | 2.0 | 29.7 | 4.1 | 0.4 | 15.8 | 7.4 | 40.6 | 100.0 | | |

The six kinds of larvae included in this report constituted over 59 percent of the fish larvae obtained in plankton hauls during 1957 (text table 2). As in all recent seasons, anchovy larvae were the most abundant kind in the survey area, and hake larvae were second in abundance. The abundance ranking of the other four kinds of larvae were as follows: fourth, rockfish larvae (Sebastes spp.); sixth, jack mackerel larvae; eleventh, sardine larvae; and eighteenth, Pacific mackerel larvae.

The abundances (standard haul totals for the year)^{1/} of the twelve most common kinds of larvae obtained in 1957 are compared with their contributions in 1956 and 1955;

| Larvae | 1957 | | 1956 | | 1955 | |
|--|---------|------|---------|------|---------|------|
| | Number | Rank | Number | Rank | Number | Rank |
| Anchovy (<u>Engraulis mordax</u>) | 146,631 | 1 | 134,931 | 1 | 140,183 | 1 |
| Hake (<u>Merluccius productus</u>) | 78,283 | 2 | 94,277 | 2 | 60,090 | 2 |
| <u>Vinciguerria lucetia</u> | 55,114 | 3 | 9,832 | 9 | 12,654 | 9 |
| Rockfish (<u>Sebastes</u> spp.) | 36,473 | 4 | 29,144 | 3 | 29,341 | 3 |
| <u>Leuroglossus stilbius</u> | 29,506 | 5 | 18,620 | 5 | 15,111 | 5 |
| Jack mackerel (<u>Trachurus symmetricus</u>) | 20,006 | 6 | 8,027 | 10 | 13,246 | 7 |
| <u>Lampanyctus leucopsarus</u> | 16,808 | 7 | 15,125 | 7 | 7,454 | 10 |
| <u>Lampanyctus mexicanus</u> | 16,207 | 8 | 10,802 | 8 | 13,165 | 8 |
| <u>Citharichthys</u> spp. | 15,813 | 9 | 23,635 | 4 | 20,411 | 4 |
| <u>Diogenichthys laternatus</u> | 11,603 | 10 | 3,158 | 13 | 4,771 | 11 |
| Sardine (<u>Sardinops caerulea</u>) | 9,833 | 11 | 15,523 | 6 | 14,121 | 6 |
| <u>Bathylagus wesethi</u> | 6,347 | 12 | 2,146 | 17 | 3,245 | 12 |
| All others | 50,925 | | 42,920 | | 25,363 | |
| Total | 493,549 | | 408,140 | | 359,155 | |

Monthly abundances of the larvae of sardine, anchovy, jack mackerel, Pacific mackerel, hake, and rockfish are summarized in text table 2.

^{1/} The method of deriving standard haul counts, totals and factors has been described by Ahlstrom (1948).

RECORD OF STANDARDIZED HAUL FACTORS FOR OBLIQUE HAULS
MADE WITH PLANKTON NETS DURING CRUISES 5701-5712

Standardized haul factors are given for all plankton hauls taken in the regular CCOFI survey area during 1957, in table I. Hauls made in the Gulf of California are not included. Additional information concerning each haul, including position of occupancy, date and time of collection, volume of water strained, average depth of haul in meters, and volume of plankton obtained is given in Zooplankton volumes off the Pacific coast, 1957 (Thraillkill: in press).

The following symbols are used in table I:

(-) - a dash indicates that the station was not occupied on the cruise under which it appears.

NS - station occupied, but sample subsequently spoiled, broken or lost.

Table I
Record of standardized haul factors for oblique hauls made
with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|-------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 60.50 | - | - | - | - | 4.59 | 1.70 | NQ | - | - | - | - | - |
| .55 | - | - | - | - | 4.85 | 3.03 | 3.38 | - | - | - | - | - |
| .60 | - | - | - | - | 4.73 | 3.12 | 3.25 | - | - | - | - | - |
| .70 | - | - | - | - | 4.52 | 2.84 | 4.51 | - | - | - | - | - |
| .80 | - | - | - | - | 4.18 | 2.91 | - | - | - | - | - | - |
| .90 | - | - | - | - | 4.35 | 4.02 | - | - | - | - | - | - |
| 63.52 | - | - | - | - | 4.36 | 3.72 | 2.64 | - | - | - | - | - |
| .55 | - | - | - | - | 4.23 | 3.64 | 2.53 | - | - | - | - | - |
| .60 | - | - | - | - | 3.32 | 3.24 | 3.57 | - | - | - | - | - |
| .70 | - | - | - | - | 3.61 | 3.35 | 3.76 | - | - | - | - | - |
| .80 | - | - | - | - | 3.87 | 3.65 | 2.26 | - | - | - | - | - |
| .90 | - | - | - | - | - | 3.60 | - | - | - | - | - | - |
| 67.50 | - | - | - | - | 2.87 | 3.28 | NS | - | - | - | - | - |
| .55 | - | - | - | - | 3.44 | 3.19 | 3.18 | - | - | - | - | - |
| .60 | - | - | - | - | 3.52 | 3.50 | 3.34 | - | - | - | - | - |
| .70 | - | - | - | - | 4.12 | 3.14 | 3.99 | - | - | - | - | - |
| .80 | - | - | - | - | 4.00 | 3.19 | 3.23 | - | - | - | - | - |
| .90 | - | - | - | - | - | 3.16 | - | - | - | - | - | - |
| 70.50 | - | - | - | - | - | - | - | - | - | - | - | - |
| .52 | - | - | - | - | 3.52 | 3.71 | 3.12 | - | - | - | - | - |
| .55 | - | - | - | - | 4.33 | 4.05 | 3.43 | - | - | - | 2.67 | - |
| .60 | - | - | - | - | 4.30 | 3.13 | 3.54 | - | - | - | 3.06 | - |
| .70 | - | - | - | - | 4.45 | 2.98 | 2.10 | - | - | - | 2.45 | - |
| .80 | - | - | - | - | 3.47 | 3.19 | - | - | - | - | 3.11 | - |
| .90 | - | - | - | - | 3.29 | 2.75 | - | - | - | - | - | - |
| 73.50 | - | - | - | - | 3.03 | 3.55 | 2.38 | - | - | - | 3.39 | - |
| .55 | - | - | - | - | 3.11 | 3.86 | 3.42 | - | - | - | 3.17 | - |
| .60 | - | - | - | - | 4.56 | 3.02 | 3.18 | - | - | - | 3.10 | - |
| .70 | - | - | - | - | 4.13 | 3.10 | 3.72 | - | - | - | - | - |
| .80 | - | - | - | - | 3.91 | 3.08 | - | - | - | - | - | - |
| .90 | - | - | - | - | - | 3.44 | - | - | - | - | - | - |
| 77.50 | - | - | - | - | 3.30 | 3.06 | 3.30 | - | - | - | - | - |
| .55 | - | - | - | - | 3.76 | 2.99 | 3.28 | - | - | - | - | - |
| .60 | - | - | - | - | 4.02 | 3.20 | 3.58 | - | - | - | - | - |
| .70 | - | - | - | - | 3.71 | 2.88 | 3.77 | - | - | - | - | - |
| .80 | - | - | - | - | 3.69 | 2.83 | 3.65 | - | - | - | - | - |
| .90 | - | - | - | - | - | 3.11 | 3.04 | - | - | - | - | - |

Table I (cont'd)
 Record of standardized haul factors for oblique hauls made
 with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|-------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 80.51 | - | 2.47 | 3.05 | 2.87 | 2.54 | 3.18 | 2.61 | - | - | 2.75 | - | 2.53 |
| .55 | - | 2.97 | 2.95 | 1.86 | 2.69 | 3.44 | 2.83 | - | - | 2.95 | 3.06 | 3.00 |
| .60 | - | 3.38 | 3.02 | 2.42 | 2.94 | 2.76 | 3.22 | - | - | 2.73 | 2.94 | 3.67 |
| .70 | - | 2.73 | 2.76 | 3.23 | 3.57 | 3.29 | 3.73 | - | - | 2.90 | 2.81 | 3.32 |
| .80 | - | 2.95 | 2.90 | 2.29 | 3.24 | 3.18 | 2.59 | - | - | NS | - | 3.01 |
| .90 | - | 3.06 | 2.96 | 2.89 | 3.65 | 3.02 | 2.72 | - | - | 2.86 | - | 3.21 |
| 82.47 | - | 2.67 | 2.75 | 2.70 | 2.33 | 3.28 | 2.64 | - | - | 3.49 | 4.68 | - |
| 83.40 | - | 1.45 | 2.05 | 1.97 | 2.82 | 1.80 | 1.16 | - | - | 1.30 | 2.26 | - |
| .43 | - | 2.99 | 2.83 | 1.65 | 2.52 | 2.62 | 1.07 | - | - | 3.55 | 5.65 | - |
| .48 | - | - | 2.52 | - | - | - | - | - | - | - | - | - |
| .51 | - | 4.52 | - | 2.41 | 3.45 | 3.55 | 2.43 | - | - | 3.05 | 3.32 | - |
| .52 | - | 2.57 | 2.70 | - | - | - | - | - | - | - | - | - |
| .55 | - | - | - | - | 3.40 | 3.37 | 2.85 | - | - | - | 2.30 | - |
| .60 | - | 2.34 | 2.94 | 1.86 | 2.89 | 3.43 | 2.68 | - | - | 3.51 | 2.50 | - |
| .65 | - | - | - | - | - | 3.15 | 3.28 | - | - | - | - | - |
| .70 | - | - | 2.87 | 2.47 | 3.04 | 3.00 | 2.89 | - | - | 2.35 | - | - |
| .75 | - | - | - | - | - | 2.60 | 3.55 | - | - | - | - | - |
| .80 | - | - | - | 2.18 | 2.85 | 3.23 | 3.94 | - | - | 2.64 | - | - |
| .85 | - | - | - | - | - | 2.93 | 2.78 | - | - | - | - | - |
| .90 | - | - | - | 2.35 | 3.28 | 2.95 | NS | - | - | 2.63 | - | 3.01 |
| 86.46 | - | 4.09 | - | 2.46 | - | - | - | - | - | - | - | - |
| 87.35 | - | 2.67 | 3.01 | 2.07 | 2.96 | 3.38 | 2.58 | - | - | 2.86 | 2.25 | - |
| .38 | - | 3.03 | - | - | - | - | - | - | - | - | - | - |
| .40 | - | 3.06 | 2.76 | 2.43 | 2.19 | 4.04 | NS | - | - | 4.17 | 3.87 | - |
| .45 | - | - | 3.12 | - | 3.48 | 4.08 | 2.69 | - | - | 3.09 | 3.49 | - |
| .50 | - | 2.97 | 2.92 | 3.44 | 1.83 | 2.89 | 2.37 | - | - | 3.18 | 3.23 | - |
| .55 | - | 2.54 | - | - | 3.25 | 2.84 | 2.86 | - | - | 2.39 | 2.59 | - |
| .60 | - | 2.82 | 2.93 | 2.98 | 2.73 | 3.82 | 2.72 | - | - | 3.35 | 3.83 | - |
| .65 | - | - | - | - | 2.89 | 2.77 | 3.40 | - | - | - | - | - |
| .70 | - | 2.82 | 2.94 | 2.84 | 3.26 | 3.24 | 4.69 | - | - | 2.96 | - | - |
| .75 | - | - | - | - | 3.35 | 2.95 | 3.82 | - | - | - | - | - |
| .80 | - | - | - | 3.19 | 3.09 | 3.13 | NS | - | - | 3.21 | - | - |
| .85 | - | - | - | - | 2.78 | 2.68 | 3.51 | - | - | - | - | - |
| .90 | - | - | - | 2.52 | 3.45 | 3.01 | 3.49 | - | - | 2.74 | - | 3.15 |
| 90.28 | - | 2.74 | 2.25 | 2.37 | 2.68 | 3.97 | 2.00 | - | - | 3.27 | 5.53 | 3.30 |
| .30 | - | 3.13 | 3.33 | 1.96 | 1.89 | 3.94 | 2.62 | - | - | 3.36 | 2.94 | 3.56 |
| .37 | - | 2.62 | 3.01 | 3.10 | 3.17 | 3.19 | 2.43 | - | - | 3.35 | 1.93 | 2.99 |
| .45 | - | 3.21 | 2.92 | 3.29 | 2.71 | 3.58 | 2.35 | - | - | 1.34 | 2.92 | 3.22 |

Table I (cont'd)
 Record of standardized haul factors for oblique hauls made
 with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|-------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 90.50 | - | - | - | - | 2.77 | 3.59 | 2.09 | - | - | 1.36 | 2.30 | 3.86 |
| .55 | - | 2.96 | 3.07 | 3.39 | 2.89 | 3.50 | 1.18 | - | - | 3.26 | 3.03 | 3.49 |
| .60 | - | 2.91 | 2.85 | 2.93 | 3.42 | 3.15 | 2.34 | - | - | 3.35 | 3.37 | 3.28 |
| .65 | - | - | - | - | 2.35 | 3.11 | 2.61 | - | - | - | - | - |
| .70 | - | 3.08 | 2.80 | 2.90 | 3.30 | 3.14 | 2.74 | - | - | 3.15 | - | 3.09 |
| .75 | - | - | - | - | 3.27 | 3.75 | 2.41 | - | - | - | - | - |
| .80 | - | 2.89 | 2.86 | 3.10 | 2.57 | 3.27 | 2.57 | - | - | 1.34 | - | 2.79 |
| .85 | - | - | - | - | 2.46 | 3.27 | 3.17 | - | - | - | - | - |
| .90 | - | 2.81 | 3.16 | 3.31 | 3.02 | 3.35 | 2.64 | - | - | 3.11 | - | 3.27 |
| 93.27 | - | 2.87 | 2.96 | 3.14 | 2.25 | 3.70 | 2.12 | - | - | 2.85 | 2.80 | - |
| .30 | - | 2.67 | 2.97 | 3.41 | 2.03 | 2.86 | 2.70 | - | - | 3.54 | 3.05 | - |
| .35 | - | - | - | 3.67 | 2.08 | 3.08 | 2.61 | - | - | 4.47 | 2.32 | - |
| .40 | - | 2.77 | 3.23 | 2.50 | 2.69 | 3.09 | 2.92 | - | - | 2.83 | 2.66 | - |
| .45 | - | - | - | 3.09 | 2.63 | 3.14 | 2.22 | - | - | - | 3.09 | - |
| .50 | - | 3.09 | 3.01 | 3.73 | 4.55 | 3.39 | 2.82 | - | - | 3.40 | 2.97 | - |
| .55 | - | - | - | 2.40 | 2.90 | 3.28 | 3.00 | - | - | - | 2.54 | - |
| .60 | - | 3.36 | 3.24 | 2.85 | 3.09 | 3.70 | 2.33 | - | - | 3.12 | 4.05 | - |
| .65 | - | - | - | - | 2.56 | 3.67 | 2.23 | - | - | - | - | - |
| .70 | - | 3.15 | 3.00 | 2.74 | 2.71 | 3.97 | 3.84 | - | - | 3.07 | - | - |
| .75 | - | - | - | - | 2.68 | 3.48 | 3.58 | - | - | - | - | - |
| .80 | - | - | - | NS | 2.70 | 3.52 | 2.59 | - | - | 3.17 | - | - |
| .85 | - | - | - | - | 3.45 | 3.56 | 2.85 | - | - | - | - | - |
| .90 | - | - | - | 1.59 | 2.99 | 3.84 | 2.46 | - | - | 3.44 | - | - |
| 97.30 | - | 2.67 | 2.53 | 1.53 | 2.23 | 3.26 | 1.99 | - | - | 2.12 | 4.21 | - |
| .32 | - | 3.29 | - | 2.83 | 3.19 | 3.01 | 2.18 | - | - | 3.76 | 3.91 | 3.16 |
| .35 | - | - | - | - | - | - | - | - | - | 3.49 | - | - |
| .40 | - | 3.09 | 3.10 | 3.01 | 1.95 | 3.01 | 2.29 | - | - | 3.40 | 5.08 | - |
| .45 | - | - | - | 3.07 | 2.60 | 4.03 | 2.73 | - | - | - | 3.70 | - |
| .50 | - | 3.04 | 2.85 | NS | 2.87 | 3.77 | 2.66 | - | - | 3.51 | 1.96 | - |
| .55 | - | - | - | NS | 3.24 | 3.29 | 2.86 | - | - | - | 2.65 | - |
| .60 | - | 3.24 | - | 1.60 | 3.92 | 3.05 | 2.43 | - | - | 3.18 | 5.18 | - |
| .65 | - | - | - | - | 2.90 | 3.54 | 2.58 | - | - | - | - | - |
| .70 | - | 2.92 | - | 2.84 | 2.37 | 3.22 | 2.95 | - | - | 2.09 | - | - |
| .75 | - | - | - | - | 3.35 | 3.11 | 2.85 | - | - | - | - | - |
| .80 | - | - | - | 2.35 | 3.48 | 2.79 | 2.77 | - | - | 3.33 | - | - |
| .85 | - | - | - | - | 2.98 | 3.03 | 2.82 | - | - | - | - | - |
| .90 | - | - | - | 3.83 | 1.89 | 3.04 | 2.51 | - | - | 3.10 | - | - |

Table I (cont'd)
 Record of standardized haul factors for oblique hauls made
 with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|--------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 100.29 | - | 2.98 | 3.49 | 2.62 | 2.26 | 2.97 | 1.89 | - | - | 3.15 | - | - |
| .30 | - | - | - | - | - | - | - | - | - | 3.21 | - | - |
| .33 | - | 2.80 | 3.42 | 2.61 | 2.13 | 2.99 | 2.82 | - | - | - | - | - |
| .35 | - | - | - | - | - | - | - | - | - | 3.28 | - | - |
| .40 | - | 2.80 | 3.06 | 2.45 | 3.70 | 3.90 | 2.65 | - | - | 2.85 | - | 2.95 |
| .45 | - | - | - | 3.15 | 2.83 | 3.04 | 2.86 | - | - | - | - | - |
| .50 | - | 2.78 | 3.18 | 3.14 | 2.40 | 2.99 | 2.79 | - | - | 3.00 | - | - |
| .55 | - | - | - | 3.44 | 2.28 | 3.31 | 2.60 | - | - | - | - | - |
| .60 | 2.97 | 3.41 | 3.00 | 2.42 | 2.98 | 3.10 | 2.72 | - | - | 3.17 | - | - |
| .65 | - | - | - | - | 3.01 | 2.70 | 2.62 | - | - | - | - | - |
| .70 | - | 3.62 | - | 1.76 | 3.42 | 2.80 | - | - | - | 2.84 | - | - |
| .75 | - | - | - | - | 2.28 | 3.04 | 3.17 | - | - | - | - | - |
| .80 | - | 3.43 | - | 3.41 | 2.17 | 3.52 | - | - | - | 2.99 | - | - |
| .85 | - | - | - | - | 3.42 | 3.09 | - | - | - | - | - | - |
| .90 | - | 3.70 | - | 2.66 | 3.20 | 2.81 | - | - | - | 3.09 | - | - |
| 103.30 | 2.45 | 3.40 | 2.93 | 1.43 | 2.26 | 2.77 | 1.63 | - | - | 2.62 | - | - |
| .35 | 2.91 | 3.46 | 2.90 | 2.65 | 3.28 | 2.93 | 3.15 | - | - | 3.34 | - | - |
| .38 | 2.91 | 3.28 | - | - | - | - | - | - | - | - | - | - |
| .40 | 3.06 | 3.14 | 3.96 | 2.82 | 3.44 | 3.82 | 2.90 | - | - | 3.15 | - | - |
| .45 | - | - | - | 3.08 | 2.56 | 3.87 | 2.83 | - | - | - | - | - |
| .50 | 3.03 | 3.59 | 3.11 | 2.68 | 2.60 | 3.35 | 2.47 | - | - | 3.41 | - | 3.20 |
| .55 | - | - | - | 2.80 | 3.57 | 3.27 | 2.56 | - | - | - | - | - |
| .60 | 3.16 | 3.04 | NS | 3.41 | 3.47 | 3.36 | 2.67 | - | - | 3.20 | - | - |
| .65 | - | - | - | - | 2.65 | 3.69 | 2.65 | - | - | - | - | - |
| .70 | - | 3.23 | 3.10 | 3.20 | 3.21 | 3.29 | 3.07 | - | - | 3.34 | - | - |
| .75 | - | - | - | - | 2.93 | 3.42 | 2.63 | - | - | - | - | - |
| .80 | - | - | - | 3.01 | 3.43 | 3.28 | - | - | - | 3.05 | - | - |
| .85 | - | - | - | - | 2.42 | 3.31 | - | - | - | - | - | - |
| .90 | - | - | - | 3.13 | 2.70 | 3.00 | - | - | - | - | - | - |
| 107.32 | 2.91 | 1.49* | 3.32 | 3.19 | 2.31 | - | 3.67 | - | - | 2.89 | - | - |
| .35 | 2.90 | - | 3.21 | 3.25 | 2.10 | 3.40 | 3.49 | - | - | 2.60 | - | - |
| .40 | 2.86 | 3.14 | 3.06 | 3.33 | 2.25 | 3.42 | 2.30 | - | - | 3.63 | - | - |
| .45 | - | - | - | 2.84 | 3.02 | 3.35 | 3.21 | - | - | - | - | - |
| .50 | 2.65 | 2.72 | 3.01 | 3.42 | 2.49 | 2.93 | 3.33 | - | - | 4.34 | - | - |
| .55 | - | - | - | 3.18 | 2.14 | 3.13 | 3.23 | - | - | - | - | - |
| .60 | 3.16 | 2.82 | 3.80 | 3.50 | 2.12 | 3.39 | 3.40 | - | - | 3.73 | - | 3.71 |
| .65 | - | - | - | - | 3.02 | 3.27 | 3.36 | - | - | - | - | - |
| .70 | - | 2.32 | 3.11 | 2.17 | 2.36 | 3.21 | 3.38 | - | - | 3.25 | - | - |

* - Samples combined. Values adjusted accordingly.

Table I (cont'd)
 Record of standardized haul factors for oblique hauls made
 with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|--------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 107.75 | - | - | - | - | 2.12 | 3.08 | 3.25 | - | - | - | - | - |
| .80 | - | - | - | 3.32 | 3.19 | 3.56 | 3.47 | - | - | 3.21 | - | - |
| .85 | - | - | - | - | 3.56 | 3.19 | 3.47 | - | - | - | - | - |
| .90 | - | - | - | 2.80 | 2.89 | 3.20 | 3.41 | - | - | - | - | - |
| 110.33 | - | 2.61 | 4.50 | 2.68 | 2.76 | 3.19 | 3.39 | 2.82 | 2.42 | 3.05 | - | - |
| .35 | - | 3.09 | 3.51 | 3.14 | 2.96 | 3.11 | 2.96 | 2.90 | 3.54 | 2.45 | - | - |
| .40 | 2.88 | 2.79 | 3.29 | 3.53 | 2.92 | 3.07 | 2.98 | 4.54 | 2.86 | 2.79 | - | - |
| .45 | - | - | - | 2.99 | 3.07 | 3.03 | 3.24 | - | - | - | - | - |
| .50 | 2.73 | 2.66 | 3.74 | 2.86 | 3.08 | 2.97 | 3.29 | - | - | 2.84 | - | - |
| .55 | - | - | - | 2.96 | 3.16 | 2.61 | 3.30 | - | - | - | - | - |
| .60 | 2.89 | 2.55 | 2.78 | 3.87 | 3.05 | 2.98 | 3.15 | - | - | 3.55 | - | - |
| .65 | - | - | - | - | 2.66 | 2.53 | 4.33 | - | - | - | - | - |
| .70 | - | 2.96 | 3.10 | 3.30 | 2.65 | 3.50 | 3.42 | - | - | 3.30 | - | 3.18 |
| .75 | - | - | - | - | 2.89 | 3.24 | 3.70 | - | - | - | - | - |
| .80 | - | 2.54 | - | 3.34 | 3.02 | 2.95 | 3.43 | - | - | 2.78 | - | - |
| .85 | - | - | - | - | 2.78 | 2.14 | 3.32 | - | - | - | - | - |
| .90 | - | 2.89 | - | 3.35 | 3.04 | 2.38 | 3.42 | - | - | - | - | - |
| 113.30 | 2.49 | 2.56 | 2.91 | 1.72 | 2.32 | 1.79 | 2.07 | 2.70 | 2.37 | 2.79 | - | - |
| .35 | 2.72 | 2.50 | 3.17 | 3.95 | 3.10 | 2.69 | 3.95 | 2.66 | 3.13 | 3.24 | - | - |
| .40 | 2.57 | 2.52 | 3.50 | 3.02 | 3.05 | 3.07 | 3.20 | 2.99 | 2.90 | 3.11 | - | - |
| .45 | - | - | - | 2.83 | 2.74 | 3.22 | 3.33 | - | - | - | - | - |
| .50 | 2.79 | 2.01 | 3.31 | 2.09 | 2.65 | 2.46 | 3.87 | - | - | 2.40 | - | - |
| .55 | - | - | - | 3.18 | 2.92 | 2.88 | 3.50 | - | - | - | - | - |
| .60 | 2.76 | 2.16 | 3.15 | 3.55 | 2.94 | 2.88 | 3.67 | - | - | 2.61 | - | - |
| .65 | - | - | - | - | 2.71 | 2.39 | 3.43 | - | - | - | - | - |
| .70 | 2.63 | 2.61 | 3.05 | 2.53 | 2.59 | 2.52 | 3.50 | - | - | 3.02 | - | - |
| .75 | - | - | - | - | 2.56 | 2.54 | 3.33 | - | - | - | - | - |
| .80 | - | - | - | 2.93 | 2.96 | 2.52 | 3.62 | - | - | 2.74 | - | - |
| .85 | - | - | - | - | - | 2.93 | - | - | - | - | - | - |
| .90 | - | - | - | - | - | 3.03 | - | - | - | - | - | - |
| 115.27 | - | - | - | - | - | - | - | 2.89 | 2.66 | - | - | - |
| .30 | - | - | - | - | - | - | - | 3.38 | 3.09 | - | - | - |
| .35 | - | - | - | - | - | - | - | 2.81 | 2.92 | - | - | - |
| .40 | - | - | - | - | - | - | - | 3.07 | 3.21 | - | - | - |
| 117.26 | 2.58 | 2.36 | 3.28 | 2.83 | 1.67 | 1.50 | 3.40 | 2.92 | 2.65 | 3.01 | - | - |
| .30 | 2.64 | 2.09 | 3.08 | 3.84 | 2.33 | 3.39 | 3.64 | 2.93 | 3.10 | 2.32 | - | - |
| .35 | 2.67 | 1.54 | 3.45 | 1.89 | 2.07 | 2.45 | 3.79 | 3.20 | 3.00 | 2.40 | - | - |
| .40 | 2.67 | 2.31 | 2.98 | 2.60 | 2.72 | 1.36 | 3.35 | 2.74 | 3.30 | 2.90 | - | - |

Table I (cont'd)
 Record of standardized haul factors for oblique hauls made
 with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|---------------------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 117.45 | - | - | - | 2.85 | 2.29 | 1.28 | 3.74 | - | - | - | - | - |
| .50 | 2.64 | 2.81 | 3.12 | 2.71 | 2.80 | 3.13 | 3.34 | - | - | 3.10 | - | - |
| .55 | - | - | - | 3.11 | 3.04 | 2.58 | 3.01 | - | - | - | - | - |
| .60 | 2.70 | 2.42 | 3.07 | 2.63 | 3.08 | 2.46 | 2.96 | - | - | 2.04 | - | - |
| .65 | - | - | - | - | 2.61 | 2.38 | 3.68 | - | - | - | - | - |
| .70 | 2.77 | 4.03 | 3.21 | 2.67 | 2.27 | 2.46 | 3.30 | - | - | 2.24 | - | - |
| .75 | - | - | - | - | 2.69 | 2.82 | 3.37 | - | - | - | - | - |
| .80 | - | - | - | 2.42 | 2.68 | 3.46 | 3.64 | - | - | 2.46 | - | - |
| .85 | - | - | - | - | - | 3.33 | - | - | - | - | - | - |
| .90 | - | - | - | - | - | 3.21 | - | - | - | - | - | - |
| 118.39 | - | - | 2.39 | 2.15 | 3.02 | 2.41 | 3.29 | - | - | 2.38 | - | - |
| 118 ⁵ 25 | - | - | - | - | - | - | - | 3.02 | 2.54 | - | - | - |
| 118 ⁵ 30 | - | - | - | - | - | - | - | 3.19 | 3.75 | - | - | - |
| 118 ⁵ 35 | - | - | - | - | - | - | - | 3.12 | 3.48 | - | - | - |
| 119.33 | - | - | 3.24 | 3.08 | 2.68 | 2.74 | 3.43 | 2.90 | 2.98 | 2.60 | - | - |
| 120.25 | 2.27 | 2.98 | 2.98 | 2.45 | 3.70 | 2.52 | 2.97 | 2.96 | 3.17 | 2.52 | - | - |
| .30 | 2.46 | 2.93 | 3.01 | 2.14 | 2.79 | 3.30 | 3.23 | 3.34 | 3.09 | 2.11 | - | - |
| .35 | 2.33 | 2.43 | 2.27 | 1.90 | 3.00 | 1.71 | 2.87 | 2.13 | 2.99 | 2.27 | - | - |
| .40 | 1.25 | 2.35 | 1.87 | - | 1.52 | 1.79 | 1.89 | 2.03 | 2.95 | 1.67 | - | - |
| .45 | 2.78 | 2.77 | - | - | 2.53 | 3.21 | 3.85 | 2.93 | 2.82 | 2.80 | - | - |
| .50 | 2.72 | 2.76 | - | 2.89 | 2.60 | 2.53 | 3.65 | - | - | 2.72 | - | - |
| .55 | 2.79 | 2.35 | - | 2.65 | 2.79 | 2.26 | 3.34 | - | - | - | - | - |
| .60 | 2.64 | 3.16 | 2.86 | 3.35 | 2.91 | 3.68 | 3.52 | - | - | 3.01 | - | - |
| .65 | - | - | - | - | 3.03 | 3.43 | 3.54 | - | - | - | - | - |
| .70 | 2.95 | 3.01 | 3.73 | 2.45 | 2.94 | 2.87 | 3.36 | - | - | 3.26 | - | - |
| .75 | - | - | - | - | 2.50 | 3.49 | 3.40 | - | - | - | - | - |
| .80 | - | 2.45 | 3.25 | 3.05 | 2.51 | 2.31 | 3.48 | - | - | 2.93 | - | - |
| .85 | - | - | - | - | - | 2.82 | - | - | - | - | - | - |
| .90 | - | - | - | - | - | 3.05 | - | - | - | 2.97 | - | - |
| 123.37 | 2.46 | 1.70 | 3.24 | 2.97 | 1.96 | 2.16 | 2.90 | 2.43 | 3.37 | 2.72 | - | - |
| .42 | 2.77 | 2.67 | 2.99 | 2.88 | 2.70 | 3.08 | 3.13 | 3.30 | 3.17 | 2.93 | - | - |
| .45 | - | - | - | - | - | - | - | 3.03 | 3.23 | - | - | - |
| .50 | 2.84 | 2.38 | 3.00 | 2.49 | - | 3.17 | 2.91 | - | - | 2.54 | - | - |
| .55 | - | 2.27 | 2.89 | 2.59 | - | 2.73 | 3.09 | - | - | - | - | - |
| .60 | 2.72 | 2.22 | 2.83 | 3.05 | - | 2.98 | 3.09 | - | - | 3.06 | - | - |
| .70 | - | - | - | 2.07 | - | - | 3.30 | - | - | 3.09 | - | - |
| .80 | - | - | - | 2.46 | - | - | 3.00 | - | - | 2.95 | - | - |

Table I (cont'd)
 Record of standardized haul factors for oblique hauls made
 with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|--------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 127.34 | 2.44 | 2.63 | 3.05 | 1.63 | 2.91 | 2.18 | 2.15 | 2.88 | 3.37 | 3.39 | - | - |
| .40 | 2.73 | 3.52 | 3.75 | 2.73 | 2.73 | 2.77 | 3.05 | 3.36 | 3.46 | 3.78 | - | - |
| .45 | - | 2.51 | 2.74 | 3.04 | 3.57 | 2.77 | 3.24 | 2.85 | 3.78 | 3.82 | - | - |
| .50 | 2.63 | 2.64 | 3.15 | 2.64 | - | 2.88 | 3.12 | - | - | 4.34 | - | - |
| .55 | - | 3.46 | 3.37 | 2.98 | - | 2.32 | 3.23 | - | - | - | - | - |
| .60 | 2.82 | 2.67 | 3.41 | 3.01 | - | 2.78 | 2.90 | - | - | 3.25 | - | - |
| .70 | - | - | - | 3.05 | - | - | 3.53 | - | - | 3.08 | - | - |
| .80 | - | - | - | 2.74 | - | - | 3.17 | - | - | 3.61 | - | - |
| 130.30 | 2.39 | 1.60 | 2.67 | 1.74 | 2.94 | 2.59 | 2.59 | 2.68 | 2.57 | 3.35 | - | - |
| .35 | 2.64 | 2.18 | 3.11 | 2.10 | 2.42 | 3.10 | 2.76 | 2.97 | 3.13 | 4.28 | - | - |
| .40 | 2.89 | 2.61 | 2.69 | 2.83 | 2.19 | 3.13 | 3.30 | 2.75 | 3.30 | 5.25 | - | - |
| .45 | - | - | - | 2.97 | - | 2.98 | 2.95 | 2.70 | 4.54 | - | - | - |
| .50 | 2.83 | 2.98 | 3.41 | 3.06 | 2.77 | 3.33 | 3.08 | - | - | 3.31 | - | - |
| .55 | - | - | - | 3.28 | - | 2.97 | 2.79 | - | - | - | - | - |
| .60 | 2.87 | 2.62 | 3.05 | 3.07 | 2.74 | 3.04 | 3.48 | - | - | 3.41 | - | - |
| .70 | - | - | - | 2.51 | - | - | 2.95 | - | - | 2.99 | - | - |
| .80 | - | - | - | 2.87 | - | - | 3.63 | - | - | 3.87 | - | - |
| 133.25 | 2.63 | 1.38 | 2.56 | 2.52 | 2.33 | 3.20 | 2.72 | 3.24 | 3.01 | 3.02 | - | - |
| .30 | 2.64 | 2.37 | 2.73 | 2.42 | 2.32 | 3.41 | 3.33 | 3.46 | 3.53 | 2.73 | - | - |
| .35 | 2.81 | - | 3.22 | 2.98 | - | 2.38 | 3.42 | - | - | 3.30 | - | - |
| .40 | 2.67 | 2.11 | 2.87 | 2.97 | 2.65 | 2.66 | 3.43 | - | - | 3.05 | - | - |
| .45 | - | - | - | 2.78 | - | 2.10 | 3.61 | - | - | - | - | - |
| .50 | - | 3.18 | 3.24 | 3.17 | 3.14 | 3.36 | 3.38 | - | - | 3.09 | - | - |
| .55 | - | - | - | 2.89 | - | - | - | - | - | - | - | - |
| .60 | - | - | - | 2.19 | - | - | 3.12 | - | - | 2.93 | - | - |
| .70 | - | - | - | 2.90 | - | - | 3.20 | - | - | 2.82 | - | - |
| .80 | - | - | - | 2.15 | - | - | 3.13 | - | - | 4.07 | - | - |
| 137.23 | 2.43 | 2.13 | 2.86 | 1.65 | 2.81 | 2.62 | 2.71 | 2.67 | 3.22 | 3.63 | - | - |
| .30 | 2.85 | 2.49 | 2.79 | 3.30 | 2.79 | 2.63 | 3.45 | 3.03 | 3.29 | 3.71 | - | - |
| .35 | 2.67 | 2.46 | 2.92 | - | - | 2.45 | 3.56 | - | - | 3.65 | - | - |
| .40 | - | 2.20 | 2.91 | 2.38 | NS | 2.53 | 3.74 | - | - | 3.15 | - | - |
| .45 | - | - | - | 2.39 | - | 2.09 | 3.30 | - | - | - | - | - |
| .50 | - | 2.38 | 3.15 | 2.93 | 2.42 | 3.25 | 3.30 | - | - | 2.92 | - | - |
| .55 | - | - | - | 2.89 | - | - | - | - | - | - | - | - |
| .60 | - | - | - | 2.77 | - | - | 3.06 | - | - | 3.09 | - | - |
| .70 | - | - | - | 3.14 | - | - | 3.16 | - | - | 3.25 | - | - |
| .80 | - | - | - | 3.20 | - | - | 2.86 | - | - | 3.14 | - | - |

Table I (cont'd)
 Record of standardized haul factors for oblique hauls made
 with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|--------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 140.30 | - | 2.05 | 2.70 | 2.35 | - | 2.17 | - | - | - | - | - | - |
| .35 | - | 1.83 | 2.75 | 2.53 | - | 3.34 | - | - | - | - | - | - |
| .40 | - | 2.85 | 3.18 | 2.04 | - | 2.78 | - | - | - | - | - | - |
| .45 | - | - | - | 2.28 | - | - | - | - | - | - | - | - |
| .50 | - | 3.37 | 2.90 | 2.51 | - | 3.83 | - | - | - | - | - | - |
| .55 | - | - | - | 2.56 | - | - | - | - | - | - | - | - |
| .60 | - | - | - | 2.99 | - | - | - | - | - | - | - | - |
| .70 | - | - | - | 2.86 | - | - | - | - | - | - | - | - |
| .80 | - | - | - | 2.48 | - | - | - | - | - | - | - | - |
| 143.26 | - | 2.07 | 2.94 | 2.61 | - | - | - | - | - | - | - | - |
| .30 | - | 2.39 | 3.31 | 2.48 | - | 4.31 | - | - | - | - | - | - |
| .35 | - | 2.08 | 2.44 | 2.52 | - | 3.89 | - | - | - | - | - | - |
| .40 | - | 2.86 | 3.07 | 2.86 | - | 3.77 | - | - | - | - | - | - |
| .50 | - | 3.01 | 3.13 | - | - | 3.52 | - | - | - | - | - | - |
| 147.20 | - | 3.09 | 3.13 | 2.99 | - | - | - | - | - | - | - | - |
| .25 | - | 3.63 | 2.59 | 2.88 | - | - | - | - | - | - | - | - |
| .30 | - | 2.87 | 2.90 | 2.98 | - | - | - | - | - | - | - | - |
| .35 | - | - | - | 2.85 | - | - | - | - | - | - | - | - |
| .40 | - | 2.26 | 3.01 | 2.46 | - | - | - | - | - | - | - | - |
| .45 | - | - | - | 2.59 | - | - | - | - | - | - | - | - |
| .50 | - | - | - | 3.02 | - | - | - | - | - | - | - | - |
| .55 | - | - | - | 3.41 | - | - | - | - | - | - | - | - |
| .60 | - | - | - | 2.41 | - | - | - | - | - | - | - | - |
| .70 | - | - | - | 2.80 | - | - | - | - | - | - | - | - |
| .80 | - | - | - | 2.89 | - | - | - | - | - | - | - | - |
| 148.20 | - | - | - | - | - | 2.91 | - | - | - | - | - | - |
| .25 | - | - | - | - | - | 3.65 | - | - | - | - | - | - |
| .30 | - | - | - | - | - | 3.47 | - | - | - | - | - | - |
| .40 | - | - | - | - | - | 3.03 | - | - | - | - | - | - |
| .50 | - | - | - | - | - | 3.32 | - | - | - | - | - | - |
| 150.16 | - | - | - | 2.89 | - | - | - | - | - | - | - | - |
| .19 | - | 2.07 | 2.31 | - | - | - | - | - | - | - | - | - |
| .20 | - | - | - | 2.97 | - | - | - | - | - | - | - | - |
| .25 | - | 3.08 | 2.81 | 2.69 | - | - | - | - | - | - | - | - |
| .30 | - | - | 2.52 | 2.87 | - | - | - | - | - | - | - | - |
| .35 | - | 3.38 | - | 2.77 | - | - | - | - | - | - | - | - |
| .40 | - | 2.29 | 2.10 | 3.24 | - | - | - | - | - | - | - | - |

Table I (cont'd)
 Record of standardized haul factors for oblique hauls made
 with plankton nets during cruises 5701-5712, 1957

| Sta. | Cruise and month | | | | | | | | | | | |
|--------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 153.16 | - | 1.84 | - | 2.47 | - | 2.98 | - | 2.78 | - | - | - | - |
| .20 | - | 3.40 | - | 3.31 | - | 2.98 | - | 3.25 | - | - | - | - |
| .25 | - | - | - | 2.17 | - | 2.46 | - | - | - | - | - | - |
| .30 | - | 3.01 | - | 3.11 | - | 2.02 | - | - | - | - | - | - |
| .35 | - | - | - | 2.40 | - | - | - | - | - | - | - | - |
| .40 | - | 2.25 | - | 1.92 | - | 2.58 | - | - | - | - | - | - |
| .45 | - | - | - | 2.72 | - | - | - | - | - | - | - | - |
| .50 | - | - | - | 2.99 | - | - | - | - | - | - | - | - |
| .55 | - | - | - | 2.53 | - | - | - | - | - | - | - | - |
| .60 | - | - | - | 2.15 | - | - | - | - | - | - | - | - |
| 157.10 | - | 1.13 | - | - | - | - | - | - | - | - | - | - |
| .15 | - | - | - | 2.79 | - | 3.32 | - | 2.97 | - | - | - | - |
| .20 | - | 3.51 | - | 2.74 | - | 2.78 | - | 2.38 | - | - | - | - |
| .25 | - | - | - | 2.86 | - | 2.77 | - | - | - | - | - | - |
| .30 | - | 2.94 | - | 2.72 | - | 2.90 | - | - | - | - | - | - |
| .35 | - | - | - | 2.69 | - | - | - | - | - | - | - | - |
| .40 | - | - | - | 2.80 | - | 2.57 | - | - | - | - | - | - |

RECORD OF SARDINE EGGS, 1957

A listing of all hauls containing sardine eggs in the 1957 plankton samples obtained on CCOFI cruises is contained in table II. The eggs at each station are reported by age in days (A to D; see below) for two categories, normal and total. Abnormal eggs have embryos that are stunted and misshapen, either due to mechanical injury during collection (rupture of the vitelline membrane) or to a diseased condition of the eggs. Deteriorating eggs are eggs in such poor condition that they cannot be classified with certainty.

The age categories into which sardine eggs are separated are the following:

- A - Eggs spawned within 24 hours of collection
- B - Eggs spawned within 24.1 to 48 hours of collection
- C - Eggs spawned within 48.1 to 72 hours of collection
- D - Eggs spawned within 72.1 to 96 hours of collection
- Unclassified (Uncl.) - deteriorating eggs that cannot be aged with certainty.

A dash (-) in table II indicates that the category was not represented at a station, either actually or potentially.

Considerable publicity has been given to the warming that occurred in the CCOFI area during 1957 and which has continued to the time of writing this report - May 1959 (Progress Report, 1 July 1956 to 1 January 1958, California Cooperative Oceanic Fisheries Investigations, sponsored by the State of California Marine Research Committee, pp. 7-26). The temperatures at which sardine eggs were obtained during 1957 are given by month and area in text table 3. A comparison with average "spawning" temperatures by area for the preceding six years (1951-1956) follows:

| Area | 1957 season | | Average temperature (1951-1956) | Difference |
|--------------------------------|------------------------|---------------------|---------------------------------|------------|
| | Number of observations | Average temperature | | |
| 60-77 (complete season) | 3 | 13.4 | - | - |
| 80-93 (complete season) | 16 | 15.0 | 14.7 | +0.3 |
| 97-107 (complete season) | 15 | 15.4 | 15.1 | +0.3 |
| 110-120 (main spawning period) | 11 | 16.4 | 15.4 | +1.0 |
| (late season spawning) | 16 | 20.8 | 18.1 | +2.7 |
| 123-137 (main spawning period) | 8 | 16.6 | 15.7 | +1.1 |
| (late season spawning) | 6 | 21.0 | 18.0 | +3.0 |
| 140-157 (complete season) | 0 | - | 18.7 | |

In the northern half of the spawning area (lines 80-107) in 1957, sardine eggs were obtained at temperatures which averaged only 0.3°C higher than during the previous six years. In the southern half of the spawning area (lines 110-137), water temperatures at the stations having sardine eggs averaged a full degree higher during the main spawning period (January through June), and 2.7°C to 3.0°C higher during the late spawning period (July through October). Hence, the increased water temperatures during 1957 were reflected mainly in the southern half of the spawning distribution.

The distribution and relative abundance of sardine eggs in 1957 are illustrated in figure 2. The abundance shown for each station represents the cumulative standard haul total of sardine eggs at the station during the year. Stations (open circles) without shading had no sardine eggs in the plankton hauls taken during the year.

Occurrences and abundance (standard haul totals) of sardine eggs are summarized by month and area in text table 4. Sardine eggs were taken in fewer hauls than in any recent year; occurrences during 1957 numbered 76, as compared to 144 in 1956, 186 in 1955, and 309 in 1954. The decrease in occurrences in 1957 reflects, in part at least, a decrease in abundance of spawning sardines. Coverage of spawning areas was as thorough in 1957 as in preceding years, except 1954. (Coverage in 1954 was more intensive than at any time before or since.)

The distribution of spawning between the several subareas, however, has been fairly similar during the past three seasons. This is shown in the following summary:

| Station lines | 1955 | | | 1956 | | | 1957 | | |
|------------------|------------------|--------|---------|------------------|--------|---------|------------------|--------|---------|
| | Occur- rences | Number | Percent | Occur- rences | Number | Percent | Occur- rences | Number | Percent |
| 60-77 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 201 | 0.6 |
| 80-93 | 39 | 6,007 | 13.3 | 29 | 11,739 | 11.9 | 16 | 5,914 | 18.9 |
| 97-107 | 69 | 16,156 | 35.7 | 53 | 34,043 | 34.6 | 15 | 8,883 | 28.4 |
| 110-120 | 60 | 18,763 | 41.4 | 47 | 41,969 | 42.7 | 28 | 12,723 | 40.7 |
| 123-137 | 13 | 3,963 | 8.8 | 13 | 10,532 | 10.7 | 14 | 3,552 | 11.4 |
| 140-157 | 5 | 309 | 0.7 | 2 | 44 | 0.1 | 0 | 0 | 0 |
| Total | 186 | 45,198 | 99.9 | 144 | 98,327 | 100.0 | 76 | 31,273 | 100.0 |

It is convenient to divide sardine spawning into two major centers: one off southern California and northern Baja California (station lines 80-107), the other off central Baja California (station lines 110-137). The two centers have been of about equal importance in each of the above seasons. In 1955, 49 percent of the eggs were obtained in the northern center; in 1956 the percentage was 46 percent; in 1957 it was 48 percent.




Figure 2. --Sardine eggs, 1957: Distribution and relative abundance.

Text table 3. --Temperature ranges and means for stations containing sardine eggs in 1957, summarized by month and area (temperatures at 10 M)

| Southern California (lines 80-93) | | | | Northern Baja California (lines 97-107) | | | |
|-----------------------------------|------------------------|-------------------|------------------|---|------------------------|-------------------|------------------|
| Cruise number | Number of observations | Temperature range | Mean temperature | Cruise number | Number of observations | Temperature range | Mean temperature |
| 5703 | 2 | 12.41-13.24 °C | 12.83 °C | 5702 | 1 | 14.46 | 14.46 °C |
| 5704 | 3 | 13.00-14.76 | 14.06 | 5703 | 5 | 13.50-15.47 | 14.15 |
| 5705 | 6 | 14.35-16.30 | 15.36 | 5704 | 3 | 12.72-15.42 | 14.38 |
| 5706 | 3 | 15.06-17.83 | 16.16 | 5705 | 3 | 15.56-16.74 | 16.09 |
| 5707 | 1 | 15.96 | 15.96 | 5706 | 0 | - | - |
| 5711 | 1 | 15.68 | 15.68 | 5707 | 3 | 16.62-20.11 | 18.02 |
| Total | 16 | 12.41-17.83 | 15.01 | Total | 15 | 12.72-20.11 | 15.38 |

| Upper central Baja California (lines 110-120) | | | | Lower central Baja California (lines 123-137) | | | |
|---|------------------------|-------------------|------------------|---|------------------------|-------------------|------------------|
| Cruise number | Number of observations | Temperature range | Mean temperature | Cruise number | Number of observations | Temperature range | Mean temperature |
| Main spawning period: | | | | | | | |
| 5701 | 2 | 16.73-17.00 °C | 16.86 °C | 5701 | 1 | 18.66 | 18.66 °C |
| 5702 | 2 | 16.07-16.72 | 16.40 | 5702 | 2 | 18.02-18.27 | 18.14 |
| 5703 | 3 | 14.78-15.72 | 15.24 | 5703 | 1 | 15.35 | 15.35 |
| 5704 | 0 | - | - | 5704 | 1 | 15.72 | 15.72 |
| 5705 | 2 | 15.58-16.88 | 16.23 | 5705 | 3 | 14.94-16.46 | 15.52 |
| 5706 | 2 | 17.22-18.06 | 17.64 | 5706 | 0 | - | - |
| Total | 11 | 14.78-18.06 | 16.36 | Total | 8 | 14.94-18.66 | 16.57 |

| Late spawning period: | | | |
|-----------------------|------------------------|-------------------|------------------|
| Cruise number | Number of observations | Temperature range | Mean temperature |
| 5707 | 4 | 17.84-21.15 | 20.10 |
| 5708 | 4 | 21.08-22.50 | 21.98 |
| 5709 | 4 | 16.61-22.40 | 20.57 |
| 5710 | 4 | 19.86-21.02 | 20.67 |
| Total | 16 | 17.84-22.50 | 20.83 |

| Total | | | |
|---------------|------------------------|-------------------|------------------|
| Cruise number | Number of observations | Temperature range | Mean temperature |
| 5707 | 1 | 16.54 | 16.54 |
| 5708 | 3 | 20.10-24.88 | 22.23 |
| 5709 | 1 | 21.68 | 21.68 |
| 5710 | 1 | 20.64 | 20.64 |
| Total | 6 | 16.54-24.88 | 20.98 |

Text table 4. --Occurrence and abundance (standard haul totals) of
sardine eggs, by month and area, in hauls made during 1957

| Cruise | Central California 60-77 | | Southern California 80-93 | | Northern Baja California 97-107 | | Upper central Baja California 110-120 | | Lower central Baja California 123-137 | | Southern Baja California 140-157 | | Total | |
|---------|-----------------------------|------|------------------------------|-------|------------------------------------|-------|--|--------|--|-------|-------------------------------------|------|--------|--------|
| | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- |
| | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber |
| 5701 | - | - | - | - | 0 | 0 | 2 | 17 | 1 | 12 | - | - | 3 | 29 |
| 5702 | - | - | 0 | 0 | 1 | 4,977 | 2 | 18 | 2 | 155 | 0 | 0 | 5 | 5,150 |
| 5703 | - | - | 2 | 15 | 5 | 386 | 3 | 88 | 1 | 3 | 0 | 0 | 11 | 492 |
| 5704 | - | - | 3 | 1,073 | 3 | 617 | 0 | 0 | 1 | 12 | 0 | 0 | 7 | 1,702 |
| 5705 | 0 | 0 | 6 | 2,015 | 3 | 2,757 | 2 | 74 | 3 | 1,703 | - | - | 14 | 6,549 |
| 5706 | 3 | 201 | 3 | 729 | 0 | 0 | 2 | 5 | 0 | 0 | 0 | 0 | 8 | 935 |
| 5707 | 0 | 0 | 1 | 1,824 | 3 | 146 | 4 | 7,807 | 1 | 47 | - | - | 9 | 9,824 |
| 5708 | - | - | - | - | - | - | 5 | 797 | 3 | 1,349 | - | - | 8 | 2,146 |
| 5709 | - | - | - | - | - | - | 4 | 3,668 | 1 | 263 | - | - | 5 | 3,931 |
| 5710 | - | - | 0 | 0 | 0 | 0 | 4 | 249 | 1 | 8 | - | - | 5 | 257 |
| 5711 | 0 | 0 | 1 | 258 | 0 | 0 | - | - | - | - | - | - | 1 | 258 |
| 5712 | - | - | 0 | 0 | - | - | - | - | - | - | - | - | 0 | 0 |
| Total | 3 | 201 | 16 | 5,914 | 15 | 8,883 | 28 | 12,723 | 14 | 3,552 | 0 | 0 | 76 | 31,273 |
| Percent | 0.64 | | 18.91 | | 28.40 | | 40.68 | | 11.36 | | 0 | | 99.99 | |

Table II
Record of sardine eggs, 1957

| Station | Number of normal eggs | | | | Total number of eggs | | | | | |
|--------------|-----------------------|-------|-------|----|----------------------|-------|-------|----|-------|-------|
| | A | B | C | D | A | B | C | D | Uncl. | n |
| Cruise 5701: | | | | | | | | | | |
| 117.26 | 0 | 3 | 5 | - | 0 | 3 | 5 | - | 0 | 8 |
| 120.25 | 9 | 0 | 0 | - | 9 | 0 | 0 | - | 0 | 9 |
| 123.37 | 0 | 12 | - | - | 0 | 12 | - | - | 0 | 12 |
| Total | 9 | 15 | 5 | - | 9 | 15 | 5 | - | 0 | 29 |
| Cruise 5702: | | | | | | | | | | |
| 103.30 | 190 | 2,353 | 1,333 | - | 380 | 2,924 | 1,673 | - | 0 | 4,977 |
| 120.25 | 0 | 6 | 6 | - | 0 | 6 | 6 | - | 0 | 12 |
| 120.40 | 0 | 2 | 0 | - | 0 | 4 | 0 | - | 2 | 6 |
| 123.37 | 76 | 8 | - | - | 113 | 10 | - | - | 0 | 123 |
| 127.34 | 0 | 32 | - | - | 0 | 32 | - | - | 0 | 32 |
| Total | 266 | 2,401 | 1,339 | - | 493 | 2,976 | 1,679 | - | 2 | 5,150 |
| Cruise 5703: | | | | | | | | | | |
| 83.48 | 0 | 0 | 0 | 0 | 0 | 5 | 2 | 0 | 2 | 9 |
| 90.60 | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 0 | 0 | 6 |
| 97.30 | 15 | 40 | 46 | - | 35 | 40 | 66 | - | 0 | 141 |
| 100.29 | 0 | 0 | 4 | 0 | 0 | 0 | 4 | 0 | 4 | 8 |
| 103.30 | 0 | 23 | 176 | 20 | 0 | 26 | 182 | 23 | 0 | 231 |
| 103.35 | 0 | 3 | - | - | 0 | 3 | - | - | 0 | 3 |
| 107.32 | 3 | 0 | 0 | - | 3 | 0 | 0 | - | 0 | 3 |
| 117.30 | 0 | 0 | 0 | 9 | 0 | 0 | 0 | 9 | 3 | 12 |
| 117.35 | 0 | 0 | 28 | - | 0 | 0 | 28 | - | 0 | 28 |
| 120.25 | 0 | 42 | - | - | 6 | 42 | - | - | 0 | 48 |
| 127.34 | 0 | 0 | 3 | - | 0 | 0 | 3 | - | 0 | 3 |
| Total | 18 | 108 | 257 | 29 | 44 | 116 | 291 | 32 | 9 | 492 |
| Cruise 5704: | | | | | | | | | | |
| 90.55 | 0 | 27 | 0 | - | 0 | 27 | 0 | - | 0 | 27 |
| 93.45 | 0 | 0 | 49 | - | 0 | 0 | 61 | - | 0 | 61 |
| 93.50 | 104 | 254 | 313 | - | 164 | 284 | 522 | - | 15 | 985 |
| 97.30 | 104 | 0 | - | - | 309 | 0 | - | - | 0 | 309 |
| 100.29 | 0 | 16 | 52 | 0 | 0 | 79 | 83 | 5 | 0 | 167 |
| 103.30 | 0 | 46 | 1 | - | 0 | 133 | 8 | - | 0 | 141 |
| 123.42 | 0 | 0 | 12 | - | 0 | 0 | 12 | - | 0 | 12 |
| Total | 208 | 343 | 427 | 0 | 473 | 523 | 686 | 5 | 15 | 1,702 |

Table II (cont'd)
Record of sardine eggs, 1957

| Station | Number of normal eggs | | | | Total number of eggs | | | | | |
|--------------|-----------------------|-------|-----|----|----------------------|-------|-------|----|-------|-------|
| | A | B | C | D | A | B | C | D | Uncl. | n |
| Cruise 5705: | | | | | | | | | | |
| 83.55 | 0 | 496 | 592 | - | 0 | 686 | 803 | - | 68 | 1,557 |
| 83.60 | 0 | 0 | 0 | - | 0 | 3 | 12 | - | 0 | 15 |
| 87.50 | 2 | 0 | - | - | 2 | 0 | - | - | 0 | 2 |
| 90.28 | 38 | 5 | 0 | - | 102 | 5 | 0 | - | 0 | 107 |
| 90.55 | 92 | 92 | - | - | 138 | 184 | - | - | 0 | 322 |
| 93.30 | 0 | 0 | 4 | - | 0 | 0 | 12 | - | 0 | 12 |
| 97.30 | 0 | 424 | 27 | - | 0 | 1,191 | 874 | - | 593 | 2,658 |
| 97.32 | 0 | 0 | 19 | - | 0 | 0 | 22 | - | 0 | 22 |
| 100.33 | 0 | 0 | 34 | - | 0 | 0 | 77 | - | 0 | 77 |
| 120.25 | 22 | 41 | - | - | 26 | 41 | - | - | 4 | 71 |
| 120.50 | 0 | 0 | 0 | - | 0 | 0 | 3 | - | 0 | 3 |
| 127.34 | 0 | 9 | 87 | - | 0 | 15 | 151 | - | 0 | 166 |
| 133.25 | 0 | 9 | 0 | - | 0 | 9 | 0 | - | 0 | 9 |
| 137.23 | 315 | 112 | 0 | - | 1,304 | 224 | 0 | - | 0 | 1,528 |
| Total | 469 | 1,188 | 763 | - | 1,572 | 2,358 | 1,954 | - | 665 | 6,549 |
| Cruise 5706: | | | | | | | | | | |
| 67.55 | 0 | 0 | 0 | 6 | 0 | 0 | 0 | 6 | 0 | 6 |
| 77.55 | 0 | 24 | 87 | 57 | 12 | 27 | 90 | 60 | 0 | 189 |
| 77.60 | 0 | 0 | 6 | - | 0 | 0 | 6 | - | 0 | 6 |
| 90.65 | 227 | 68 | 3 | - | 603 | 87 | 6 | - | 0 | 696 |
| 93.27 | 0 | 0 | - | - | 7 | 0 | - | - | 0 | 7 |
| 93.55 | 7 | 3 | 3 | - | 20 | 3 | 3 | - | 0 | 26 |
| 120.25 | 0 | 2 | - | - | 0 | 2 | - | - | 0 | 2 |
| 120.45 | 0 | 0 | - | - | 0 | 3 | - | - | 0 | 3 |
| Total | 234 | 97 | 99 | 63 | 642 | 122 | 105 | 66 | 0 | 935 |
| Cruise 5707: | | | | | | | | | | |
| 82.47 | 3 | 1,750 | - | - | 3 | 1,821 | - | - | 0 | 1,824 |
| 97.30 | 22 | 8 | - | - | 76 | 24 | - | - | 2 | 102 |
| 103.30 | 0 | 3 | 11 | - | 0 | 3 | 27 | - | 0 | 30 |
| 107.32 | 0 | 7 | - | - | 0 | 14 | - | - | 0 | 14 |
| 117.26 | 422 | 119 | - | - | 589 | 136 | - | - | 0 | 725 |
| 120.25 | 1,307 | 1,782 | - | - | 1,485 | 1,907 | - | - | 0 | 3,392 |
| 120.30 | 472 | 2,862 | - | - | 562 | 2,956 | - | - | 0 | 3,518 |
| 120.40 | 0 | 146 | - | - | 0 | 172 | - | - | 0 | 172 |
| 123.37 | 9 | 38 | - | - | 9 | 38 | - | - | 0 | 47 |
| Total | 2,235 | 6,715 | 11 | - | 2,724 | 7,071 | 27 | - | 2 | 9,824 |

Table II (cont'd)
Record of sardine eggs, 1957

| Station | Number of normal eggs | | | | Total number of eggs | | | | | |
|--------------|-----------------------|-------|---|---|----------------------|-------|---|---|-------|-------|
| | A | B | C | D | A | B | C | D | Uncl. | n |
| Cruise 5708: | | | | | | | | | | |
| 113.30 | 0 | 8 | - | - | 0 | 8 | - | - | 0 | 8 |
| 120.25 | 477 | - | - | - | 524 | - | - | - | 0 | 524 |
| 120.30 | 0 | 57 | - | - | 0 | 77 | - | - | 0 | 77 |
| 120.35 | 2 | 87 | - | - | 4 | 119 | - | - | 2 | 125 |
| 120.40 | 6 | 49 | - | - | 12 | 51 | - | - | 0 | 63 |
| 123.37 | 54 | - | - | - | 103 | - | - | - | 2 | 105 |
| 127.34 | 0 | 881 | - | - | 0 | 1,241 | - | - | 0 | 1,241 |
| 127.40 | 0 | 0 | - | - | 0 | 3 | - | - | 0 | 3 |
| Total | 539 | 1,082 | - | - | 643 | 1,499 | - | - | 4 | 2,146 |
| Cruise 5709: | | | | | | | | | | |
| 113.30 | 0 | 2 | - | - | 0 | 2 | - | - | 0 | 2 |
| 120.25 | 653 | - | - | - | 1,373 | - | - | - | 0 | 1,373 |
| 120.30 | 241 | - | - | - | 420 | - | - | - | 22 | 442 |
| 120.35 | 724 | - | - | - | 1,851 | - | - | - | 0 | 1,851 |
| 123.37 | 111 | - | - | - | 256 | - | - | - | 7 | 263 |
| Total | 1,729 | 2 | - | - | 3,900 | 2 | - | - | 29 | 3,931 |
| Cruise 5710: | | | | | | | | | | |
| 119.33 | 0 | 10 | - | - | 0 | 10 | - | - | 0 | 10 |
| 120.25 | 44 | 70 | - | - | 70 | 134 | - | - | 0 | 204 |
| 120.30 | 6 | 0 | - | - | 6 | 0 | - | - | 6 | 12 |
| 120.35 | 23 | - | - | - | 23 | - | - | - | 0 | 23 |
| 123.37 | 8 | - | - | - | 8 | - | - | - | 0 | 8 |
| Total | 81 | 80 | - | - | 107 | 144 | - | - | 6 | 257 |
| Cruise 5711: | | | | | | | | | | |
| 83.55 | 5 | 0 | - | - | 14 | 244 | - | - | 0 | 258 |
| Total | 5 | 0 | - | - | 14 | 244 | - | - | 0 | 258 |
| Cruise 5712: | | | | | | | | | | |
| | | | | | | | | | | 0 |




Figure 3. --Sardine larvae, 1957: Distribution and relative abundance.

RECORD OF SARDINE LARVAE, 1957

Sardine larvae are grouped by size classes in table III, which have the following midpoints and ranges:

| Midpoint (in mm.) | Range (in mm.) | Midpoint (in mm.) | Range (in mm.) |
|----------------------|-------------------|----------------------|-------------------|
| 3.00 | 2.00-4.25 | 12.75 | 12.26-13.25 |
| 4.75 | 4.26-5.25 | 13.75 | 13.26-14.25 |
| 5.75 | 5.26-6.25 | 14.75 | 14.26-15.25 |
| 6.75 | 6.26-7.25 | 15.75 | 15.26-16.25 |
| 7.75 | 7.26-8.25 | 17.25 | 16.26-18.25 |
| 8.75 | 8.26-9.25 | 19.25 | 18.26-20.25 |
| 9.75 | 9.26-10.25 | 21.25 | 20.26-22.25 |
| 10.75 | 10.26-11.25 | 23.25 | 22.26-24.25 |
| 11.75 | 11.26-12.25 | 25.25 | 24.26-26.25 |

Dis. - Disintegrating larvae that cannot be measured accurately.

The distribution of sardine larvae in 1957 is illustrated in figure 3. Stations occupied during the year are shown as open circles. Four categories of abundance (other than 0) are used. The value shown for a station is the cumulative standard haul total of sardine larvae taken in all occupancies during the year.

There were considerably more hauls containing sardine larvae in 1957 than eggs: 175, as compared to 76. The distribution of catches of eggs and larvae is summarized by area in the following table:

| Station lines | Sardine eggs | | | Sardine larvae | | |
|---------------------------|------------------|-------------|---------|------------------|-------------|---------|
| | occur- rences | num- ber | percent | occur- rences | num- ber | percent |
| 60-77 (complete season) | 3 | 201 | 0.6 | 6 | 94 | 1.0 |
| 80-93 (complete season) | 16 | 5,914 | 18.9 | 21 | 2,028 | 20.6 |
| 97-107 (complete season) | 15 | 8,883 | 28.4 | 22 | 757 | 7.7 |
| 110-120 (Jan.-June) | 11 | 202 | 0.6 | 35 | 1,364 | 13.9 |
| (July-Oct.) | 17 | 12,521 | 40.0 | 48 | 3,889 | 39.6 |
| 123-137 (Jan.-June) | 8 | 1,885 | 6.0 | 21 | 957 | 9.7 |
| (July-Oct.) | 6 | 1,667 | 5.3 | 10 | 316 | 3.2 |
| 140-157 (complete season) | 0 | | | 12 | 428 | 4.3 |
| Total | 76 | 31,273 | 99.8 | 175 | 9,833 | 100.0 |

As has been noted in previous reports, the center of distribution of sardine larvae tends to be somewhat south of the distribution of eggs, due to the pre-dominantly southward transport of eggs and larvae. Off California and northern Baja California (lines 60-107) there were 49 occurrences of larvae to 34 of eggs, or a ratio of occurrences of larvae to eggs of 1.4. Off central Baja California (lines 110-137) there were 114 occurrences of larvae to 42 occurrences of eggs, or a ratio of 2.7 occurrences of larvae to each occurrence of eggs. Only larvae were obtained off southern Baja California (lines 140-157).

The temporal distributions of eggs and larvae agree quite well. In the northern part of the range (lines 60-77), eggs were obtained in June, larvae in June and July (text table 5). Off southern California, eggs were obtained during March through July and in October; larvae were taken in April through July and October; off northern Baja California (lines 97-107) eggs and larvae were taken during February through July; off central Baja California eggs and larvae were taken throughout the period of sampling.

Larger sardine larvae (i.e., larvae 12.26 mm. and larger) were markedly more abundant in the southern part of the distribution. In fact, larger larvae were taken in the northern half of the spawning distribution (lines 60-107) only during June and July, but were taken on all cruises off central and southern Baja California except May and July. The larger larvae occurred in greatest abundance in February and September.

Larger sardine larvae (standard haul totals)

| | California and northern Baja | | | Central and southern Baja | | |
|-----------|------------------------------|-----------------------|-------|----------------------------|-----------------------|-------|
| | California (lines 60-107) | | | California (lines 110-157) | | |
| | 12.26- 15.25 mm. | 15.26 mm. and over | Total | 12.26- 15.25 mm. | 15.26 mm. and over | Total |
| January | 0 | 0 | 0 | 40 | 6 | 46 |
| February | 0 | 0 | 0 | 216 | 74 | 290 |
| March | 0 | 0 | 0 | 2 | 43 | 45 |
| April | 0 | 0 | 0 | 3 | 5 | 8 |
| May | 0 | 0 | 0 | 0 | 0 | 0 |
| June | 12 | 40 | 52 | 0 | 4 | 4 |
| July | 0 | 4 | 4 | 0 | 0 | 0 |
| August | - | - | - | 27 | 6 | 33 |
| September | - | - | - | 91 | 27 | 118 |
| October | 0 | 0 | 0 | 5 | 7 | 12 |
| November | 0 | 0 | 0 | - | - | - |
| December | 0 | 0 | 0 | - | - | - |
| Total | 12 | 44 | 56 | 384 | 172 | 556 |

Text table 5. —Occurrence and abundance (standard haul totals) of sardine larvae, by month and area, in hauls made during 1957

| Cruise | Central California | | Southern California | | Northern Baja California | | Upper central Baja California | | Lower central Baja California | | Southern Baja California | | Total | |
|---------|--------------------|-----------------|---------------------|-----------------|--------------------------|-----------------|-------------------------------|-----------------|-------------------------------|-----------------|--------------------------|-----------------|-------------|-------|
| | 60-77 | 80-93 | 80-93 | 97-107 | 97-107 | 110-120 | 110-120 | 123-137 | 123-137 | 140-157 | 140-157 | 140-157 | | |
| | occur- reces | occur- reces | occur- reces | occur- reces | occur- reces | occur- reces | occur- reces | occur- reces | occur- reces | occur- reces | occur- reces | occur- reces | num- ber | |
| 5701 | - | - | - | 0 | 0 | 7 | 183 | 6 | 576 | - | - | 13 | 759 | |
| 5702 | - | 0 | 0 | 3 | 114 | 10 | 317 | 5 | 164 | 5 | 349 | 23 | 944 | |
| 5703 | - | 0 | 0 | 4 | 254 | 10 | 671 | 8 | 209 | 4 | 31 | 26 | 1,165 | |
| 5704 | - | 3 | 370 | 3 | 44 | 2 | 8 | 0 | 0 | 0 | 0 | 8 | 422 | |
| 5705 | 0 | 8 | 1,332 | 3 | 239 | 2 | 151 | 2 | 8 | - | - | 15 | 1,730 | |
| 5706 | 4 | 8 | 60 | 7 | 55 | 4 | 34 | 0 | 0 | 3 | 48 | 26 | 284 | |
| 5707 | 2 | 1 | 48 | 2 | 51 | 11 | 1,338 | 1 | 3 | - | - | 17 | 1,447 | |
| 5708 | - | - | - | - | - | 14 | 1,257 | 5 | 256 | - | - | 19 | 1,513 | |
| 5709 | - | - | - | - | - | 15 | 1,016 | 1 | 20 | - | - | 16 | 1,036 | |
| 5710 | - | 0 | 0 | 0 | 0 | 8 | 278 | 3 | 37 | - | - | 11 | 315 | |
| 5711 | 0 | 1 | 218 | 0 | 0 | - | - | - | - | - | - | 1 | 218 | |
| 5712 | - | 0 | 0 | - | - | - | - | - | - | - | - | 0 | 0 | |
| Total | 6 | 94 | 21 | 2,028 | 22 | 757 | 83 | 5,253 | 31 | 1,273 | 12 | 428 | 175 | 9,833 |
| Percent | 0.96 | 20.62 | 7.70 | 53.42 | 12.95 | 4.35 | 100.00 | | | | | | | |

Table III
Record of sardine larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Dis. | Total |
|--------------|---------------------------------|-------|-------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|
| | 3.00 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | | |
| Cruise 5701: | | | | | | | | | | | | | | | | | | |
| 117.26 | | | | | 10.6 | | | 21.1 | 2.6 | | | | | | | | | 2.6 |
| .30 | | | | | | | | 21.4 | 10.6 | | | | | | | | | 42.3 |
| .40 | | | | | | | | | | | | | | | | | | 21.4 |
| 120.30 | | | 19.6 | 2.3 | 29.6 | 14.7 | | | | | | | | | | | | 63.9 |
| .35 | 2.3 | 4.7 | 2.3 | 9.3 | 11.7 | | | 2.3 | | | | | | | | | | 37.3 |
| .40 | 2.4 | 1.2 | | | | 4.7 | | 2.4 | 1.2 | | | | | | | | | 7.2 |
| .45 | 2.8 | 2.8 | | | | 2.4 | | 2.8 | | | | | | | | | | 8.4 |
| 123.37 | | | | | | | | | | | | | | | | | | 2.5 |
| 127.34 | 39.0 | 146.4 | 22.0 | 2.4 | 17.1 | 24.4 | 12.2 | 4.8 | 7.3 | 4.8 | | | | | | | | 280.4 |
| .40 | 5.5 | 5.5 | 27.3 | 65.5 | 65.6 | 49.1 | 21.9 | 16.4 | 5.5 | 5.5 | 5.5 | 5.5 | 5.5 | | | | | 267.8 |
| 130.35 | 5.3 | | | | | | | | | | | | | | | | | 5.3 |
| .40 | 5.8 | | | | | 11.6 | | | | | | | | | | | | 17.4 |
| 137.23 | | 2.4 | | | | | | | | | | | | | | | | 2.4 |
| Total | 2.8 | 57.6 | 160.2 | 43.9 | 79.2 | 109.0 | 130.1 | 87.5 | 41.1 | 23.7 | 10.3 | 5.5 | 5.5 | 5.5 | | | 2.5 | 758.9 |
| Cruise 5702: | | | | | | | | | | | | | | | | | | |
| 103.30 | | 27.2 | | | | | | | | | | | | | | | | 27.2 |
| .35 | 27.7 | | | | | | | | | | | | | | | | | 27.7 |
| 107.32* | 11.9 | 47.7 | | | | | | | | | | | | | | | | 59.6 |
| .35* | | | | | | | | | | | | | | | | | | |
| 110.33 | 10.4 | | | | | | | | | | | | | | | | | 10.4 |
| .40 | | 5.6 | | | | | | | | | | | | 2.8 | | | | 8.4 |
| 113.40 | 2.5 | 2.5 | 7.5 | | | | | | | | | | | | | | | 12.5 |
| 117.26 | | | | | | | | | | | | | | | | | | 37.6 |
| .30 | | | | | | | 9.4 | | 18.8 | | | | | | | | | 200.5 |
| 120.25 | 12.0 | | | | | | 33.4 | | 33.4 | | | | | | | | | 12.0 |
| .30 | | | | | | | | | | | | | | | | | | 11.7 |
| .35 | | | | 4.9 | | | | | | | | | | | | | | 4.9 |
| .40 | 4.7 | 2.4 | 7.1 | 2.4 | | | | | | | | | | | | | | 16.6 |
| .45 | | | | | | | | | | | | | | | | | | 2.8 |

* - Samples combined. Values were adjusted accordingly.

Table III (cont'd)
Record of sardine larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Total | | |
|-----------------------|---------------------------------|-------|-------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | 3.00 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | | 23.25 | 25.25 |
| Cruise 5702 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 123.37 | 13.6 | | | | | | | | | | | 4.5 | | 4.5 | | | | | 13.6 |
| .55 | | | | | | | | | | | | | | | | | | | 9.0 |
| 127.34 | 57.9 | | | | | | | | | | | | | | | | | | 57.9 |
| 130.30 | | | | | | 6.4 | | | | | | | | | | | | | 6.4 |
| 137.23 | | 4.3 | 25.5 | 25.6 | 12.8 | 4.3 | 4.3 | | | | | | | | | | | | 76.8 |
| 140.35 | | | | | | | | 7.3 | 3.7 | 3.7 | | | | | | | | | 14.7 |
| .40 | | | | | | | | 2.8 | | | | | | | | | | | 8.4 |
| 147.20 | | | | | | | | 3.1 | 12.4 | 3.1 | 15.5 | 3.1 | | 2.8 | | | | | 8.4 |
| .25 | 3.6 | 21.8 | 32.7 | | | | | 3.1 | 3.1 | 3.1 | 3.1 | | 3.1 | 3.1 | | | | | 46.5 |
| 150.25 | 3.1 | 9.2 | 24.6 | 40.1 | 33.9 | 24.6 | 30.8 | 21.5 | 30.8 | 3.1 | | | | | | | | | 58.1 |
| Total | 114.8 | 85.9 | 125.6 | 72.8 | 55.3 | 38.2 | 38.4 | 62.7 | 61.7 | 50.0 | 61.2 | 104.8 | 48.7 | 19.0 | 3.1 | 2.8 | | | 945.0 |
| Cruise 5703: | | | | | | | | | | | | | | | | | | | |
| 103.30 | 29.4 | 35.1 | 23.5 | 14.7 | 2.9 | | | | | | | | | | | | | | 105.6 |
| .35 | | 26.1 | 2.9 | | | | | | | | | | | | | | | | 29.0 |
| 107.32 | | 36.5 | 63.1 | 16.6 | | | | | | | | | | | | | | | 116.2 |
| .35 | | | 3.2 | | | | | | | | | | | | | | | | 3.2 |
| 110.40 | | | 3.3 | | | | | | | | | | | | | | | | 3.3 |
| 113.35 | 12.7 | | | 12.7 | | | | | | | | | | | | | | | 25.4 |
| .40 | | 14.0 | | | | | | | | | | | | | | | | | 14.0 |
| 117.30 | | 18.5 | 224.8 | 6.2 | | | | | | | | | | | | | | | 249.5 |
| .35 | | 151.8 | 151.8 | 13.8 | | | | | | | | | | | | | | | 317.4 |
| .40 | | 9.0 | 6.0 | | | | | | | | | | | | | | | | 15.0 |
| .50 | | | 6.2 | | | | | | | | | | | | | | | | 6.2 |
| .70 | | 3.2 | | | | | | | | | | | | | | | | | 3.2 |
| 119.33 | | | | 6.5 | 6.5 | | | | | | | | | | | | | | 13.0 |
| 120.25 | 24.0 | | | | | | | | | | | | | | | | | | 24.0 |
| 123.42 | | | | | | | | | | | | | | | | | | | 12.0 |
| 127.40 | | 7.5 | 15.0 | | | | | 7.5 | | | | | | | | | | | 30.0 |
| .45 | | 43.8 | 43.8 | | | | | | | | | | | | | | | | 98.6 |
| .50 | | 6.3 | 18.9 | 6.3 | 6.3 | | | | | | | | | | | | | | 37.8 |
| .55 | | | | | | | | | | | | | | | | | | | 13.5 |

Table III (cont'd)
Record of sardine larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Total | | | |
|-----------------------|---------------------------------|-------|-------|-------|-------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| | 3.00 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | | 23.25 | 25.25 | Dis. |
| Cruise 5703 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 130.30 | | | | | | | | | | | | | | | 10.7 | | | | 10.7 | |
| 133.35 | | | 3.2 | | | | | | | | | | | | | | | | | 3.2 |
| 137.23 | | | | | | | | | | | | | | | | | | | | 2.9 |
| 140.35 | | | | | | | | | | | | | | 2.8 | 11.0 | | | | | 13.8 |
| 143.50 | | | | | | | | | | | | | | 3.1 | | | | | | 3.1 |
| 150.30 | | | | | | | | | | | | | | | 2.5 | | | | | 7.5 |
| .40 | | | | | | | | | | | | | | | 4.2 | | | | | 6.3 |
| Total | 66.1 | 345.5 | 549.9 | 92.1 | 35.2 | 6.3 | 6.5 | 7.5 | 9.6 | 2.5 | | | 13.8 | 16.2 | 13.2 | | | | | 1,164.4 |
| Cruise 5704: | | | | | | | | | | | | | | | | | | | | |
| 90.45 | 26.4 | 13.2 | 13.2 | 13.2 | | | | | | | | | | | | | | | | 66.0 |
| 93.45 | | 61.8 | 185.4 | | 12.4 | | | | | | | | | | | | | | | 259.6 |
| .50 | | 14.9 | 29.8 | | | | | | | | | | | | | | | | | 44.7 |
| 100.33 | | | 5.2 | | | | | | | | | | | | | | | | | 5.2 |
| 103.30 | | 11.5 | 14.3 | 7.2 | 2.9 | | | | | | | | | | | | | | | 35.9 |
| 107.35 | | | 3.2 | | | | | | | | | | | | | | | | | 3.2 |
| 117.60 | | | | | | | | | | | | | | | | | | | 5.3 | 5.3 |
| .70 | | | | | | | | | | | | | | | | | | | | 2.7 |
| Total | 26.4 | 101.4 | 251.1 | 20.4 | 15.3 | | | | | | | | | | | | | | | 422.6 |
| Cruise 5705: | | | | | | | | | | | | | | | | | | | | |
| 83.51 | | | 3.4 | | | | | | | | | | | | | | | | | 3.4 |
| 90.50 | | 2.8 | 5.6 | | | | | | | | | | | | | | | | | 8.4 |
| .55 | 138.8 | 505.7 | 66.5 | | 5.8 | 2.9 | | | | | | | | | | | | | | 719.7 |
| .60 | 13.7 | 41.0 | | | 10.3 | | 10.2 | | | | | | | | | | | | | 75.2 |
| 93.55 | | | | | | | | | | | | | | | | | | | | 40.6 |
| .60 | | 24.7 | 151.4 | 5.8 | 29.0 | 5.8 | | | | | | | | | | | | | | 463.5 |
| .65 | | | | 5.1 | 132.9 | 55.6 | | | | | | | | | | | | | | 10.2 |
| .70 | | | | | 5.1 | | | | | | | | | | | | | | | 10.8 |
| 97.30 | 178.4 | | | 4.5 | 8.1 | 2.7 | | | | | | | | | | | | | | 196.4 |
| .32 | 9.6 | | | | 4.5 | 4.5 | 4.5 | | | | | | | | | | | | | 12.8 |
| 100.33 | 12.8 | | 3.2 | | | | | | | | | | | | | | | | | 29.7 |
| 117.45 | 137.4 | | 2.1 | | | | | | | | | | | | | | | | | 137.4 |
| .50 | 5.6 | 8.4 | | | | | | | | | | | | | | | | | | 14.0 |
| 127.34 | 5.8 | | | | | | | | | | | | | | | | | | | 5.8 |
| 133.30 | 2.3 | | | | | | | | | | | | | | | | | | | 2.3 |
| Total | 504.4 | 595.3 | 232.2 | 114.3 | 190.6 | 78.7 | 14.7 | | | | | | | | | | | | | 1,730.2 |

Table III (cont'd)
Record of sardine larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | Dis. | Total | |
|--------------|---------------------------------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|-------|
| | 3.00 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | | | 21.25 |
| Cruise 5706: | | | | | | | | | | | | | | | | | | |
| 73.70 | | | | 3.1 | | 12.4 | 21.7 | 12.4 | 3.1 | | | | | | | | | 52.7 |
| .80 | | | | | | | | 6.2 | | | | | | | | | | 6.2 |
| 77.55 | | | | 6.0 | | 3.0 | | | | | | | | | | | | 9.0 |
| .60 | | 6.4 | | 6.4 | | | | | 6.4 | | | | | | | | | 19.2 |
| 80.60 | | | | 11.0 | | | | | | 5.5 | | | | | | | | 16.5 |
| .80 | | | | | | | | | 6.4 | | | | | | | | | 6.4 |
| 83.85 | | | | | | | | | | 2.9 | | | | | | | | 2.9 |
| 87.90 | | | | | | | | | | | | | | | 3.0 | | | 3.0 |
| 90.65 | 3.1 | | | | | | | | | | | | | | | | | 3.1 |
| 93.55 | | 6.6 | | | | | | | | | 3.3 | | | | | | | 16.5 |
| .60 | | | | | | | | | | | | | | 6.6 | | | | 7.4 |
| .65 | | | | | | | | | | | | | | 7.4 | | | | 3.7 |
| 97.55 | | | | | | | | | | | | | | 3.7 | | | | 6.6 |
| .60 | | | | | | | | | | | | | | 6.6 | | | | 3.0 |
| 100.33 | | 26.9 | 6.0 | | | | | | | | | | | 3.0 | | | | 32.9 |
| .55 | | | | | | | | | | | | | | | | 3.3 | | 3.3 |
| .70 | | 2.8 | | | | | | | | | | | | | | | | 2.8 |
| 103.35 | | | | | | | | | | | | | | | | | | 2.9 |
| .60 | | | | | | | | | | | | | | | | | | 3.4 |
| 110.45 | 6.1 | 6.1 | | | | | | | | | | | | | | | | 3.4 |
| 117.45 | 5.1 | 11.5 | | | | | | | | | | | | | | | | 12.2 |
| 120.35 | | | | 3.4 | | | | | | | | | | | | | | 16.6 |
| .40 | | 1.8 | | | | | | | | | | | | | | | | 3.4 |
| 148.20 | | 17.4 | | | 2.9 | | | | 2.9 | | | | | | | | | 1.8 |
| .25 | | | | | | | | | | | | | | | | | | 23.2 |
| .30 | | | | | | | | | 10.9 | | | | | | | | | 18.1 |
| | | | | | | | | 6.9 | | | | | | | | | | 6.9 |
| Total | 14.3 | 79.5 | 27.2 | 9.1 | 20.3 | 15.4 | 21.7 | 18.6 | 18.8 | 8.4 | | 3.3 | 27.3 | 12.9 | | 3.3 | 3.6 | 283.7 |

Table III (cont'd)
Record of sardine larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Dis. | Total |
|--|---------------------------------|-------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|
| | 3.00 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | | |
| Cruise 5707: | | | | | | | | | | | | | | | | | | |
| 67.80 | | 3.2 | | | | | | | | | | | | | | | 4.2 | 3.2 |
| 70.70 | | | | | | | | | | | | | | | | | | 4.2 |
| 82.47 | 7.9 | 10.5 | 5.2 | | 5.2 | 5.2 | 7.9 | 2.6 | 2.6 | | | | | | | | | 47.1 |
| 97.30 | | 2.0 | | | | | | | | | | | | | | | | 2.0 |
| 103.30 | 39.1 | 8.2 | 1.6 | | | | | | | | | | | | | | | 48.9 |
| 110.33 | | | | 3.4 | | | | | | | | | | | | | | 3.4 |
| 113.30 | | | 6.2 | 4.2 | 8.2 | 14.5 | 16.6 | 6.2 | | | | | | | | | | 55.9 |
| .35 | 4.0 | | | | | | | | | | | | | | | | | 4.0 |
| .35 | 3.8 | | | 3.4 | | 3.4 | | 3.4 | | | | | | | | | | 20.4 |
| .40 | | | | | 3.4 | | | | | | | | | | | | | 3.8 |
| 119.33 | 10.3 | 3.4 | 3.4 | | 3.4 | | | | | | | | | | | | | 3.4 |
| 120.25 | 665.3 | 35.7 | 3.0 | | 3.4 | | | | | | | | | | | | | 20.5 |
| .30 | 361.7 | 38.8 | 6.4 | 3.2 | | | | | | | | | | | | | | 704.0 |
| .35 | | | 8.6 | 37.3 | 5.7 | | | | | | | | | | | | | 410.1 |
| .40 | 3.8 | 18.9 | 7.6 | 18.9 | 7.6 | 1.9 | | 1.9 | | | | | | | | | | 51.6 |
| 130.45 | | 3.0 | | | | | | | | | | | | | | | | 60.6 |
| Total 1095.9 133.9 42.0 70.4 26.7 31.8 24.5 14.1 2.6 | | | | | | | | | | | | | | | | | | |
| Cruise 5708: | | | | | | | | | | | | | | | | | | |
| 113.30 | | | 2.7 | | 2.7 | | | | | | | | | | | | | 5.4 |
| .35 | | | | | | 2.7 | | | | | | | | | | | | 2.7 |
| 115.27 | | 2.9 | 28.9 | 69.4 | 49.1 | 28.9 | 11.6 | 8.7 | | | | | | | | | | 199.5 |
| 117.26 | | 11.7 | 17.6 | 32.1 | 46.7 | 43.8 | 55.5 | 26.2 | 5.8 | 5.8 | 5.8 | 2.9 | | | | | | 253.9 |
| .30 | | | | | 5.8 | 2.9 | 8.8 | 2.9 | | | | | | | | | | 20.4 |
| .35 | | | | | 6.4 | | | | | | | 3.2 | | | | | | 9.6 |
| 118.25 | 3.0 | 33.2 | 3.0 | 3.0 | | | | | | | | | | | | | | 42.2 |
| .35 | 78.0 | 137.3 | | | | | | | | | | | | | | | | 215.3 |
| 119.33 | 34.8 | 2.9 | | | | | | | | | | | | | | | | 37.7 |
| 120.25 | 8.9 | 41.5 | 62.2 | 32.5 | 20.7 | 38.5 | 14.8 | 6.0 | 3.0 | 3.0 | | | | | | | | 231.1 |
| .30 | | 6.7 | 3.3 | 3.3 | 10.0 | | | | | | | | | | | | | 23.3 |
| .35 | 55.4 | 19.2 | | 2.1 | | | | | | | | | | | | | | 76.7 |
| .40 | 4.0 | 12.2 | 16.3 | 20.3 | 24.4 | 28.4 | 12.2 | 10.2 | 4.1 | 4.1 | | | | | | | | 136.2 |
| .45 | | | | | | | | | 2.9 | | | | | | | | | 2.9 |
| 123.37 | 4.8 | 12.2 | 12.2 | 4.8 | 2.4 | | | | | | | | | | | | | 36.4 |
| .45 | | 3.0 | | | | | | | | | | | | | | | | 3.0 |
| 127.34 | 5.8 | 8.7 | 8.6 | 31.7 | 60.5 | 34.6 | 20.2 | 8.7 | 5.8 | | | | | | | | | 184.6 |

Table III (cont'd)
Record of sardine larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | Dis. | Total | | | | |
|-----------------------|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|-------|-------|-------|---------|
| | 3.00 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.76 | 11.75 | 12.75 | 13.76 | 14.75 | | | 15.75 | 17.25 | 19.25 | 21.25 |
| Cruise 5708 (cont'd): | | | | | | | | | | | | | | | | | | |
| 127.40 | | 6.7 | 3.4 | 13.4 | 3.4 | | | | | | | | | | | | | 26.9 |
| 130.30 | | | | 2.7 | 2.7 | | | | | | | | | | | | | 5.4 |
| Total | 194.7 | 276.9 | 126.9 | 150.3 | 207.1 | 225.8 | 140.9 | 97.6 | 59.5 | 15.8 | 5.8 | 5.8 | 3.2 | 2.9 | | | | 1,513.2 |
| Cruise 5709: | | | | | | | | | | | | | | | | | | |
| 110.33 | | | | | | | 7.1 | 9.5 | 4.7 | | | | | | | 2.4 | | 2.4 |
| 113.30 | | | 2.4 | 3.1 | | | | | | | | | | | | | | 23.7 |
| .35 | | | 3.1 | 3.1 | | | | | | | | | | 3.1 | | | | 6.2 |
| 115.30 | | | 5.3 | 2.6 | | | | | | | | | | | | | | 3.1 |
| 117.26 | | | 3.0 | 3.0 | | | | | | | | | | | | | | 7.9 |
| .35 | | | | | | | | | | | | | | | | | | 6.0 |
| .40 | | 3.3 | | | 6.6 | 19.8 | 6.6 | 3.3 | | | 6.6 | 3.3 | | | | | | 49.5 |
| 118.25 | 2.5 | 10.1 | 5.0 | 2.5 | 7.6 | 5.1 | 2.5 | 5.0 | | | | | | | | | | 40.3 |
| .30 | 15.0 | 15.0 | 41.2 | 37.4 | 67.5 | 78.7 | 78.7 | 37.5 | 11.3 | | | | | | | | | 393.6 |
| 119.33 | 6.0 | 11.9 | 17.8 | 17.9 | 6.0 | 9.0 | 6.0 | 6.0 | 3.0 | | | | | | | | | 83.6 |
| 120.25 | 6.3 | 3.2 | 3.2 | | 3.2 | 6.3 | | | | | | | | | | | | 22.2 |
| .30 | 3.1 | 12.4 | 6.2 | | 3.1 | | | | | | | | | | | | | 27.9 |
| .35 | 3.0 | 18.0 | 3.0 | 12.0 | 12.0 | 15.0 | 12.0 | 3.0 | 3.0 | | | | | | | | | 93.0 |
| .40 | | 5.9 | 26.6 | 8.9 | 8.9 | 17.6 | 47.2 | 50.1 | 38.3 | 23.6 | 8.8 | 5.9 | 3.0 | 3.0 | | | | 247.8 |
| .45 | | | | | 5.6 | 2.8 | | | | | | | | | | | | 8.4 |
| 123.37 | 3.4 | 10.1 | 3.4 | | 3.4 | | | | | | | | | | | | | 20.3 |
| Total | 36.8 | 79.0 | 114.8 | 92.6 | 114.5 | 110.6 | 142.4 | 143.2 | 83.5 | 55.9 | 26.6 | 8.8 | 12.5 | 9.4 | 3.0 | 2.4 | | 1,035.9 |
| Cruise 5710: | | | | | | | | | | | | | | | | | | |
| 113.30 | | | | | | 2.8 | | | | | | | | | | | | 2.8 |
| 117.26 | 6.0 | 69.2 | 21.0 | 3.0 | 6.0 | 6.0 | 3.0 | 3.0 | 3.0 | | | | | | | | | 120.2 |
| .35 | | | | | | | 2.4 | | 2.4 | | | | | | | | | 4.8 |
| 119.33 | 5.2 | 10.4 | 10.4 | 2.6 | 7.8 | 7.8 | 7.8 | | | | | | | | | | | 52.0 |
| 120.25 | 1.8 | 1.8 | 11.1 | 11.0 | 7.3 | | | 1.8 | | | | | | | | | | 34.8 |
| .30 | | 27.5 | 10.5 | 2.1 | 2.1 | | | | | | | | | | | | | 42.2 |
| .35 | 4.6 | 9.1 | 2.3 | | 2.3 | | | | | | | | | | | | | 18.3 |
| .60 | | | | | | | | | 3.0 | | | | | | | | | 3.0 |
| 123.37 | 10.8 | 13.6 | 2.7 | 2.7 | | | | | | | | | | 3.1 | | | | 29.8 |
| .60 | | | | | | | | | | | | | | | | | | 3.1 |
| 127.40 | | | | | | | | | | | | | | 3.8 | | | | 3.8 |
| Total | 28.4 | 131.6 | 58.0 | 21.4 | 25.5 | 16.6 | 13.2 | 4.8 | 3.0 | 6.4 | | | | 3.1 | 3.8 | | | 314.8 |
| Cruise 5711: | | | | | | | | | | | | | | | | | | |
| 83.55 | 167.9 | 13.8 | 18.4 | 6.9 | 6.9 | 4.6 | | | | | | | | | | | | 218.5 |
| Total | 167.9 | 13.8 | 18.4 | 6.9 | 6.9 | 4.6 | | | | | | | | | | | | 218.5 |




Figure 4. --Anchovy larvae, 1957: Distribution and relative abundance.

RECORD OF ANCHOVY LARVAE, 1957

The larvae of the northern anchovy, *Engraulis mordax*, are recorded by size in table IV. The size classes are similar to those used for sardine larvae except for the two smallest size categories which have the following midpoints and ranges:

| Midpoint (mm.) | Size range (mm.) |
|-------------------|---------------------|
| 2.50 | 1.76-3.25 |
| 3.75 | 3.26-4.25 |

The distribution and relative abundance of anchovy larvae are illustrated in figure 4. The value for each station is the cumulative standard haul total for the year.

Anchovy larvae ranked first in abundance in 1957, as in all recent years. The average number of anchovy larvae taken per haul has been exceptionally constant during the past three years:

| Year | Total hauls taken | Occurrences | Total anchovy larvae (standard haul totals) | Average per haul |
|------|----------------------|-------------|--|---------------------|
| 1955 | 1,375 | 616 | 140,183 | 102 |
| 1956 | 1,397 | 536 | 134,931 | 97 |
| 1957 | 1,493 | 580 | 146,631 | 98 |

In the previous report in this series, the unusual distribution of anchovy larvae during 1956 was noted. During that year, considerably fewer anchovy larvae were taken in the northern half of the survey area (lines 60-107) than in the southern portion. The most unusual feature of the distribution was the presence of large numbers of larvae off southern Baja California (lines 140-157), an area in which anchovy larvae are usually uncommon.

| Station lines | 1955 | | 1956 | | 1957 | |
|---------------|---------|---------------------|---------|---------------------|---------|---------------------|
| | Number | Percent of total | Number | Percent of total | Number | Percent of total |
| 60-77 | 38 | 0.03 | 629 | 0.5 | 206 | 0.1 |
| 80-93 | 30,147 | 21.5 | 17,838 | 13.3 | 79,923 | 54.5 |
| 97-107 | 30,092 | 21.5 | 8,463 | 6.3 | 10,791 | 7.4 |
| 110-120 | 68,568 | 48.9 | 61,565 | 45.8 | 32,583 | 22.2 |
| 123-137 | 11,269 | 8.0 | 20,884 | 15.1 | 23,014 | 15.7 |
| 140-157 | 69 | 0.05 | 25,552 | 19.0 | 114 | 0.1 |
| Total | 140,183 | 99.98 | 134,931 | 100.0 | 146,631 | 100.0 |

Only a few anchovy larvae were taken off southern Baja California in 1957 - less than 0.1 percent of the number taken in the CCOFI survey area, as compared to 19.0 percent in this area in 1956. A much higher percentage of the larvae were taken in the northern half of the survey area (station lines 60-107), however: 62.0 percent as compared to 20.1 percent in 1956. Water temperatures in the upper mixed layer, the stratum in which anchovy larvae occur, were higher in 1957 than in 1956, especially in the area off central and southern Baja California. The northward shift in larval distribution is probably related, directly or indirectly, to the increase in water temperatures in the CCOFI area.

The abundance of anchovy larvae in different parts of the CCOFI survey area in 1957 is summarized by month in text table 6. Almost a third of the larvae were obtained during the peak month, February. This was the month of peak abundance in all major subareas, an unusual circumstance as can be seen by comparing the 1957 areal abundance with that given for 1956 (Ahlstrom 1958, text table 5) or 1955 (Ahlstrom and Kramer 1957, text table 4). Anchovy spawning ordinarily takes place throughout the year in the area between Point Conception and Point San Juanico (station lines 80-137). There was a marked decrease in the number of larvae in the southern part of the range (lines 123-137) during the latter part of the year. There were only three occurrences in 64 hauls taken in this area during August through October, and these contained only small numbers of larvae. The increase in temperatures in this area (3°C higher than the average of the preceding six years during these months) may account for the decrease in anchovy larvae.

The monthly abundance of anchovy larvae during 1957 is compared with that of the preceding four years in text table 7. Throughout the five-year period, more anchovy larvae were taken during the winter months (January-March) than in the other seasons. Coverage during the last five months of the year has varied in intensity during this period, but in no year was it as thorough as during the earlier months, hence abundance during these months is underestimated. Even allowing for this fact, there is a marked decrease in numbers of larvae during the summer and fall months as compared to the other two seasons.

The abundance of anchovy larvae of different sizes is summarized for the five-year period, 1953-1957, in text table 8. A comparison of the larger larvae (12.26 mm. and over) is as follows:

| Year | Standard haul total | Larvae over 12.26 mm. | Percent of total |
|------|---------------------|-----------------------|------------------|
| 1953 | 99,160 | 2,294 | 2.31 |
| 1954 | 161,326 | 4,366 | 2.71 |
| 1955 | 140,188 | 3,773 | 2.69 |
| 1956 | 134,931 | 4,599 | 3.41 |
| 1957 | 146,631 | 4,253 | 2.90 |

Text table 6. —Occurrence and abundance (standard haul totals) of anchovy larvae (*Engraulis mordax*), by month and area, in hauls made during 1957

| Cruise | Central California 60-77 | | Southern California 80-93 | | Northern Baja California 97-107 | | Upper central Baja California 110-120 | | Lower central Baja California 123-137 | | Southern Baja California 140-157 | | Total | |
|---------|--------------------------|------|---------------------------|--------|---------------------------------|--------|---------------------------------------|--------|---------------------------------------|--------|----------------------------------|------|--------|---------|
| | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- |
| | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber |
| 5701 | - | - | - | - | 4 | 604 | 17 | 6,581 | 11 | 3,314 | - | - | 32 | 10,499 |
| 5702 | - | - | 23 | 23,280 | 17 | 3,486 | 17 | 9,279 | 22 | 12,202 | 3 | 13 | 82 | 48,260 |
| 5703 | - | - | 23 | 6,723 | 9 | 1,052 | 19 | 4,725 | 17 | 1,750 | 5 | 79 | 73 | 14,329 |
| 5704 | - | - | 25 | 16,879 | 21 | 2,237 | 13 | 734 | 12 | 364 | 3 | 18 | 74 | 20,232 |
| 5705 | 3 | 14 | 21 | 15,203 | 12 | 2,090 | 30 | 2,906 | 11 | 582 | - | - | 77 | 20,795 |
| 5706 | 7 | 122 | 33 | 11,098 | 8 | 103 | 31 | 5,107 | 10 | 1,973 | 1 | 4 | 90 | 18,407 |
| 5707 | 5 | 23 | 24 | 3,834 | 9 | 493 | 17 | 1,735 | 12 | 2,801 | - | - | 67 | 8,886 |
| 5708 | - | - | - | - | - | - | 17 | 1,440 | 2 | 6 | - | - | 19 | 1,446 |
| 5709 | - | - | - | - | - | - | 7 | 35 | 0 | 0 | - | - | 7 | 35 |
| 5710 | - | - | 13 | 311 | 8 | 487 | 4 | 41 | 1 | 22 | - | - | 26 | 861 |
| 5711 | 2 | 47 | 19 | 2,153 | 5 | 173 | - | - | - | - | - | - | 26 | 2,373 |
| 5712 | - | - | 6 | 442 | 1 | 66 | 0 | 0 | - | - | - | - | 7 | 508 |
| Total | 17 | 206 | 187 | 79,923 | 94 | 10,791 | 172 | 32,583 | 98 | 23,014 | 12 | 114 | 580 | 146,631 |
| Percent | | 0.14 | | 54.51 | | 7.36 | | 22.22 | | 15.70 | | 0.07 | | 100.00 |

Text table 7.--Monthly abundance of anchovy larvae
1953-1957, based on standard haul summations

| Month | 1953 | 1954 | 1955 | 1956 | 1957 |
|-----------|--------|---------|---------|---------|---------|
| January | 12,820 | 24,853 | 40,143 | 8,847 | 10,499 |
| February | 16,199 | 32,709 | 30,951 | 29,136 | 48,261 |
| March | 15,841 | 34,314 | 25,081 | 16,642 | 14,328 |
| April | 6,920 | 32,833 | 16,490 | 22,857 | 20,232 |
| May | 3,496 | 12,494 | 4,201 | 11,937 | 20,796 |
| June | 4,886 | 6,497 | 12,656 | 18,261 | 18,407 |
| July | 5,639 | 8,661 | 7,058 | 14,717 | 8,886 |
| August | 2,856 | 1,875 | - | 9,635 | 1,446 |
| September | 3,009 | - | 720 | 373 | 35 |
| October | 6,096 | 1,248 | 647 | 825 | 862 |
| November | 7,470 | - | 1,155 | 1,424 | 2,371 |
| December | 13,928 | 5,842 | 1,086 | 277 | 508 |
| Total | 99,160 | 161,326 | 140,188 | 134,931 | 146,631 |

Text table 8.--Abundance of anchovy larvae by size categories,
1953-1957, based on standard haul summations

| Size (mm.) | 1953 | 1954 | 1955 | 1956 | 1957 |
|---------------|--------|---------|---------|---------|---------|
| 2.50 | 15,112 | 33,466 | 19,707 | 13,993 | 19,983 |
| 3.75 | 19,453 | 24,407 | 29,109 | 21,246 | 23,640 |
| 4.75 | 17,462 | 19,898 | 17,187 | 15,463 | 19,915 |
| 5.75 | 12,153 | 19,428 | 18,540 | 14,635 | 18,276 |
| 6.75 | 10,474 | 19,992 | 15,738 | 16,710 | 17,126 |
| 7.75 | 7,780 | 14,596 | 12,646 | 15,105 | 14,592 |
| 8.75 | 5,764 | 10,747 | 9,691 | 12,351 | 11,930 |
| 9.75 | 4,283 | 6,902 | 6,850 | 9,946 | 8,422 |
| 10.75 | 2,627 | 4,406 | 4,386 | 6,839 | 5,179 |
| 11.75 | 1,619 | 2,848 | 2,392 | 4,027 | 3,237 |
| 12.75 | 804 | 1,812 | 1,458 | 2,159 | 1,817 |
| 13.75 | 562 | 999 | 838 | 1,046 | 942 |
| 14.75 | 325 | 639 | 523 | 535 | 587 |
| 15.75 | 187 | 371 | 302 | 274 | 302 |
| 17.25 | 236 | 304 | 412 | 310 | 381 |
| 19.25 | 70 | 178 | 144 | 217 | 119 |
| 21.25 | 42 | 37 | 43 | 15 | 66 |
| 23.25 | 68 | 26 | 53 | 43 | 39 |
| Disintegrated | 139 | 270 | 169 | 17 | 78 |
| Total | 99,160 | 161,326 | 140,188 | 134,931 | 146,631 |

Table IV
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Total | | | | | |
|--------------|---------------------------------|--------|--------|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------------|
| | 2.60 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | | 17.25 | 19.25 | 21.25 | 23.25 | Dis. |
| Cruise 5701: | | | | | | | | | | | | | | | | | | | | |
| 103.30 | 7.3 | 14.8 | 22.1 | 17.2 | 7.3 | 2.4 | 2.4 | | | | | | | | | | | | | 73.5 |
| .35 | 2.9 | 11.6 | 14.5 | 2.9 | | | | | | | | | | | | | | | | 31.9 |
| 107.32 | 168.2 | 98.6 | 23.2 | 29.0 | 29.0 | 11.6 | 17.4 | 5.8 | | | | | | | | | | | | 382.8 |
| .35 | 66.7 | 23.2 | 8.7 | 5.8 | | 5.8 | 5.8 | | | | | | | | | | | | | 116.0 |
| 110.40 | | | | | 2.9 | | | | | | | | | | | | | | | 2.9 |
| 113.30 | 57.3 | 15.0 | 29.9 | 34.8 | 77.2 | 44.8 | 22.4 | 20.0 | 2.5 | 2.5 | | | | | | | | | | 306.4 |
| .35 | 2.7 | 5.4 | 5.4 | 8.1 | 13.6 | 2.7 | | | | | | | | | | | | | | 37.9 |
| .40 | | | | | | | | | 2.6 | | | | | | | | | | | 2.6 |
| 117.26 | 111.0 | 719.8 | 221.9 | 252.9 | 134.1 | 85.2 | 43.8 | 15.5 | 5.2 | 5.2 | | | | | | | | | | 1,594.6 |
| .30 | 31.7 | 211.2 | 380.1 | 1108.8 | 591.4 | 147.8 | 73.9 | 95.0 | 63.3 | 31.7 | | | | | | | | | | 2,734.9 |
| .35 | 21.4 | 42.7 | 32.1 | 53.4 | 53.4 | 10.7 | 21.4 | 10.7 | 10.7 | | | | | | | | | | | 256.5 |
| .40 | | 235.0 | 309.7 | 149.5 | 53.4 | 53.4 | 85.4 | 21.4 | 21.4 | 21.4 | 10.7 | 10.7 | 10.7 | 10.7 | 10.7 | | | | | 1,004.1 |
| .50 | | 15.8 | 21.1 | | | 13.2 | 18.5 | 29.1 | 18.5 | 7.9 | 10.5 | 5.2 | 2.6 | | | | | | | 142.4 |
| .60 | | 5.4 | 8.1 | 2.7 | 2.7 | 5.4 | 2.7 | 2.7 | | | | | | | | | | | 2.7 | 29.7 |
| 120.25 | | | | | | 18.2 | 18.2 | 27.3 | | | | | | | | | | | | 63.7 |
| .30 | | 9.8 | 14.8 | 39.4 | 9.8 | 19.6 | 68.9 | 54.1 | 64.0 | 44.3 | 9.8 | 4.9 | | | | | | | | 339.4 |
| .35 | | 4.6 | | | | | | | | 2.3 | | | | | | | | | | 6.9 |
| .40 | 2.4 | 6.3 | 3.8 | | | | | | | | | | | | | | | | | 12.5 |
| .45 | | 5.6 | 5.6 | | | | | | | | | | | | | | | | | 22.4 |
| .55 | | | | | | 5.6 | 2.8 | 2.8 | | 2.8 | 2.8 | | | | | | | | | 5.6 |
| .60 | | | | | | 2.8 | 2.6 | 2.6 | 10.5 | | | | | | | | | | | 18.3 |
| 123.37 | 86.2 | 12.3 | | 4.9 | 2.5 | 7.4 | 2.5 | 2.5 | 2.5 | 5.0 | | | | | | | | | | 125.8 |
| .42 | 2.8 | | | 2.8 | 5.5 | 5.6 | 5.5 | | | | | | | | | | | | | 22.2 |
| .50 | | | | | | | | | | | | | | | | | | | | 2.8 |
| 127.34 | 21.9 | 22.0 | 46.3 | 100.1 | 207.4 | 156.2 | 90.3 | 65.9 | 56.1 | 22.0 | 14.6 | 7.3 | | | | | | | | 810.1 |
| .40 | | 11.0 | 43.7 | 120.2 | 109.2 | 54.6 | 65.6 | 71.0 | 43.7 | 32.7 | 27.3 | 5.5 | 16.4 | | | | | | | 600.9 |
| 130.30 | 71.7 | 4.8 | 9.6 | 4.8 | 4.8 | 4.8 | | | | | | | | | | | | | | 100.5 |
| .35 | | 5.3 | 5.3 | 15.8 | 5.3 | 21.1 | 5.3 | | | | | | | | | | | | | 58.1 |
| .40 | | 28.9 | 52.1 | 46.2 | 46.2 | 92.5 | 80.9 | 23.1 | 17.4 | 11.6 | 5.8 | 5.8 | | | | | | | 5.8 | 370.1 |
| .50 | | | | 5.7 | | | | | | | | | | | | | | | | 5.7 |
| 133.25 | | 21.1 | 128.9 | 381.3 | 349.8 | 239.3 | 76.3 | 13.1 | | | | | | | | | | | | 1,212.4 |
| 137.30 | | | | | 2.8 | 2.8 | | | | | | | | | | | | | | 5.6 |
| Total | 654.2 | 1480.1 | 1355.7 | 2413.0 | 1716.6 | 997.9 | 712.6 | 450.9 | 348.5 | 189.4 | 81.3 | 32.1 | 34.4 | 13.3 | 10.7 | | | | | 2,710,499.2 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | | Total | |
|--------------|---------------------------------|--------|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | | 23.25 |
| Cruise 5702: | | | | | | | | | | | | | | | | | | | |
| 80.51 | | | 2.5 | 7.4 | 12.4 | 49.4 | 64.2 | 32.1 | 7.4 | 7.4 | 5.0 | | | | | | | | 187.8 |
| .55 | | 3.4 | 6.8 | 13.6 | 16.9 | 190.0 | 47.5 | 47.5 | 95.0 | | | | | | | | | | 380.0 |
| .60 | | | | | | 10.2 | 3.4 | 3.4 | 6.8 | | | | | | | | | | 64.5 |
| 83.40 | 31.9 | 29.0 | 46.4 | 37.7 | 34.8 | 40.6 | 75.4 | 52.2 | 26.1 | 20.3 | 5.8 | 5.8 | | | | | | | 406.0 |
| .43 | 12.0 | 24.0 | 24.0 | 23.9 | 12.0 | 12.0 | 12.0 | 12.0 | 12.0 | | | | | | | | | | 119.9 |
| .51 | 54.2 | 90.4 | 90.4 | 126.5 | 18.1 | 54.3 | 18.1 | 18.1 | | | | | | | | | | | 470.1 |
| .52 | 41.1 | 41.1 | 30.8 | 10.3 | 41.1 | | | | | | | | | | | | | | 164.4 |
| 86.46 | 32.8 | 65.5 | 49.1 | 32.8 | 32.8 | 49.1 | 16.4 | | | | | | | | | | | | 245.7 |
| 87.36 | 90.8 | 288.3 | 165.6 | 128.2 | 165.5 | 154.8 | 133.5 | 144.2 | 48.0 | 37.4 | 21.4 | | | | | | | | 1,377.7 |
| .38 | 1236.3 | 2593.6 | 739.3 | 351.5 | 303.0 | 278.8 | 242.4 | 181.8 | 97.0 | 36.3 | 12.1 | 12.1 | | | | | | | 6,084.2 |
| .40 | 1689.1 | 2080.8 | 2252.1 | 1003.7 | 832.3 | 465.1 | 367.2 | 98.0 | 73.5 | 24.5 | | | | | | | | | 8,886.3 |
| .50 | | 6.0 | | | 3.0 | 3.0 | 3.0 | | | | | | | | | | | | 15.0 |
| .55 | 20.3 | 20.3 | | | | 20.3 | | | | | | | | | | | | | 60.9 |
| 90.28 | 11.0 | 60.3 | 54.8 | 60.3 | 32.8 | 27.4 | 27.4 | 5.5 | | | | | | | | | | | 279.5 |
| .30 | | 25.0 | 25.0 | | | 25.0 | | | | | | | | | | | | | 75.0 |
| .37 | | | 41.9 | | | | | | | | | | | | | | | | 41.9 |
| .55 | 71.0 | 71.1 | 47.4 | | 59.2 | 23.7 | 11.8 | 59.2 | 23.6 | 23.7 | 11.8 | | | | | | | | 402.5 |
| .60 | 2.9 | | | | 5.8 | | 14.5 | 23.3 | 14.5 | 5.8 | | | | | | | | | 66.8 |
| 93.27 | 516.7 | 424.8 | 309.9 | 126.3 | 91.8 | 114.8 | 11.5 | 23.0 | | | | | | | | | | | 1,618.8 |
| .30 | 26.7 | 53.4 | 21.3 | 21.3 | 21.4 | 5.3 | | 5.3 | | | | | | | | | | | 154.7 |
| .40 | 110.8 | 487.5 | 265.9 | 44.3 | | | | | | | | | | | | | | | 908.5 |
| .50 | 630.3 | 259.6 | 74.1 | 86.5 | 37.1 | 24.7 | 24.7 | 24.7 | 12.4 | | | | | | | | | | 1,149.4 |
| .60 | 40.3 | 67.2 | 13.4 | | | | | | | | | | | | | | | | 120.9 |
| 97.30 | 10.6 | 10.7 | 8.0 | 18.7 | 32.0 | 16.0 | 8.0 | 2.7 | | | | | | | | | | | 106.7 |
| .40 | 6.2 | 6.2 | 30.9 | 30.9 | 24.8 | 24.7 | 6.2 | 6.2 | 6.2 | | | | | | | | | | 142.3 |
| .60 | | | 51.8 | 77.7 | 25.9 | 25.9 | 25.9 | 51.8 | 51.8 | 51.8 | 25.9 | | | | | | | | 336.7 |
| .70 | | | | | | 2.9 | 2.9 | | | | 2.9 | | | | | | | | 5.8 |
| 100.29 | | | 17.9 | 44.7 | 17.9 | 12.0 | 3.0 | | | | | | | | | | | | 95.5 |
| .33 | | 112.0 | 89.6 | 22.4 | 44.8 | 11.2 | 11.2 | | | | | | | | | | | | 291.2 |
| .40 | 380.8 | 224.0 | 179.2 | 89.6 | 67.2 | 44.8 | 22.4 | | | | | | | | | | | | 1,008.0 |
| .50 | | | | | | | | | | 44.5 | | | | | | | | | 44.5 |
| .60 | | | | | | 6.8 | 3.4 | 13.6 | | | 3.4 | | | | | | | | 27.2 |
| 103.30 | | | 27.2 | 81.6 | 27.2 | 108.8 | 54.4 | 27.2 | 27.2 | | | | | | | | | | 353.6 |
| .35 | 27.7 | 166.1 | | 83.0 | 55.4 | 27.7 | 27.7 | 55.4 | | | | | | | | | | | 443.0 |
| .38 | 52.5 | 111.5 | 78.7 | | 6.6 | 6.6 | 19.7 | | | | | | | | | | | | 276.6 |
| .40 | | | 6.3 | | 12.6 | | | 6.3 | | | | | | | | | | | 25.2 |
| .50 | 14.4 | 14.4 | | | | 7.2 | 7.2 | | | | 7.2 | | | | | | | | 57.6 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Total | |
|-----------------------|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | | 23.25 |
| Cruise 5702 (cont'd): | | | | | | | | | | | | | | | | | | |
| 107.32* | 23.8 | 23.8 | 23.8 | 47.6 | 11.9 | 59.6 | 11.9 | 11.9 | 11.9 | 11.9 | | | | | | | | 202.4 |
| .35* | | 38.0 | 27.2 | | | | | | | | | | | | | | | 65.2 |
| .60 | | | | | | | | | | | 5.6 | | | | | | | 5.6 |
| 110.33 | | 94.0 | 177.5 | 292.3 | 177.4 | 135.7 | 219.3 | 93.9 | 82.7 | 41.7 | 41.8 | 10.4 | | | 10.4 | | | 1,357.1 |
| .35 | 12.4 | 12.4 | 49.5 | 49.5 | 24.7 | 24.8 | 61.8 | 37.1 | 24.7 | 37.1 | 24.7 | | | 12.4 | | | | 371.1 |
| .40 | 2.8 | 69.7 | 16.8 | 14.0 | 36.2 | 67.0 | 44.7 | 36.2 | 30.7 | 2.8 | 14.0 | | | | | | | 934.9 |
| .50 | | | 2.7 | 2.7 | | | | | | | | | | | | | | 5.4 |
| 113.30 | 5.1 | 10.2 | 20.5 | 35.9 | 51.2 | 66.3 | 51.2 | 15.3 | 15.3 | 5.1 | | 5.1 | | | | | | 271.2 |
| .35 | | 20.0 | 85.0 | 110.0 | 80.0 | 15.0 | 30.0 | 5.0 | | | | | | | | | | 325.0 |
| .40 | 10.1 | 55.4 | 58.0 | 50.4 | 37.8 | 32.8 | 5.0 | 2.5 | | | | | | | | | | 252.0 |
| 117.26 | | 56.6 | 18.9 | 28.3 | 132.1 | 141.8 | 188.8 | 198.2 | 169.9 | 84.9 | 37.8 | 47.2 | 37.7 | 9.4 | 16.9 | | | 1,170.3 |
| .30 | | | 133.7 | 100.3 | 234.1 | 200.6 | 588.5 | 588.4 | 367.9 | 267.6 | 167.2 | | 33.4 | 33.4 | 33.4 | | | 2,708.5 |
| .35 | | | 24.6 | 24.6 | 98.5 | | | | | | | | | | | | | 147.7 |
| .40 | | 9.2 | | 2.3 | 6.9 | 2.3 | | | 4.6 | | | 2.3 | | | | | | 27.6 |
| .50 | 2.8 | 28.1 | 2.8 | 8.4 | | | 2.8 | | | | | | | | | | | 44.9 |
| 120.25 | 23.8 | 12.0 | 89.4 | 53.7 | 71.5 | 89.4 | 77.5 | 53.6 | 29.8 | 29.8 | | 6.0 | | 6.0 | | | | 542.5 |
| .30 | | | 23.4 | 152.3 | 246.1 | 187.5 | 328.2 | 281.3 | 187.6 | 46.8 | 35.2 | 11.7 | 11.7 | | | | | 1,511.8 |
| .35 | | | | 19.5 | 4.9 | 4.9 | | 4.9 | | | | 4.9 | | | | | | 39.1 |
| .40 | | | 7.1 | 7.1 | 7.1 | | 7.1 | 7.1 | | 2.4 | 4.8 | | | | | | | 42.7 |
| .45 | | 13.9 | 44.3 | 8.3 | 5.5 | 8.3 | 2.8 | 27.7 | 11.1 | 2.8 | 2.8 | | | | | | | 127.5 |
| 123.37 | 22.1 | 25.5 | 13.6 | 10.2 | 3.4 | 1.7 | 6.8 | 5.1 | 5.1 | 11.9 | 10.2 | 1.7 | 1.7 | | | | | 119.0 |
| .42 | | | 13.4 | 13.3 | 10.6 | 8.0 | 8.0 | 2.7 | 2.7 | | | | | | | | | 58.7 |
| .50 | | | 19.1 | 4.8 | 4.8 | | 4.8 | | | | | | | | | | | 33.5 |
| .55 | | | | 9.0 | 4.5 | 9.1 | 54.5 | 49.9 | 9.0 | 4.5 | 4.5 | 9.0 | | | | | | 154.0 |
| .60 | | | | | | | | | | | | | | | | | 4.4 | 4.4 |
| 127.34 | 34.1 | 29.0 | 7.9 | 5.2 | 2.6 | 2.6 | 2.6 | 7.9 | 5.3 | | | | | | | | | 94.6 |
| .40 | | 7.0 | 14.0 | | | | 7.0 | | | | | | | | | | | 28.0 |
| .45 | | 10.0 | 10.0 | 50.2 | 80.3 | 100.4 | 120.4 | 130.6 | 70.3 | 30.1 | | | | | | | | 602.3 |
| .55 | | | | | | | 3.5 | | | | | | | | | | | 3.5 |
| .60 | | | | | | | 2.7 | 2.7 | | | | | | | | | | 5.4 |
| 130.30 | 25.6 | 281.6 | 748.8 | 614.4 | 153.6 | 44.8 | 25.6 | 12.8 | 6.4 | 19.2 | 6.4 | 6.4 | | | | | | 1,945.6 |
| .35 | | 21.8 | 34.9 | 30.5 | 21.8 | 52.3 | 65.4 | 52.4 | 43.6 | 39.3 | 34.8 | 34.9 | 13.1 | 8.8 | | | | 453.6 |
| .40 | | 5.2 | 13.0 | 13.0 | 15.6 | 15.6 | 5.2 | 7.8 | 2.6 | 2.6 | 2.6 | | | | | | | 80.6 |
| .50 | | | 6.0 | | | | | | | | | | | | | | | 6.0 |

* - Samples combined. Values adjusted accordingly.

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Total | | | | |
|-----------------------|---------------------------------|---------|--------|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | | 17.25 | 19.25 | 21.25 | 23.25 |
| Cruise 5702 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 133.25 | 242.9 | 2130.7 | 2252.2 | 1236.5 | 452.7 | 353.3 | 88.4 | 11.0 | 44.1 | 11.0 | 22.1 | 33.1 | | | | | | | 6,878.0 |
| .30 | 19.0 | 151.7 | 720.5 | 274.9 | 113.7 | 85.3 | 19.0 | 9.5 | 38.0 | 9.5 | 9.5 | | | | | | | | 1,469.6 |
| .50 | | | | | | | 3.2 | | | | | | | | | | | | 3.2 |
| 137.23 | | | 4.3 | | 8.6 | 8.5 | | | 4.3 | | | | | | | | | | 25.7 |
| .30 | | | | | | | | 10.0 | | | | | | | | | | | 10.0 |
| .35 | | | 9.8 | 24.6 | 19.6 | 49.2 | 34.5 | 29.6 | 4.9 | 9.8 | 9.8 | 9.8 | 4.9 | | | | | | 206.5 |
| .40 | | | 2.2 | | 13.2 | 2.2 | | | | | | | | | | | | | 17.6 |
| .50 | | | | | 2.4 | | | | | | | | | | | | | | 2.4 |
| 140.35 | | 3.7 | | | | | | | | | | | | | | | | | 3.7 |
| .40 | | | | | | | | | 2.8 | | | | | | | | | | 2.8 |
| .50 | | | | | | | | | 3.4 | | | | | | | | | | 3.4 |
| Total | 5518.1 | 9382.9 | 4157.8 | 3274.3 | 1663.1 | 918.0 | 558.5 | 213.4 | 172.2 | 85.0 | 71.5 | | | | | | | | 4.4 49,260.8 |
| | | 10420.7 | 5775.7 | 3546.0 | 2499.2 | | | | | | | | | | | | | | |
| Cruise 5703: | | | | | | | | | | | | | | | | | | | |
| 80.51 | | | | | | | 3.0 | | | | | | | | | | | | 3.0 |
| 82.47 | | 2.8 | | | | | | | | | | | | | | | | | 2.8 |
| 83.40 | | | | | | | 2.0 | | | | | | | | | | | | 2.0 |
| .43 | | | | | | 2.5 | | | | | | | | | | | | | 2.5 |
| .48 | | | | | 2.7 | 5.4 | 2.7 | 2.7 | 2.7 | | | | | | | | | | 10.0 |
| .51 | | | | | 2.7 | 5.4 | 2.7 | 2.7 | 2.7 | | | | | | | | | | 16.2 |
| 87.36 | | 6.0 | 12.0 | 24.1 | 84.3 | 48.2 | 54.2 | 30.1 | 30.1 | 12.0 | | | | | | | | | 301.0 |
| .40 | 11.0 | 22.1 | 66.2 | 55.2 | 143.6 | 154.6 | 187.7 | 132.5 | 33.1 | 22.1 | 44.2 | | | | | | | | 927.5 |
| .45 | 6.2 | 25.0 | | | 6.2 | 24.9 | | 6.2 | | | | | | | | | | | 68.5 |
| .50 | | | | | | | | 23.4 | 11.7 | | | | | | | | | | 35.1 |
| .70 | | | | | 82.4 | 94.1 | 199.9 | 117.6 | 82.3 | | | | | | | | | | 588.1 |
| 90.28 | 65.2 | 229.6 | 204.7 | 101.2 | 31.5 | 15.8 | 24.8 | 15.8 | 18.0 | 11.3 | 2.2 | | | | | | | | 720.1 |
| .30 | 73.3 | 60.0 | | 13.3 | | 13.4 | 46.6 | 13.3 | | | 13.3 | | | | | | | | 233.2 |
| .37 | 96.3 | 108.4 | 84.3 | 108.4 | 216.7 | 156.5 | 84.3 | 24.0 | | | | | | | | | | | 878.9 |
| .45 | 11.7 | 17.5 | 23.4 | 29.2 | 37.9 | 20.5 | 32.1 | | | | | | | | | | | | 172.3 |
| .60 | 34.2 | 17.1 | 5.7 | | 11.4 | 17.1 | 5.7 | | | | 5.7 | | | | | | | | 96.9 |
| .70 | | | | | 33.6 | 156.8 | 246.4 | 257.6 | 134.4 | | | | | | | | | | 828.8 |
| .80 | | | 22.8 | 5.7 | 74.4 | 97.3 | 108.7 | 28.6 | 5.7 | | | | | | | | | | 343.2 |
| 93.27 | 115.5 | 35.5 | 11.9 | 14.8 | 32.6 | 68.0 | 79.9 | 38.4 | 5.9 | 3.0 | | | | | | | | | 405.5 |
| .30 | 133.7 | 86.1 | 14.9 | 5.9 | 3.0 | 3.0 | 3.0 | 3.0 | | | | | | | | | | | 252.6 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Total | | | |
|-----------------------|---------------------------------|-------|-------|-------|-------|-------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | | 21.25 | 23.25 | Dis. |
| Cruise 5703 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 93.40 | 32.3 | 25.8 | 3.2 | 3.2 | 6.5 | 22.7 | 12.9 | 19.4 | | | | | | | | | | | | 129.2 |
| .50 | 12.0 | 18.0 | 72.3 | 72.2 | 72.2 | 24.0 | 48.1 | 90.3 | 42.2 | 6.0 | 6.0 | | | | | | | | | 463.3 |
| .60 | | | | 16.2 | 16.2 | 38.9 | 58.4 | 74.5 | 22.7 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | | | | | | 239.7 |
| 97.30 | 65.8 | 222.6 | 25.3 | 10.1 | 30.3 | | 10.2 | | 5.1 | | | | | | | | | | | 369.4 |
| .40 | 99.2 | 142.6 | 52.7 | 18.6 | 3.1 | | | | | | | | | | | | | | | 316.2 |
| .50 | | 22.8 | | | | | | 5.7 | | | | | | | | | | | | 34.2 |
| 100.29 | | | | 7.0 | 3.5 | 10.5 | | 3.5 | | | | | | | | | | | | 34.2 |
| .33 | | | | | | 6.8 | 27.4 | 17.1 | 24.0 | 30.8 | 6.8 | 6.8 | 3.4 | | | | | | | 28.0 |
| .40 | | | 3.1 | | | | 6.1 | | 6.2 | 6.1 | 3.1 | 6.1 | | | | | | | | 123.1 |
| 103.30 | 2.9 | 26.3 | 2.9 | 2.9 | 5.9 | 8.8 | | 2.9 | 5.8 | | | | | | | | | | | 30.7 |
| .35 | 2.9 | 5.8 | | | | | | | | | | | | | | | | | | 58.4 |
| 107.32 | | 26.6 | 6.6 | 33.2 | 13.3 | | 3.3 | | | | | | | | | | | | | 8.7 |
| 110.33 | | | | | | 4.5 | | | | | | | | | | | | | | 83.0 |
| .35 | | | | 14.0 | 35.1 | 10.5 | 10.5 | 7.0 | | | | | | | | | | | | 105.1 |
| .40 | | | | | | | | | | | | | | | | | | | | 3.3 |
| 113.30 | 5.8 | 5.8 | 11.6 | 5.8 | 5.8 | | | 11.6 | 11.6 | 23.2 | | | | | | | | | | 87.0 |
| .35 | | 12.7 | 50.7 | 12.7 | 12.7 | 50.8 | | 12.7 | 12.7 | 12.7 | | | | | | | | | | 190.4 |
| .40 | | | | 7.0 | 14.0 | 7.0 | | | | | | | | | | | | | | 28.0 |
| .50 | | | | 3.3 | | | | | | | | | | | | | | | | 3.3 |
| .60 | | | | | | | | | | | | | | | | | | | | 3.2 |
| 117.30 | | | | | 3.2 | | | | | | | | | | | | | | | 3.2 |
| .35 | 82.8 | 110.4 | 262.2 | 386.4 | 372.6 | 124.2 | | 13.8 | 13.8 | | | | | | | | | | | 160.3 |
| .40 | 3.0 | 6.0 | 17.9 | 20.9 | 9.0 | 3.0 | 3.0 | | | | | | | | | | | | | 1,366.2 |
| .50 | | | | 18.7 | 18.7 | 18.7 | 12.4 | 12.4 | 6.2 | | | | | | | | | | | 3.0 |
| .60 | | | | | | | 3.1 | | | | | | | | | | | | | 68.8 |
| .70 | 3.2 | | | 6.1 | | | | | | | | | | | | | | | | 87.1 |
| 118.39 | 95.6 | 57.3 | 28.7 | 38.3 | 9.6 | | | | | | | | | | | | | | | 12.3 |
| 119.33 | 45.3 | 13.0 | 110.2 | 123.1 | 64.8 | 13.0 | | | | | | | | | | | | | | 6.4 |
| 120.25 | 429.1 | 862.1 | 429.1 | 65.5 | | | | 6.0 | 6.5 | | | | | | | | | | | 239.1 |
| .30 | 48.2 | 12.0 | 36.1 | 36.1 | | | 12.0 | 12.0 | | | | | | | | | | | | 375.9 |
| .60 | 11.4 | | | | | | | | | | | | | | | | | | | 1,811.8 |
| 123.37 | | | | | | | | | | | | | | | | | | | | 156.4 |
| 127.34 | 76.3 | 18.3 | | 3.0 | | | 3.0 | | 3.2 | | | | | | | | | | | 11.4 |
| .40 | | 7.5 | | 22.5 | 22.5 | 15.0 | 15.0 | 3.0 | 9.1 | 3.0 | | | | | | | | | | 6.4 |
| .45 | 54.8 | 21.9 | 54.8 | 54.8 | 54.8 | | | | 15.0 | 7.5 | | | | | | | | | | 112.7 |
| .50 | | | 6.3 | | 31.5 | 37.8 | 6.3 | | | | | | | | | | | | | 112.5 |
| | | | | | | | | | 22.0 | | | | | | | | | | | 241.2 |
| | | | | | | | | | | | | | | | | | | | | 81.9 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | Total | | | | |
|-----------------------|---------------------------------|--------|--------|--------|--------|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|----------|---------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | | 19.25 | 21.25 | 23.25 | Dts. |
| Cruise 5703 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 130.30 | | | | | | | 10.7 | | | | | | 21.4 | 10.7 | | | | | | 42.8 |
| .40 | | | | | | | | 2.7 | | | | | | | | | | | | 2.7 |
| .50 | | | | | | 3.4 | | | 3.4 | | | | | | | | | | | 6.8 |
| 133.25 | | | | 20.5 | 30.7 | 20.4 | 30.7 | | | | | | | | | | | | | 102.3 |
| .30 | | 10.9 | 21.8 | 21.8 | 54.6 | 10.9 | 10.9 | 10.9 | 10.9 | | | | | | | | | | | 152.7 |
| .35 | 29.0 | 35.4 | 9.7 | | | | | | | | | | | | | | | | | 74.1 |
| .40 | | | | | 2.9 | 2.9 | | | | | | | | | | | | | | 5.8 |
| 137.23 | | | | | 8.6 | 2.9 | | | | | | | | | | | | | | 11.5 |
| .30 | | | | 2.8 | 2.8 | | | | | | | | | | | | | | | 5.6 |
| .35 | 11.7 | 99.3 | 332.9 | 99.3 | 5.8 | | | | | 5.8 | | | | | | | | | | 554.8 |
| .40 | 17.4 | 46.6 | 58.2 | 66.9 | 34.9 | 2.9 | 2.9 | 2.9 | | | | | | | | | | | | 232.7 |
| .50 | | | | 3.2 | | | | | | | | | | | | | | | | 3.2 |
| 140.30 | | | | | 2.7 | 5.4 | 5.4 | 2.7 | | 5.4 | 5.4 | 2.7 | | 2.7 | | | | | | 18.9 |
| .35 | | | | | 5.5 | 8.3 | 13.7 | 5.6 | 5.5 | 2.8 | | | | | | | | | | 41.4 |
| 143.26 | | | | | | | 2.9 | | | | | | | | | | | | | 5.8 |
| .30 | | | | | | | | | | | | | | | | | | | 3.3 | 3.3 |
| .35 | | | | | | | | | | | | | 9.8 | | | | | | | 9.8 |
| Total | 1603.6 | 2440.7 | 2006.7 | 1591.3 | 1805.2 | 1438.9 | 1390.0 | 994.0 | 498.0 | 202.4 | 120.3 | 134.6 | 17.6 | 2.7 | 42.2 | 21.9 | 8.8 | 9.4 | 14,328.3 | |
| Cruise 5704: | | | | | | | | | | | | | | | | | | | | |
| 82.47 | | 5.4 | 2.7 | 8.1 | 8.1 | 10.8 | 16.2 | 10.8 | | 8.1 | 2.7 | | | | | | | | | 72.9 |
| 83.40 | 78.8 | 7.9 | 23.6 | 94.5 | | | | | | | | | | | | | | | | 204.8 |
| .43 | | | | 13.2 | 13.2 | 6.6 | | | | | | | | | | | | | | 33.0 |
| .51 | | | | 19.3 | 9.6 | 24.1 | | | | 14.4 | 19.3 | | | | | | | | | 96.3 |
| 86.46 | | 39.4 | 88.6 | 118.0 | 98.4 | 78.7 | 34.5 | 24.6 | 4.9 | | | | | | | | | | | 487.1 |
| 87.36 | 20.7 | 132.5 | 385.0 | 285.6 | 248.4 | 120.1 | 24.9 | 20.7 | 8.2 | | | | | | | | | | | 1,246.1 |
| .40 | | 9.7 | 48.6 | 68.1 | 136.1 | 68.1 | 106.9 | 48.6 | 38.8 | 9.7 | 19.4 | | | | | | | | | 554.0 |
| .50 | | 55.0 | 220.1 | 440.3 | 522.9 | 440.3 | 385.2 | 275.2 | 82.5 | | 27.5 | | | | | | | | | 2,449.0 |
| 90.28 | 554.6 | 1303.5 | 474.0 | 331.8 | 132.7 | 85.3 | 71.1 | 14.2 | 4.7 | | | | | | | | | | | 2,971.9 |
| .30 | 70.5 | 274.4 | 62.7 | 23.5 | 23.5 | 7.8 | 23.5 | 47.0 | 23.5 | 23.5 | | | | | | | | | | 579.9 |
| .37 | 260.4 | 111.6 | 74.4 | 161.2 | 111.6 | 74.4 | 74.4 | 12.4 | 24.8 | | | | | | | | | | | 905.2 |
| .45 | | | | 13.2 | 79.0 | 144.8 | 144.8 | 210.5 | 79.0 | 92.1 | 13.2 | | | | | | | | | 776.6 |
| .55 | 13.6 | | | | | | | 13.6 | | | | | | | | | | | | 40.8 |
| .60 | | | | | | | | 2.9 | | | | | | | | | | | | 2.9 |
| .80 | | | | | | | | 12.4 | | | | | | | | | | | | 12.4 |
| .90 | | | | | | 13.2 | 79.5 | 53.0 | 66.2 | 13.2 | | | | | 6.6 | | | | | 231.7 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Dis. | Total | | |
|-----------------------|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|-------|---------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | | | 17.25 | 19.25 |
| Cruise 5704 (cont'd): | | | | | | | | | | | | | | | | | | |
| 93.27 | 62.8 | 270.0 | 389.3 | 219.8 | 138.1 | 43.9 | 18.9 | 25.1 | 12.6 | | | | | | | | | 1,180.5 |
| .30 | 477.5 | 300.1 | 272.8 | 122.8 | 81.8 | 95.5 | 27.2 | 13.6 | 27.2 | | | | | | | | | 1,418.5 |
| .35 | 73.5 | 110.1 | 55.1 | 73.4 | 113.7 | 84.4 | 44.1 | 44.1 | 33.1 | 11.0 | | | | | | | | 642.5 |
| .40 | 100.0 | 40.0 | 20.0 | 80.0 | 200.0 | 280.0 | 180.0 | 140.0 | 100.0 | 40.0 | 20.0 | | | | | | | 1,200.0 |
| .45 | 14.9 | 12.4 | 49.4 | 160.7 | 111.2 | 185.4 | 284.3 | 173.0 | 160.7 | 61.8 | 24.8 | | | | 12.4 | | | 1,236.1 |
| .50 | | | 14.9 | 14.9 | 14.9 | 44.7 | 44.7 | 44.8 | 14.9 | 59.6 | 14.9 | | | | | | | 268.3 |
| .55 | | | 9.6 | 33.6 | 14.4 | 28.8 | 19.2 | 33.6 | 38.4 | 4.8 | 4.8 | | | 4.8 | | | | 192.0 |
| .60 | | | | 34.2 | 11.4 | | | | 11.4 | | | | | | | | | 57.0 |
| .90 | | | | | | | | 6.4 | 12.7 | | | | | | | | | 19.1 |
| 97.30 | 55.0 | 101.0 | 73.4 | 27.5 | 42.8 | 33.7 | 15.3 | 18.3 | | | | | | | | | | 367.0 |
| .32 | 203.8 | 17.0 | | 5.7 | 17.0 | 50.9 | 17.0 | 22.6 | 5.7 | 5.7 | | | | | | | | 345.4 |
| .40 | | | | | | | 6.1 | | 6.0 | 6.0 | | | | | | | | 12.0 |
| .45 | | | | | | | | | 12.2 | | | | | | | | | 18.3 |
| .60 | | | | | | | | | | | | | | 12.8 | | | | 38.4 |
| .70 | | | | | | | | | | | | | | | | | | 2.8 |
| .90 | | | | | | | | 2.8 | | | | | | | | | | 23.0 |
| 100.29 | 5.2 | 5.2 | 10.5 | 10.5 | 15.7 | 20.9 | 94.3 | 78.6 | 57.6 | 15.7 | | 7.7 | | 5.2 | 5.2 | | | 324.6 |
| .33 | | 26.1 | | 5.2 | 5.2 | 31.4 | 62.6 | 26.1 | 31.3 | 5.2 | | | | | | | | 193.1 |
| .40 | 4.9 | 19.6 | 4.9 | | | | | 4.9 | 14.7 | 14.7 | | | | | | | | 63.7 |
| .45 | | | 6.3 | 9.5 | 3.2 | 6.4 | | 3.2 | | 3.2 | 3.2 | | | | | | | 35.0 |
| .50 | | | | | | | | | | | | | | | | | | 3.1 |
| .60 | | | | | | 4.8 | | | | 4.8 | 19.2 | 4.8 | | | | | | 33.6 |
| .70 | | | | | | | | | | 1.8 | | | | | | | | 1.8 |
| 103.30 | 2.9 | 1.4 | 26.6 | 61.6 | 40.0 | 7.1 | 10.0 | 17.2 | 24.4 | 62.9 | 60.1 | 37.2 | 12.9 | 8.6 | 4.3 | | | 379.1 |
| .40 | | | | | | | | | | 2.8 | 2.8 | | 2.8 | 2.8 | 2.8 | | | 8.4 |
| .90 | | | | | | | | 3.1 | | | | | | | | | | 3.1 |
| 107.32 | | | 6.4 | 6.4 | | | 12.8 | 6.4 | 31.9 | 44.7 | 12.8 | | | | | | | 121.4 |
| .35 | 3.2 | | 22.8 | 48.8 | 58.4 | 22.7 | 13.0 | 22.8 | 16.2 | 16.3 | 9.7 | | | | | | | 233.9 |
| .40 | | | 10.0 | | | 3.3 | 3.3 | 3.3 | 3.3 | 3.3 | 3.3 | | | | | 3.3 | | 26.5 |
| .45 | | | | | | 2.8 | | | | | | | | | | | | 2.8 |
| 110.33 | | | | | | | | | 21.4 | | | | | | | | | 42.8 |
| .35 | | | | | 6.3 | 18.9 | 18.9 | 6.3 | 6.3 | 6.3 | | | 12.6 | | 12.6 | 6.3 | | 125.9 |
| .40 | | | | | | 14.1 | 14.1 | 7.1 | 7.1 | | | 35.3 | 28.2 | 21.2 | 7.1 | 35.4 | 7.1 | 176.8 |
| .45 | | | | | | 3.0 | | | | | | | | | | | | 3.0 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Dis. | Total | | |
|-----------------------|---------------------------------|--------|--------|--------|--------|--------|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|---------|----------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | | | 21.25 | 23.25 |
| Cruise 5704 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 113.30 | | | | 1.7 | | | 1.7 | 1.7 | 5.1 | 24.0 | 39.5 | 46.5 | 43.0 | 32.7 | 24.2 | 11.9 | | 232.0 | | |
| .35 | | | | | | | | | 4.0 | 8.0 | | | | | | | | 12.0 | | |
| 117.26 | | 8.5 | 2.8 | 8.5 | 8.5 | 2.8 | 5.7 | | | | 3.8 | | | | 3.8 | | | 36.8 | | |
| .30 | 3.8 | 3.8 | | 3.8 | | | | | | | | | 5.3 | 5.3 | | | | 19.0 | | |
| .60 | | | | | | | | | | | | | | | | | | 21.2 | | |
| 120.25 | | | | | 4.8 | 7.3 | | 2.4 | | | | | | | | | | 16.9 | | |
| .30 | | | 2.1 | | | | | | | | | | | | | | | 2.1 | | |
| .35 | | | 7.6 | | 15.2 | | | | | | | | | | | | | 22.8 | | |
| .50 | | | | | | | | | | | | | | | | | | 23.1 | | |
| 123.42 | 5.8 | 5.8 | | | | 5.8 | | | | | | | | | | | | 17.4 | | |
| .70 | | | | | | | | 2.1 | | | | | | | | | | 2.1 | | |
| 127.34 | | | | 1.6 | 1.6 | | 1.6 | 1.6 | | | 1.6 | | 1.6 | | 1.6 | | | 11.2 | | |
| .45 | 6.1 | 3.0 | | 3.0 | | 6.1 | | 6.1 | | | | | | | | | | 24.3 | | |
| .50 | | | | | 10.6 | | | | | | | | | | | | | 10.6 | | |
| 130.35 | 4.2 | 25.2 | | | | | | | | | | | | | | | | 29.4 | | |
| 133.30 | | 19.4 | | | | | | | | | | | | | | | | 19.4 | | |
| .35 | 3.0 | 8.9 | | 3.0 | | | | | | | | | | | | | | 14.9 | | |
| .40 | | 107.0 | | | | | | | | | | | | | | | | 107.0 | | |
| 137.23 | 39.6 | 6.6 | | | | | | | | | | | | | | | | 46.2 | | |
| .40 | | 38.1 | 19.0 | | | | | | | | | | | | | | | 57.1 | | |
| .45 | 9.6 | 14.4 | | | | | | | | | | | | | | | | 24.0 | | |
| 140.50 | 5.0 | 5.0 | | | | | | | | | | | | | | | | 10.0 | | |
| 143.26 | | | | | | | | | | | 2.6 | | | | | | | 5.2 | | |
| 147.20 | | | | | | | 2.6 | | | | 3.0 | | | | | | | 3.0 | | |
| Total | 2079.4 | 3088.0 | 2364.7 | 2468.6 | 2315.9 | 2024.5 | 1778.8 | 1520.0 | 980.8 | 670.3 | 340.0 | 214.8 | 122.2 | 63.7 | 120.2 | 30.2 | 37.3 | 9.3 | 3.1 | 20,231.8 |
| Cruise 5705: | | | | | | | | | | | | | | | | | | | | |
| 60.55 | | | | | | | | | | | | | | | | | | | 4.8 | |
| 63.52 | | | | | | | | | | | | | | | | | | | 4.4 | |
| 73.60 | | | | | 4.6 | | | | | | | | | | | | | | 4.6 | |
| 82.47 | | | | | | | | 37.3 | 27.9 | 46.6 | 28.0 | 9.3 | | 9.3 | | | | | 158.4 | |
| 83.40 | 2566.3 | 372.3 | 225.6 | 530.2 | 569.7 | 220.0 | 73.4 | 22.6 | 11.3 | 5.6 | 5.6 | 5.6 | 5.6 | | | | | | 4,613.8 | |
| .43 | | | | 50.4 | 60.5 | 80.6 | 20.2 | 30.3 | 10.1 | 10.1 | 10.1 | | | 10.1 | | | | | 282.4 | |
| .51 | | | 3.4 | | | | | 6.9 | | 3.4 | | | | | | | | | 13.7 | |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | | Dis. | Total |
|-----------------------|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|---------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | | |
| Cruise 5705 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 87.36 | 455.9 | 799.2 | 787.4 | 538.7 | 219.0 | 88.8 | 29.6 | 17.7 | | | | | | | | | | | 2,936.3 |
| .40 | 26.3 | 157.7 | 280.3 | 289.1 | 481.8 | 402.9 | 297.9 | 87.6 | 61.4 | 17.6 | 8.8 | | | | | | | | 2,111.4 |
| .45 | 13.9 | 111.3 | 236.7 | 501.2 | 473.3 | 167.0 | 97.5 | 41.7 | 13.9 | | | | | | | | | | 1,656.5 |
| .50 | 28 | | 5.5 | 14.6 | 7.4 | 11.0 | 1.8 | 1.8 | | | | | | | 1.8 | | | | 43.9 |
| 90.28 | 107.2 | 61.6 | 50.9 | 42.9 | 34.8 | 18.7 | 10.8 | 8.1 | 2.7 | | | | | | | | | | 337.7 |
| .30 | 226.9 | 85.1 | 43.5 | 45.4 | 35.9 | 32.1 | 11.3 | 1.9 | | | | | | | | | | | 487.8 |
| .37 | 50.7 | 101.4 | 44.4 | 38.0 | 34.8 | 19.0 | 6.4 | | 3.2 | | | | | | | | | | 297.9 |
| .45 | 81.3 | 16.3 | | 5.4 | | | | | | | | | | | | | | | 103.0 |
| .50 | | | 2.8 | 11.1 | 19.4 | 30.5 | | 2.8 | | | | | | | | | | | 66.6 |
| .65 | 28.2 | | | | | | | | | | | | | | | | | | 28.2 |
| .75 | | | | | | | | | | | | | | | | | | | |
| .85 | | | | | | | | | | | | | | | | | | | |
| 93.27 | 189.0 | 45.0 | | | 27.0 | 27.0 | | 36.0 | 9.0 | | | | | | 2.5 | | | | 6.6 |
| .30 | 158.3 | 227.3 | 184.8 | 194.9 | 115.7 | 69.0 | 58.9 | 52.8 | 24.4 | 6.1 | 4.0 | 2.0 | 2.0 | | | | | | 2.5 |
| .35 | 143.6 | 37.4 | 39.5 | 104.0 | 62.4 | 35.3 | 16.6 | 2.1 | 4.2 | | | | | | | | | | 333.0 |
| .45 | 10.5 | 47.3 | 42.1 | 28.9 | 18.4 | 7.9 | 2.6 | | | | | | | | | | | | 1,104.2 |
| .50 | | | 4.6 | 9.2 | | | | | | | | | | | | | | | 445.1 |
| 97.30 | | 13.4 | 49.1 | 75.8 | 49.1 | 44.6 | 49.1 | 26.8 | 4.5 | | | | | | | | | | 160.3 |
| .32 | | | | | 6.4 | 12.8 | | | | | | | | | | | | | 13.8 |
| .45 | | | | | | | | | | | | | | | | | | | 312.4 |
| .50 | | | | | | | | | | | | | | | | | | | 19.2 |
| 100.29 | | | 9.0 | 4.5 | 9.0 | 18.1 | 13.5 | 18.0 | 2.9 | | | | | | | | | | 5.2 |
| .33 | | | 34.0 | 110.7 | 153.4 | 136.4 | 119.3 | 102.2 | 8.5 | 42.6 | 8.5 | 17.0 | 17.0 | 8.5 | 17.0 | 8.5 | 8.5 | | 2.9 |
| .40 | | | | | | | | | | 3.7 | | | | | | | | | 72.1 |
| .50 | | | | | | | | | | | | | | | | | | | 792.1 |
| .55 | | | | | | | | | | | | | | | | | | | 3.7 |
| 103.30 | | | | 4.5 | 9.0 | 45.2 | 72.3 | 63.2 | 9.1 | 76.8 | 36.2 | 13.5 | 22.6 | 9.0 | 4.5 | 4.5 | | | 2.4 |
| .35 | | | | | | | | | | 3.3 | | | | | | | | | 9.1 |
| 107.32 | | | | | | | | | | 92.4 | 64.7 | 36.9 | 9.2 | 9.2 | | | | | 424.6 |
| 110.40 | | | | | | | | | | 23.4 | | | | | | | | | 3.3 |
| .45 | | | | | | | | | | | | | | | | | | | 443.3 |
| .60 | | | 6.0 | | 3.0 | | | | | | | 3.1 | | 6.2 | | | | | 23.4 |
| .65 | 5.3 | 5.3 | | | | | | | | | | | | | | | | | 9.3 |
| .70 | 13.2 | 5.3 | 2.6 | | | | | | | | | | | | | | | | 9.0 |
| .75 | | 28.9 | 5.8 | 2.9 | 2.9 | | | | | | | | | | | | | | 10.6 |
| | | | | | | | | | | | | | | | | | | | 21.1 |
| | | | | | | | | | | | | | | | | | | | 40.5 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Total | | | |
|-----------------------|---------------------------------|--------|--------|--------|--------|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|----------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | | 21.25 | 23.25 | Dis. |
| Cruise 5705 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 113.35 | 105.4 | 80.6 | | | | | | | | | | | | | | | | | | 186.0 |
| .40 | 835.7 | 73.2 | | | | | | | | | | | | | | | | | | 908.9 |
| .45 | 2.7 | | | | | | | | | | | | | | | | | | | 2.7 |
| .50 | 15.8 | 21.2 | 2.6 | | | | | | | | | | | | | | | | | 39.8 |
| .65 | 10.8 | | 2.7 | 2.7 | | | | | | | | | | | | | | | | 16.2 |
| .75 | 2.6 | | | | | | | | | | | | | | | | | | | 2.6 |
| 117.26 | | | 6.6 | 6.7 | 6.7 | 3.3 | 6.7 | | | | | | | | | | | | | 30.0 |
| .30 | 23.3 | 9.4 | | | 9.4 | 4.7 | 4.7 | | 4.7 | | | | | | | | | | | 51.5 |
| .35 | | | | | | 4.1 | 4.1 | | | | | | | | | | | | | 4.1 |
| .40 | 8.1 | | | | | | | | | | | | | | | | | | | 8.1 |
| .50 | | | 8.4 | 5.6 | 2.8 | 2.8 | 2.8 | | 2.8 | | | | | | | | | | | 22.4 |
| .55 | 449.9 | 73.0 | | | | | | | | | | | | | | | | | | 522.9 |
| .60 | | | | 6.2 | | | | | | | | | | | | | | | | 6.2 |
| 118.39 | 3.0 | | | | 12.0 | 12.1 | 3.0 | 6.0 | | | | | | | | | | | | 36.1 |
| 119.33 | | | 5.4 | 32.1 | 80.4 | 107.2 | 26.8 | 5.4 | | | | | | | | | | | | 257.3 |
| 120.25 | | 7.4 | 3.7 | | 7.4 | 3.7 | 3.7 | | | | | | | | | | | | | 25.9 |
| .30 | | | 33.5 | 22.4 | 55.8 | 11.2 | 22.3 | 11.2 | | | | | | | | | | | | 156.4 |
| .40 | 7.6 | 3.0 | | | | | | | | | | | | | | | | | | 10.6 |
| .45 | | 5.1 | | 7.6 | 2.5 | 2.5 | 2.5 | | | | | | | | | | | | | 20.2 |
| .50 | 2.6 | 2.6 | 2.6 | 20.8 | 15.6 | 7.8 | | | | | | | | | | | | | | 52.0 |
| .55 | | | 2.8 | 2.8 | 2.8 | 2.8 | | | | | | | | | | | | | | 8.4 |
| .60 | | | | 23.3 | 11.6 | 46.6 | 23.3 | 34.9 | | | | | | | | | | | | 139.7 |
| .65 | 36.4 | 66.7 | 24.2 | 36.3 | 54.5 | 30.3 | 12.2 | 6.1 | 6.1 | 6.1 | | | | | | | | | | 278.9 |
| .70 | | | | | 5.9 | | | | | | | | | | | | | | | 5.9 |
| 123.37 | | | | 7.9 | 7.8 | 19.6 | 9.8 | 2.0 | 2.0 | | | | | | | | | | | 49.1 |
| .42 | | | | 8.1 | 10.8 | 5.4 | 2.7 | | | | | | | | | | | | | 27.0 |
| 127.34 | 8.7 | 2.9 | | | | | | | | | | | | | | | | | | 14.5 |
| 130.30 | 47.1 | | | | | | | | | | | | | | | | | | | 47.1 |
| .40 | | | | | | | 4.4 | | | | | | | | | | | | | 4.4 |
| 133.25 | 9.3 | 37.3 | 18.6 | 28.0 | 18.6 | | | | | | | | | | | | | | | 111.8 |
| .30 | 37.1 | 39.4 | 14.0 | 4.6 | | | | | | | | | | | | | | | | 95.1 |
| .40 | | | | | 2.6 | | | | | | | | | | | | | | | 2.6 |
| .50 | 3.1 | | | | | | 3.1 | 3.1 | | | | | | | | | | | | 9.3 |
| 137.23 | | | 11.2 | 44.9 | 45.0 | 11.2 | 11.2 | | | | | | | | | | | | | 123.5 |
| .30 | 2.8 | 25.1 | 22.3 | 19.6 | 25.2 | 2.8 | 2.8 | | | | | | | | | | | | | 97.8 |
| Total | 5693.0 | 2577.7 | 2235.3 | 2843.4 | 2706.4 | 1782.6 | 1098.0 | 763.5 | 405.8 | 283.3 | 131.3 | 76.6 | 56.4 | 50.7 | 38.1 | 28.8 | 15.4 | 3.3 | 5.7 | 20,795.9 |
| | | | | | | | | | | | | | | | | | | | | 2.6* |

* - 2.6 at 24.5 mm.

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | Total | | | |
|--------------|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.76 | 13.75 | 14.75 | 15.75 | 17.25 | | 19.25 | 21.25 | 23.25 |
| Cruise 5706: | | | | | | | | | | | | | | | | | | | |
| 63.60 | | | | | | | | 13.0 | | | | | | | | | | | 13.0 |
| 67.50 | | | | | | | | | 3.3 | | | | | | | | | | 3.3 |
| 70.60 | | | | | | 6.3 | | | | | | | | | | | | | 6.3 |
| .80 | | | | | | | | | | 12.8 | | | | | | | | | 25.6 |
| 73.55 | | | | | | | | 15.4 | | 15.4 | 15.4 | | | | | | | | 46.2 |
| .60 | | | | | | | | | | | 12.1 | | | | | | | | 24.2 |
| .70 | | | | | | | | | | | | | | | | | | | 3.1 |
| 80.51 | | | | | | | | | | | | | | | | | | | 25.4 |
| .90 | | | | | | | | | | | | | | | | | | | 6.0 |
| 83.40 | 199.8 | 111.6 | 104.4 | 68.4 | 32.4 | 3.6 | | | | | | | | | | | | | 520.2 |
| .43 | 2.6 | 15.7 | 36.7 | 52.4 | 68.1 | 34.1 | 18.4 | 13.1 | 13.1 | 2.6 | | | | | | | | | 256.8 |
| .51 | 35.5 | 92.4 | 60.4 | 71.0 | 95.9 | 80.3 | 32.0 | 31.9 | 21.3 | 7.1 | | | | | | | | | 507.8 |
| .55 | | 3.4 | 10.1 | 23.6 | 37.1 | 23.6 | 16.8 | 6.8 | 10.1 | 3.4 | | | | | | | | | 134.9 |
| 87.35 | 500.3 | 473.2 | 121.7 | 121.7 | 54.0 | 81.1 | 40.5 | | | | | | | | | | | | 1,392.5 |
| .40 | 32.3 | 153.6 | 56.6 | 16.2 | | | | | | | | | | | | | | | 258.7 |
| .45 | 8.2 | 28.5 | 24.4 | | | | | | | | | | | | | | | | 134.6 |
| .50 | | | 17.3 | 23.1 | 17.4 | 52.0 | 34.6 | 69.3 | 63.6 | 46.2 | 17.4 | 5.8 | | | | | | | 346.7 |
| .55 | | | 11.4 | 5.7 | 22.8 | 11.4 | 11.4 | 22.7 | 39.8 | 68.2 | 68.1 | 39.8 | 22.7 | 11.4 | 28.5 | 5.7 | | | 369.8 |
| .60 | | | 15.2 | 45.8 | 198.6 | 313.3 | 198.6 | 114.6 | 45.8 | 15.3 | | | | | | | | | 954.8 |
| .65 | | | | | | 5.5 | | | | | | | | | | | | | 5.5 |
| .70 | | | | | | 13.0 | 13.0 | 13.0 | | 13.0 | | | | | | | | | 52.0 |
| .75 | | | | | | 6.0 | 5.9 | 8.9 | | | | | | | 3.0 | | | | 23.8 |
| .80 | | | | | | | 3.1 | | | | | | | | | | | | 6.2 |
| 90.28 | 762.3 | 571.7 | 254.1 | 55.6 | 87.3 | 31.7 | 31.7 | 31.7 | 7.9 | | | | | | | | | | 1,834.0 |
| .30 | 701.3 | 228.5 | 47.3 | 55.1 | 47.3 | 15.8 | 15.8 | | | | | | | | | | | | 1,111.1 |
| .37 | 446.6 | 459.4 | 191.4 | 308.2 | 382.8 | 165.8 | 38.3 | 38.3 | 25.6 | | | | | | | | | | 2,054.4 |
| .45 | 78.8 | 64.5 | 71.6 | 57.2 | 50.1 | 35.8 | 57.3 | 35.8 | 28.7 | 21.5 | 14.4 | 7.2 | | | | | | | 530.1 |
| .50 | | | 3.6 | 3.6 | | | | | 7.2 | | | | | | | | | | 14.4 |
| .55 | | | | | | | 7.0 | | | | | | | | | | | | 28.0 |
| .60 | | | | | | | 6.3 | 6.3 | 37.8 | 69.3 | 37.8 | | | 6.3 | 6.3 | | | | 226.8 |
| .65 | | | | | | | | | 3.1 | 3.1 | | | | 3.1 | | | | | 9.3 |
| .70 | | | | | | | | | | | | | | | | | | | 3.1* |
| .75 | | | | | | | | | | | | | | | | | | | 15.0 |
| .90 | | | | | | | | | | | | | | | | | | | 6.2 |
| | | | | | | | | | | | | | | | | | | | 15.0 |
| | | | | | | | | | | | | | | | | | | | 10.2 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Total | | |
|-----------------------|---------------------------------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | | 21.25 | 23.25 |
| Cruise 5706 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 93.27 | | | | 14.8 | 14.8 | 29.6 | 22.2 | 22.2 | 14.8 | 11.1 | | | | | | | | | 129.5 |
| .30 | | | 2.9 | | | | | | | | | | | | | | | | 2.9 |
| .35 | | | | | | | | 3.1 | | | | | | | | | | | 3.1 |
| .40 | 3.1 | 18.6 | | | | | | | | | | | | | | | | | 21.7 |
| .50 | | | 3.4 | | | | | | | | | | | | | | | | 3.4 |
| .55 | | 6.6 | 13.2 | 55.8 | 26.3 | | | | | | | | | | | | | | 101.9 |
| 97.55 | | | | 19.7 | 19.8 | 19.8 | 6.6 | | | | | | | | | | | | 65.9 |
| .60 | | | | | | | | | | | | | | | | | | | 3.0 |
| 100.29 | | | | | | | | | | 11.9 | | | | | | | | | 11.9 |
| .45 | | | | 3.0 | | | | | | | | | | | | | | | 3.0 |
| .55 | | | | 3.3 | | 3.3 | | | | | | | | | | | | | 3.0 |
| .60 | | | | 6.2 | | | | | | | | | | | | | | | 6.6 |
| 107.55 | | | | | | | | | | | | | | | | | | | 6.2 |
| .65 | | | | | | | | | | | | | | | | | | | 3.1 |
| 110.33 | | | | | | | | | | | | | | | | | | | 3.3 |
| .35 | | 12.4 | | | | | 3.2 | | | | | | | | | | | | 3.2 |
| .75 | | | | 9.7 | 3.2 | 16.2 | 3.1 | | | | | | | | | | | | 15.5 |
| .80 | | | | 5.9 | | | 22.7 | | | 5.9 | | | | | | | | | 55.0 |
| 113.30 | | | | 1.8 | 3.6 | | 1.8 | | | | | | | | | | | | 11.8 |
| .35 | 2.7 | | | | | | | | | | | | | | | | | | 7.2 |
| .40 | | | | 3.1 | 3.1 | | | | | | | | | | | | | | 2.7 |
| .45 | | | | 4.9 | | | | | | | | | | | | | | | 6.2 |
| .50 | | | 9.8 | | | | | | | | | | | | | | | | 3.2 |
| .55 | 2.9 | | | | | 2.9 | 4.9 | | | | | | | | | | | | 6.2 |
| .65 | | | | | | 2.4 | 2.9 | | | 5.8 | | | | | | | | | 3.2 |
| 117.26 | | | 2.4 | | | | | | | | | | | | | | | | 6.2 |
| .30 | 94.9 | 6.8 | 6.0 | 24.0 | 36.0 | 42.0 | 36.0 | | | | | | | | | | | | 144.0 |
| .35 | 2.4 | 2.4 | 20.4 | 27.2 | 13.6 | 13.6 | 27.1 | 27.2 | | 6.8 | | | | | | | | | 237.6 |
| .40 | 32.6 | 81.6 | 4.9 | 12.3 | 12.3 | 2.4 | 4.8 | | 4.9 | | | | | | | | | | 46.4 |
| .45 | | | 39.5 | 13.6 | | 6.8 | 2.8 | | 2.8 | 1.4 | | | | | | | | | 181.1 |
| .50 | | | | | | 1.3 | 1.3 | | 1.3 | | | | | | | | | | 3.9 |
| .55 | 5.2 | | | 3.1 | | | | | | | | | | | | | | | 3.1 |
| .60 | | | | | | | | | | | | | | | | | | | 7.8 |
| .65 | 2.4 | 2.4 | | 2.4 | | 2.4 | 2.4 | | | | | | | | | | | | 2.5 |
| .70 | | | | | | | | | | | | | | | | | | | 12.0 |
| 118.39 | 4.8 | | | | 2.5 | | | | | | | | | | | | | | 2.5 |
| | | | | | 4.8 | | | | | | | | | | | | | | 14.4 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Dis. | Total |
|---|---------------------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|---------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | | |
| Cruise 5706 (cont'd): | | | | | | | | | | | | | | | | | | |
| 119.33 | 5.5 | 2.7 | 5.5 | 13.7 | 19.2 | 16.5 | 8.2 | 2.7 | 5.5 | 2.7 | 2.7 | 2.7 | 2.7 | 2.7 | | | | 84.9 |
| 120.25 | | 32.8 | 68.0 | 128.5 | 189.0 | 131.0 | 68.1 | 40.4 | 25.2 | 17.7 | 2.5 | | | | | | | 723.4 |
| .30 | | 6.6 | 13.2 | 29.7 | 89.1 | 145.2 | 99.0 | 52.8 | 19.8 | 26.4 | 6.6 | | | | | | | 488.4 |
| .35 | | 10.3 | 34.2 | 109.4 | 355.6 | 820.8 | 776.3 | 352.3 | 153.9 | 188.1 | 102.6 | 44.5 | 17.1 | 6.8 | | | | 2,971.9 |
| .40 | | 1.8 | 1.8 | 3.6 | 3.6 | 1.8 | | | | | | | | | | | | 9.0 |
| .45 | | | | 3.2 | | | | | | | | | | | | | | 6.4 |
| .50 | | | | | | | | | 2.5 | | | | | | | | | 2.5 |
| .65 | | | 3.4 | | 3.4 | 3.4 | | | | | | | | | | | | 10.2 |
| .70 | | 8.6 | 2.9 | | | | | | | | | | | | | | | 11.5 |
| 123.37 | | | | | 4.3 | | | | | | | | | | | | | 6.5 |
| 130.30 | | | | | | 2.6 | | 6.2 | | | | | | 2.2 | | | | 2.6 |
| .35 | 6.2 | | | | | | | | | | | | | | | | | 12.4 |
| .40 | 28.1 | | 3.1 | | | | | | | | | | | | | | | 31.2 |
| 133.25 | | 12.8 | 204.8 | 307.2 | 384.0 | 448.0 | 230.4 | 25.6 | | | | | | | | | | 1,612.8 |
| .30 | | | | 13.6 | 13.6 | 13.6 | | | | | | | | | | | | 54.4 |
| .35 | | | | 9.5 | 19.0 | 28.5 | | 28.5 | | | | | | | | | | 85.5 |
| .40 | 2.7 | 2.7 | 16.0 | 10.6 | | 2.7 | 2.7 | 2.7 | | | | | | | | | | 40.1 |
| 137.23 | | | | 41.9 | 41.9 | 41.9 | | 41.9 | | | | | | | | | | 125.7 |
| .45 | 2.1 | | | | | | | | | | | | | | | | | 2.1 |
| 143.40 | | | | | | | | | | | | | | | | | | 3.8 |
| Total 2969.3 2420.1 1477.6 1726.5 2387.1 2601.4 1970.7 1032.2 626.2 512.4 330.2 145.2 59.7 34.3 61.2 15.7 7.2 3.1* 26.4 18,406.5 | | | | | | | | | | | | | | | | | | |
| Cruise 5707: | | | | | | | | | | | | | | | | | | |
| 63.70 | | | | | | | | | | | | | | | | | | 3.8 |
| 70.70 | | | | | | | | | | | | | | 4.2 | | | | 4.2 |
| 73.50 | | | | | | | | | | | | | | | | | | 4.8 |
| 77.50 | | | | 3.3 | 3.3 | | | | | | | | | | | | | 6.6 |
| .70 | | | | | | | | | | | | | | | | | | 3.8 |
| 80.51 | 7.8 | 10.4 | | | 2.6 | 7.8 | 13.0 | 15.6 | 7.8 | 10.4 | | | | | | | | 78.0 |
| .55 | | | 11.3 | 22.7 | 62.3 | 107.6 | 130.1 | 113.2 | 62.3 | 39.6 | 34.0 | | | | | | | 588.8 |
| .60 | | | | | 12.9 | 6.4 | 25.8 | 32.2 | 19.3 | 19.3 | 38.7 | 6.4 | 19.3 | 6.4 | 6.4 | | | 206.0 |
| 82.47 | | | | | 2.6 | 2.6 | 2.6 | 2.6 | | | | | | | | | | 10.4 |
| 83.40 | 36.0 | 24.4 | 17.4 | 29.0 | 53.4 | 39.5 | 25.5 | 19.7 | 5.8 | 1.2 | 1.2 | | | | | | | 253.1 |
| .43 | 38.5 | 70.6 | 59.9 | 66.4 | 179.8 | 211.9 | 111.3 | 44.9 | 19.3 | 8.6 | 4.3 | | | | | | | 815.5 |
| .51 | 2.4 | | 2.4 | | 2.4 | | | 2.4 | | | | | | | | | | 9.6 |

* - 3.1 - 24.5 mm.

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Dis. | Total | |
|-----------------------|---------------------------------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|-------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | | | 21.25 |
| Cruise 5707 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 83.55 | 8.6 | 2.8 | | | 2.8 | | 2.8 | 8.5 | 2.8 | | | | | | | | | | 28.3 |
| .70 | | | | | | | | | | | | | | | 2.9 | | | | 2.9 |
| 87.36 | 12.9 | 7.8 | 7.7 | 2.6 | | | | | | | | | | | | | | | 31.0 |
| .45 | | | | | | 5.4 | 16.2 | 21.6 | 16.2 | | | | | | | | | | 59.4 |
| .50 | 90.0 | 35.5 | 2.4 | 9.5 | 11.9 | 52.1 | 64.0 | 35.6 | 23.7 | 4.7 | 2.4 | | | | | | | | 331.8 |
| .55 | | | | 8.6 | 17.1 | 14.3 | 8.6 | 20.0 | 11.4 | 2.9 | | | | | | | | | 82.9 |
| .60 | | | | | 5.4 | | 2.7 | 2.7 | 2.7 | 2.7 | | | | | | | | 2.7 | 18.9 |
| .90 | | | | | | | | | | 28.0 | 27.9 | 28.0 | 14.0 | | | | | | 97.9 |
| 90.30 | 21.0 | 2.6 | | | 2.6 | | | | | | | | | | | | | | 26.2 |
| .37 | 29.2 | 21.9 | 31.6 | 58.3 | 41.3 | 19.5 | 9.8 | 4.9 | | 4.9 | 2.4 | 2.4 | 2.4 | | | | | | 228.6 |
| .45 | | | 4.7 | 42.3 | 42.3 | 32.9 | 98.7 | 84.6 | 42.3 | 47.0 | 23.5 | 32.9 | 9.4 | | | 4.7 | | | 465.3 |
| .50 | | | | 33.5 | | | | | 16.7 | 16.7 | 8.4 | | | | | | | | 75.3 |
| .55 | 2.4 | 4.7 | 14.1 | 23.6 | 26.0 | 23.6 | 7.1 | 7.1 | 2.4 | 7.1 | 23.6 | 7.1 | 2.4 | | | | | 7.2* | 160.8 |
| .60 | | | | | 2.3 | 2.3 | 4.6 | | | 2.3 | 2.3 | 7.0 | | | 7.0 | 11.6 | | 2.3 | 41.7 |
| .80 | | | | | | | | | | | | | | | | | | | 2.6 |
| 93.30 | 37.8 | 62.1 | 43.2 | 35.1 | 27.0 | 8.1 | | | | | | | | | | | | | 213.3 |
| .35 | | 2.6 | | | | 2.6 | | | | | | | | | | | | | 5.2 |
| 97.30 | 14.0 | 2.0 | 2.0 | 4.0 | | | 2.0 | | | | | | | | | | | | 26.0 |
| .32 | | 8.7 | 6.6 | | 2.2 | | | | | | | | | | | | | | 17.5 |
| .40 | 329.8 | 59.5 | 16.1 | 9.2 | | | | | | | | | | | | | | | 414.6 |
| 100.32 | | | | | 2.8 | | | | | | | | | | | | | | 5.6 |
| .40 | 2.6 | 2.6 | | | | 2.6 | | | | | | | | | | | | | 7.8 |
| 103.35 | | | | | | | | | | | | | | | | | | | 3.2 |
| .40 | 8.7 | 2.9 | | | | | | | | | | | | | | | | | 11.6 |
| .45 | 2.8 | | | | | | | | | | | | | | | | | | 2.8 |
| 107.32 | 3.7 | | | | | | | | | | | | | | | | | | 3.7 |
| 110.33 | 125.5 | 81.4 | 10.2 | 6.8 | 13.6 | 10.2 | 10.2 | 6.8 | 3.4 | | | | | | | | | | 268.1 |
| .35 | | | | | | | | | | | | | | | | | | | 53.4 |
| 113.30 | 2.1 | | 2.1 | | 2.1 | 4.1 | 4.1 | 4.1 | 10.3 | | | | | | | | | | 24.8 |
| .35 | 83.0 | 59.3 | 27.6 | 59.3 | 67.2 | 23.7 | 15.8 | | | | | | | | | | | | 335.9 |
| .40 | 3.2 | | | | | | | | | | | | | | | | | | 3.2 |
| 117.26 | | | | | | | | | | | | | | | | | | | 6.8 |
| .30 | | | | | | | | | | | | | | | | | | | 7.2 |
| .35 | | | | | | | | | | | | | | | | | | | 11.4 |
| .40 | | | | | | | | | | | | | | | | | | | 53.7 |
| .50 | | | | | | | | | | | | | | | | | | | 3.3 |

* - 4.8 - 23.25 mm.; 2.4 - 25.5 mm.

Table IV (cont'd)
Record of anchovy larvae, 1957

Midpoint of size class (in mm.)

| Station | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | 23.25 | Dis. | Total |
|-----------------------|--------|-------|-------|-------|--------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|---------|
| Cruise 5707 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 119.33 | | 3.4 | 17.2 | 27.5 | 10.3 | 10.3 | 10.3 | | | | | | | | | | | | | 79.0 |
| 120.30 | | 48.4 | 87.2 | 113.1 | 129.2 | 64.6 | 38.8 | 3.2 | 6.4 | | 3.2 | | | | | | | | | 494.1 |
| .35 | | 2.9 | 40.2 | 40.2 | 37.4 | 40.2 | 20.1 | 25.9 | 8.6 | 2.9 | | | | | | | | | | 218.4 |
| .40 | 3.8 | 22.7 | 24.5 | 20.7 | 32.1 | 15.2 | 15.1 | 7.6 | 1.9 | 7.6 | 1.9 | 7.6 | | 1.9 | | | | | | 162.6 |
| .55 | 3.3 | | | | | | | | | | | | | | | | | | | 3.3 |
| .70 | | | | | 3.4 | | | 3.4 | | | | | | | | | | | | 6.8 |
| .75 | | | | | 3.4 | | | | | | | | | | | | | | | 3.4 |
| 123.37 | | | | | | 2.9 | 2.9 | | | | | | | | | | | | | 5.8 |
| .42 | | | | | | | 3.1 | | | | | | | | | | | | | 3.1 |
| .50 | | | 2.9 | 17.5 | 23.2 | 29.2 | 2.9 | 2.9 | | 2.9 | | | | | | | | | | 81.5 |
| .55 | | | | 3.1 | 3.1 | | | | | | | | | | | | | | | 6.2 |
| 130.35 | | | | | 132.4 | 176.6 | 88.3 | 66.2 | | 22.1 | 44.2 | | 44.2 | | | | | | | 574.0 |
| .40 | | | 6.6 | 191.4 | 475.2 | 580.8 | 198.0 | 33.0 | 6.6 | 13.2 | | | 6.6 | | | | | | | 1,511.4 |
| .45 | 3.0 | 8.8 | 3.0 | 6.0 | 20.6 | 6.0 | | | | | | | | | | | | | | 47.4 |
| .50 | | | 3.1 | 3.1 | | 6.2 | | | | | | | | | | | | | | 12.4 |
| 133.35 | | | | | 3.4 | | | | | | | | | | | | | | | 3.4 |
| .40 | 3.4 | 41.2 | 27.5 | 13.8 | | | | | | | | | | | | | | | | 85.9 |
| .50 | 165.6 | 148.8 | 74.4 | 44.0 | 33.8 | | | | | | | | | | | | | | | 466.6 |
| 137.45 | | 3.3 | | | | | | | | | | | | | | | | | | 3.3 |
| Total | 1041.1 | 748.5 | 580.4 | 914.0 | 1524.9 | 1534.0 | 936.2 | 563.8 | 274.2 | 245.2 | 185.3 | 110.6 | 114.9 | | 31.7 | 27.5 | 8.5 | | 12.2 | 8,885.9 |

Cruise 5708:

| | | | | | | | | | | | | | | | | | | | | |
|--------|-------|------|------|------|------|------|-------|-------|------|------|-----|-----|--|-----|-----|--|--|--|--|-------|
| 110.33 | | 2.8 | | 2.8 | 2.8 | 2.8 | 8.4 | 8.4 | 2.8 | | 2.8 | | | | | | | | | 33.6 |
| .35 | 5.8 | 2.9 | | 2.9 | 2.9 | 2.9 | 2.9 | 2.9 | | | | | | | | | | | | 23.2 |
| 113.30 | 153.9 | 40.5 | 8.1 | 2.7 | 2.7 | 10.8 | 13.5 | 10.8 | 5.4 | | | | | | | | | | | 248.4 |
| .35 | 2.7 | 10.7 | 31.9 | 8.0 | 13.3 | 10.6 | 13.3 | 26.6 | 13.3 | 2.7 | 2.7 | 2.7 | | | | | | | | 138.5 |
| 115.27 | | | | 5.8 | 11.6 | 2.9 | 17.4 | 14.5 | 8.7 | | | | | | 2.9 | | | | | 63.8 |
| .30 | | 16.9 | 6.8 | 13.5 | 3.4 | | | 6.8 | 3.4 | | | | | | | | | | | 50.8 |
| 117.26 | | | | 2.9 | | 5.8 | 2.9 | 5.8 | 14.6 | 2.9 | | | | 2.9 | | | | | | 37.8 |
| .30 | | 8.8 | 14.7 | 8.8 | 32.2 | 90.9 | 175.8 | 164.1 | 90.8 | 43.9 | 5.9 | 5.8 | | | | | | | | 641.7 |
| .35 | 3.2 | | | | 16.0 | 38.4 | 22.4 | 9.6 | 3.2 | 3.2 | | | | | | | | | | 96.0 |
| .40 | | | | | 2.7 | | 2.7 | | | | | | | | | | | | | 5.4 |
| 118.25 | | | 6.0 | 3.0 | 3.0 | | | | | | | | | | | | | | | 12.0 |
| .30 | | | | | | | 3.2 | | | | | | | | | | | | | 3.2 |
| 120.25 | | | 8.9 | 5.9 | 6.0 | | 3.0 | | | | | | | | | | | | | 26.8 |
| .30 | | | 3.3 | 3.3 | 3.3 | 3.3 | | | | | | | | | | | | | | 9.9 |

Table IV (cont'd)
Record of anchovy larvae, 1957

Midpoint of size class (in mm.)

| Station | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | 23.25 | Dls. | Total |
|-----------------------|-------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|---------|
| Cruise 5708 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 120.35 | | | | 4.2 | 2.1 | 8.4 | 6.4 | 2.1 | | | | | | | | | | | | 21.2 |
| .40 | | | 2.0 | 6.1 | 4.1 | 4.1 | | 4.1 | 2.0 | | | | | | 2.0 | | | | | 24.4 |
| .45 | | | 2.9 | | | | | | | | | | | | | | | | | 2.9 |
| 127.40 | | | 3.4 | | | | | | | | | | | | | | | | | 3.4 |
| .45 | | | | | | | 2.8 | | | | | | | | | | | | | 2.8 |
| Total | 165.6 | 82.6 | 84.7 | 69.9 | 106.1 | 178.9 | 274.7 | 255.7 | 144.2 | 55.7 | 11.4 | 8.5 | | | 2.9 | 4.9 | | | | 1,445.8 |
| Cruise 5709: | | | | | | | | | | | | | | | | | | | | |
| 110.33 | | | | | | | 2.4 | | | 3.5 | | | | | | | | | | 2.4 |
| .35 | | | | | | | | | 3.1 | | | | | | | | | | | 3.5 |
| 113.35 | | | | | | | | | | | | | | | | | | | | 3.1 |
| 115.35 | | | | | | | | | | | | | | | | | | | | 3.1 |
| 117.35 | | | | | | 3.0 | | | | | | | | | | | | | 5.8 | 5.8 |
| .40 | 3.3 | | 3.3 | | | 3.3 | | | 3.3 | | | | | | | | | | | 3.0 |
| 118.30 | | | | | | 3.8 | | | | | | | | | | | | | | 13.2 |
| Total | 3.3 | | 3.3 | | 6.8 | 3.3 | 3.3 | 5.5 | 3.3 | 3.5 | | | | | | | | | 5.8 | 34.8 |
| Cruise 5710: | | | | | | | | | | | | | | | | | | | | |
| 80.51 | | | | | 2.8 | 2.8 | 2.8 | 2.8 | | | | | | | | | | | | 14.0 |
| .60 | | | | | | | | | | | | | | | | | | | 2.7 | 2.7 |
| 82.47 | | | | | 3.5 | | | | 3.5 | | 3.5 | | | | | | | | | 10.5 |
| 83.40 | | 2.6 | | 1.3 | | | | 2.6 | 1.3 | | 1.3 | | | | | | | | | 9.1 |
| .43 | | 10.6 | 7.1 | 3.6 | 3.6 | | 7.1 | 7.2 | 17.7 | 7.2 | | | | | | | | | | 64.1 |
| .70 | | | 2.4 | | | | | | | | | | | | | | | | | 2.4 |
| 87.45 | | | | | | | | 3.1 | | | | | | | | | | | | 3.1 |
| .50 | | | | | | | | | 6.4 | | | | | 3.2 | | | | | | 9.6 |
| 90.30 | | | 3.4 | 3.4 | 16.8 | 10.1 | 16.8 | | | | 3.4 | | | | | | | | | 53.9 |
| .37 | | | | | 13.4 | 40.2 | 33.5 | 26.8 | 6.8 | 3.4 | | | | | | | | | | 124.1 |
| .50 | | | | | | | | | 1.4 | | | | | | | | | | | 1.4 |
| 93.27 | | | | | | 2.8 | | | | | | | | | | | | | | 2.8 |
| .35 | | | | | | 9.0 | | 4.5 | | | | | | | | | | | | 13.5 |
| 97.30 | | 2.1 | | | 4.2 | | | 2.1 | | | | | | | | | | | | 8.4 |
| .32 | | | | | 7.6 | | | 7.5 | 3.8 | | 7.5 | | | | | | | | | 26.4 |
| .35 | | 3.5 | | | | | 7.0 | | | 3.5 | | | | | | | | | | 14.0 |
| .40 | 10.2 | 13.6 | 34.0 | 23.8 | 23.8 | 37.4 | 27.2 | 3.4 | 6.8 | 3.4 | | | | | | | | | | 183.6 |
| 100.30 | | | | | | | 3.2 | | | | | | | | | | | | | 3.2 |
| .35 | | | | | | | | | 6.6 | | | | | | | | | | | 6.6 |

Table IV (cont'd)
Record of anchovy larvae, 1957
Midpoint of size class (in mm.)

| Station | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | 17.25 | 19.25 | 21.25 | 23.25 | Dis. | Total |
|-----------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|---------|
| Cruise 5710 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 103.30 | | | | 2.6 | 18.3 | 49.8 | 55.0 | 60.3 | 26.2 | 7.8 | | | | | | | | | | 220.0 |
| .40 | 15.7 | 9.5 | | | | | | | | | | | | | | | | | | 25.2 |
| 110.33 | | | | | 2.8 | 3.0 | | | | | | | | | | | | | | 3.0 |
| 113.30 | | | | | | | 9.0 | 18.0 | 6.0 | | | | | | | | | | | 2.8 |
| 117.26 | | | | | | | 2.3 | | | | | | | | | | | | | 33.0 |
| .30 | | | 3.6 | 3.6 | 7.3 | 7.3 | | | | | | | | | | | | | | 2.3 |
| 137.23 | | | | | | | | | | | | | | | | | | | | 21.8 |
| Total | 15.8 | 42.5 | 60.0 | 38.3 | 92.3 | 180.4 | 172.9 | 126.3 | 80.5 | 25.3 | 15.7 | 3.2 | | 2.7 | 5.6 | | | | | 861.5 |
| Cruise 5711: | | | | | | | | | | | | | | | | | | | | |
| 73.51 | | | | | 6.8 | 13.6 | 6.8 | | 3.4 | | 3.4 | | | | | | | | | 34.0 |
| .55 | | | | | | | 6.3 | | | 3.2 | 3.2 | | | | | | | | | 12.7 |
| 80.55 | 15.3 | 39.8 | 116.3 | 97.9 | 79.6 | 67.3 | 67.3 | 52.0 | 61.2 | 39.8 | 12.3 | | 6.2 | 6.2 | | | | | | 593.9 |
| .60 | | 5.9 | 14.7 | 2.9 | 8.8 | 8.8 | 8.8 | 11.8 | 5.9 | 5.9 | 2.9 | | | | 2.9 | | | | | 70.5 |
| .70 | | | 2.8 | | 2.8 | | | | | | | | | | | | | | | 5.6 |
| 82.47 | 4.7 | 56.1 | 70.2 | 70.2 | 60.8 | 51.5 | 56.2 | 23.4 | 23.4 | 18.7 | 4.7 | | | | | | | | | 439.9 |
| 83.40 | 4.5 | 20.3 | 27.1 | 18.1 | 2.3 | 2.3 | 4.6 | | 4.6 | | | | | | | | | | | 83.8 |
| .43 | | 33.8 | 84.7 | 33.8 | 11.2 | 16.9 | 62.1 | 28.2 | 11.2 | 11.3 | 5.6 | | | | | | | | | 298.8 |
| .51 | 6.6 | 13.3 | 19.9 | 23.3 | 29.9 | 16.6 | | | | | | 3.3 | | | | | | | | 112.9 |
| .55 | | 4.6 | 2.3 | | | | | 2.3 | 2.3 | | | | | | | | | | | 11.5 |
| 87.36 | | 2.2 | 4.4 | 4.4 | | | | 2.2 | 2.2 | | | | | | | | | | | 15.4 |
| .50 | | | | | | | | 6.4 | | | | | | | | | | | | 6.4 |
| .55 | | | | 2.6 | | | | | | | | | | | | | | | | 2.6 |
| .60 | | | 11.5 | 19.2 | 11.5 | 11.5 | 3.8 | 3.8 | | 3.8 | | | | | | | | | 3.8 | 88.9 |
| 90.28 | | 16.6 | 49.8 | 49.7 | 33.2 | 86.3 | 33.2 | 27.6 | 33.2 | | | | 11.1 | | | 5.5 | | | | 328.2 |
| .30 | | 2.9 | 8.8 | 8.8 | 5.9 | 8.8 | | 2.9 | 2.9 | 2.9 | | | | | | | | | | 41.0 |
| .37 | | 17.4 | | 11.6 | 7.8 | 5.8 | 1.9 | | 1.9 | | | | | | | | | | | 46.4 |
| 93.27 | | 8.4 | 2.8 | 2.8 | 5.6 | | | | | | | | | | | | | | | 19.8 |
| .30 | | | | | | | | 3.0 | | | | | | | | | | | | 3.0 |
| .35 | | | 2.3 | | | | | | | | | | | | | | | | | 2.3 |
| .60 | | | | | | | 4.0 | | | | | | | | | | | | | 4.0 |
| 97.30 | | 12.6 | 21.0 | 8.4 | 8.4 | 8.4 | 8.4 | 12.6 | | 4.2 | | | | | | | | | | 75.6 |
| .32 | | 11.7 | | 3.9 | 3.9 | | | | | | | | | | | | | | | 19.5 |
| .40 | 10.2 | 35.6 | 15.3 | 10.2 | | | | | | | | | | | | | | | 3.7 | 71.3 |
| .45 | | | | | | | | | | | | | | | | | | | | 3.7 |
| .55 | | | 2.6 | | | | | | | | | | | | | | | | | 2.6 |
| Total | 26.0 | 192.9 | 343.9 | 401.9 | 304.6 | 268.4 | 309.2 | 180.8 | 144.7 | 124.9 | 32.1 | 3.3 | 6.2 | 17.3 | 2.9 | 5.5 | | | 7.5 | 2,372.1 |

Table IV (cont'd)
Record of anchovy larvae, 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Dis. | Total | | |
|--------------|---------------------------------|-------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|------|-------|-------|-------|
| | 2.50 | 3.75 | 4.75 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.75 | | | 17.25 | 19.25 |
| Cruise 5712: | | | | | | | | | | | | | | | | | | |
| 80.51 | 164.5 | 96.1 | 7.6 | 10.2 | 2.5 | 7.6 | 2.5 | 2.5 | 2.5 | 2.5 | | | | | | | | 296.0 |
| .55 | 30.0 | 12.0 | 9.0 | 3.0 | 3.0 | | | | | | | | | | | | | 57.0 |
| .60 | | | | 7.3 | | 18.3 | 3.7 | 7.4 | 7.4 | 3.7 | 11.0 | | 3.7 | | 3.7 | | | 66.2 |
| .70 | | | | 3.3 | | | | | | | | | | | | | | 3.3 |
| 90.28 | 3.3 | 6.6 | 3.3 | | | | | | | | | | | | | | | 13.2 |
| .37 | | | 3.0 | 3.0 | | | | | | | | | | | | | | 6.0 |
| 97.32 | 15.8 | 28.5 | | 6.3 | 3.2 | 6.3 | 3.2 | 3.2 | 3.2 | | | | | | | | | 66.5 |
| Total | 213.6 | 146.2 | 22.9 | 30.1 | 8.7 | 32.2 | 9.4 | 10.6 | 9.9 | 6.2 | 11.0 | | 3.7 | | 3.7 | | | 508.2 |

RECORD OF THE LARVAE OF THE JACK MACKEREL
(TRACHURUS SYMMETRICUS), 1957

Jack mackerel larvae are recorded by size in table V. The data are summarized in text table 9, by month and area. The distribution and relative abundance of jack mackerel larvae in 1957 are illustrated in figure 5. Four categories of abundance are used in the chart. The value shown for a station is the cumulative standard haul total of jack mackerel larvae taken in all occupancies during the year.

The size groupings of larvae in table V have the following midpoints and ranges:

| Midpoint (in mm.) | Range (in mm.) | Midpoint (in mm.) | Range (in mm.) |
|----------------------|-------------------|----------------------|-------------------|
| 2.00 | 1.76-2.25 | 7.75 | 7.26-8.25 |
| 2.50 | 2.26-2.75 | 8.75 | 8.26-9.25 |
| 3.00 | 2.76-3.25 | 9.75 | 9.26-10.25 |
| 3.50 | 3.26-3.75 | 10.75 | 10.26-11.25 |
| 4.00 | 3.76-4.25 | 11.75 | 11.26-12.25 |
| 4.50 | 4.26-4.75 | 12.75 | 12.26-13.25 |
| 5.00 | 4.76-5.25 | 13.75 | 13.26-14.25 |
| 5.75 | 5.26-6.25 | 14.75 | 14.26-15.25 |
| 6.75 | 6.26-7.25 | 15.26 and over | |

The relative abundance of jack mackerel larvae in different parts of the CCOFI survey area in 1957 is compared with their areal abundance during the preceding two years in the following summary:

| Lines | 1955 | | | 1956 | | | 1957 | | |
|---------|------------------|-------------|---------|------------------|-------------|---------|------------------|-------------|---------|
| | Occur- rences | Num- ber | Percent | Occur- rences | Num- ber | Percent | Occur- rences | Num- ber | Percent |
| 60-77 | 22 | 660 | 5.0 | 20 | 992 | 11.5 | 29 | 2,740 | 13.7 |
| 80-93 | 83 | 5,716 | 43.2 | 39 | 1,198 | 14.9 | 82 | 7,174 | 35.9 |
| 97-107 | 118 | 4,192 | 31.6 | 81 | 3,556 | 44.3 | 130 | 9,488 | 47.4 |
| 110-120 | 121 | 2,457 | 18.5 | 67 | 2,285 | 28.5 | 47 | 602 | 3.0 |
| 123-137 | 25 | 221 | 1.7 | 8 | 66 | 0.8 | 1 | 3 | 0.01 |
| 140-157 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 369 | 13,246 | 100.0 | 215 | 8,027 | 100.0 | 289 | 20,007 | 100.01 |




Figure 5. --Jack mackerel larvae, 1957: Distribution and relative abundance.

Text table 9. --Occurrence and abundance (standard haul totals) of jack mackerel larvae (*Trachurus symmetricus*), by month and area, in hauls made during 1957

| Cruise | Central California | | Southern California | | Northern Baja California | | Upper central Baja California | | Lower central Baja California | | Southern Baja California | | Total | |
|---------|--------------------|-------------|---------------------|-------------|--------------------------|-------------|-------------------------------|-------------|-------------------------------|-------------|--------------------------|-------------|------------------|-------------|
| | 60-77 | 80-93 | 80-93 | 80-93 | 97-107 | 97-107 | 110-120 | 110-120 | 123-137 | 123-137 | 140-157 | 140-157 | | |
| | occur- rences | num- ber | occur- rences | num- ber | occur- rences | num- ber | occur- rences | num- ber | occur- rences | num- ber | occur- rences | num- ber | occur- rences | num- ber |
| 5701 | - | - | - | - | 1 | 2 | 0 | 0 | 0 | 0 | - | - | 1 | 2 |
| 5702 | - | - | 0 | 0 | 9 | 562 | 1 | 3 | 0 | 0 | 0 | 0 | 10 | 565 |
| 5703 | - | - | 4 | 293 | 4 | 27 | 1 | 11 | 0 | 0 | 0 | 0 | 9 | 331 |
| 5704 | - | - | 5 | 378 | 22 | 735 | 10 | 187 | 1 | 3 | 0 | 0 | 38 | 1,303 |
| 5705 | 8 | 1,195 | 25 | 3,363 | 41 | 6,054 | 15 | 238 | 0 | 0 | - | - | 89 | 10,850 |
| 5706 | 16 | 1,467 | 27 | 2,778 | 38 | 1,863 | 9 | 80 | 0 | 0 | 0 | 0 | 90 | 6,188 |
| 5707 | 5 | 78 | 21 | 362 | 14 | 242 | 7 | 71 | 0 | 0 | - | - | 47 | 753 |
| 5708 | - | - | - | - | - | - | 2 | 8 | 0 | 0 | - | - | 2 | 8 |
| 5709 | - | - | - | - | - | - | 0 | 0 | 0 | 0 | - | - | 0 | 0 |
| 5710 | - | - | 0 | 0 | 1 | 3 | 2 | 4 | 0 | 0 | - | - | 3 | 7 |
| 5711 | 0 | 0 | 0 | 0 | 0 | 0 | - | - | - | - | - | - | 0 | 0 |
| 5712 | - | - | 0 | 0 | 0 | 0 | 0 | 0 | - | - | - | - | 0 | 0 |
| Total | 29 | 2,740 | 82 | 7,174 | 130 | 9,488 | 47 | 602 | 1 | 3 | 0 | 0 | 289 | 20,007 |
| Percent | 13.70 | | 35.86 | | 47.42 | | 3.01 | | 0.01 | | 0 | | 100.00 | |

Data for the several years are only roughly comparable since the coverage was somewhat different in each year. Only a portion of the spawning range of the jack mackerel is surveyed on CCOFI cruises, and it is likely that a variable portion of the population is within the survey area in successive seasons. The most striking change in distribution of jack mackerel larvae in 1957 was the marked decrease in abundance off central and southern Baja California. Nearly 30 percent of jack mackerel larvae were taken in the area off central Baja California (station lines 110-137) in 1956, and over 20 percent in 1955, while only 3 percent occurred in this area in 1957. The northward shift in distribution of jack mackerel larvae is probably related to the warmer water temperatures prevailing in 1957.

The seasonal distribution of larger jack mackerel larvae (6.26 mm. and larger) during 1957 is compared with the distributions of the preceding five years in the following tabulation:

Seasonal abundance of larger jack mackerel larvae (6.26 mm. and over), 1952-1957 (standard haul totals)

| Months | 1952 | 1953 | 1954 | 1955 | 1956 | 1957 |
|----------|------|------|------|------|------|------|
| January | 0 | 0 | 0 | 0 | 0 | 0 |
| February | 0 | 3 | 0 | 3 | 0 | 0 |
| March | 11 | 56 | 8 | 9 | 105 | 13 |
| April | 47 | 38 | 42 | 223 | 54 | 13 |
| May | 56 | 77 | 46 | 80 | 58 | 62 |
| June | 286 | 310 | 246 | 80 | 108 | 635 |
| July | 189 | 89 | 57 | 335 | 67 | 56 |
| August | 31 | 3 | 18 | - | 0 | 0 |
| October | 0 | 44 | 0 | 0 | 0 | 0 |
| Total | 620 | 620 | 417 | 730 | 392 | 779 |

There was a marked peak in the abundance of larger larvae in June, 1957. A similar pattern of seasonal abundance was found in 1952 through 1954. The seasonal distributions in 1955 and 1956 were bimodal, however; peaks in abundance of larger larvae occurred in April and July during 1955 and in March and June of 1956. The numbers of larger larvae taken in June 1957 were markedly greater than during any other one month, but the season total was not much larger than that for 1955.

Table V
Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | Total | | | | | | | |
|---------------------|---------------------------------|------|-------|-------|------|------|------|------|------|------|------|-------|------|-------|-------|-------|-------|-------|-------------------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | 14.75 | 15.26 and over |
| Cruise 5701: | | | | | | | | | | | | | | | | | | | |
| 103.30 | 2.4 | | | | | | | | | | | | | | | | | | |
| Total | 2.4 | | | | | | | | | | | | | | | | | | |
| Cruise 5702: | | | | | | | | | | | | | | | | | | | |
| 100.60 | 6.8 | 13.6 | 30.7 | 20.5 | | | | | | | | | | | | | | | 71.6 |
| .70 | | | 43.4 | 14.5 | | | | | | | | | | | | | | | 57.9 |
| .80 | 6.9 | 13.7 | 17.2 | 20.6 | 3.4 | | | | | | | | | | | | | | 61.8 |
| .90 | 7.4 | 3.7 | | | | | | | | | | | | | | | | | 11.1 |
| 103.50 | 7.2 | | | | | | | | | | | | | | | | | | |
| .70 | | | 12.9 | 25.8 | 19.4 | | | | | | | | | | | | | | 58.1 |
| 107.50 | 16.3 | 27.2 | 81.6 | 97.9 | 16.3 | | | | | | | | | | | | | | 239.3 |
| .60 | | | 5.6 | 22.6 | 16.9 | 5.6 | | | | | | | | | | | | | 50.7 |
| .70 | | | 2.3 | 2.3 | | | | | | | | | | | | | | | 4.6 |
| 110.50 | 2.7 | | | | | | | | | | | | | | | | | | |
| Total | 37.4 | 58.2 | 193.7 | 214.1 | 56.0 | 5.6 | | | | | | | | | | | | | 565.0 |
| Cruise 5703: | | | | | | | | | | | | | | | | | | | |
| 83.60 | 5.9 | | | | | | | | | | | | | | | | | | |
| .70 | | | 11.5 | 11.5 | 11.5 | | | | | | | | | | | | | | 5.9 |
| 90.90 | 9.5 | 53.7 | 75.8 | 53.7 | 37.9 | 9.5 | 3.2 | 3.2 | | | | | | | | | | | 34.5 |
| 93.70 | 6.0 | | | | | | | | | | | | | | | | | | |
| 103.70 | 6.2 | | | | | | | | | | | | | | | | | | |
| 107.35 | 3.2 | | | | | | | | | | | | | | | | | | 6.2 |
| .60 | | | 3.8 | 3.8 | 3.8 | 3.8 | 3.8 | | | | | | | | | | | | 3.2 |
| .70 | 3.1 | | | | 3.1 | | | | | | | | | | | | | | 11.4 |
| 110.50 | 7.5 | | | | | | | | | | | | | | | | | | |
| Total | 6.3 | 9.5 | 67.4 | 91.1 | 68.3 | 49.4 | 13.3 | 12.9 | 3.2 | 6.0 | 3.7 | | | | | | | | 331.1 |
| Cruise 5704: | | | | | | | | | | | | | | | | | | | |
| 83.90 | 2.4 | | | | | | | | | | | | | | | | | | |
| 90.45 | 39.5 | | | | | | | | | | | | | | | | | | 39.5 |
| .80 | | | 49.6 | 86.8 | 24.8 | | | | | | | | | | | | | | 161.2 |
| 93.50 | | | 14.9 | | | | | | | | | | | | | | | | 14.9 |
| .60 | 22.8 | 22.8 | 22.8 | 57.0 | 34.2 | 22.8 | | | | | | | | | | | | | 159.6 |

Table V (cont'd)
 Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Total | | | | | | |
|-----------------------|---------------------------------|-------|-------|-------|-------|-------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|----------------|-------|---------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | | 12.75 | 13.75 | 14.75 | 15.26 and over | | |
| Cruise 5704 (cont'd): | | | | | | | | | | | | | | | | | | | | | |
| 97.45 | | 12.8 | 76.8 | 179.2 | 128.0 | 25.6 | 12.8 | | | | | | | | | | | | 6.1 | 435.2 | |
| .60 | | | | | | | | | | | | | | | | | | 4.9 | | | |
| 100.40 | | 3.2 | | | | | | | | | | | | | | | | | | 3.2 | 38.5 |
| .45 | | | 4.8 | 9.6 | 14.5 | 4.8 | | 4.8 | | | | | | | | | | | | | 80.9 |
| .60 | | 7.0 | 31.7 | 21.1 | 12.3 | 5.3 | 3.5 | | | | | | | | | | | | | | 17.0 |
| .70 | | 3.4 | 3.4 | 6.8 | | | | | | | | | | | | | | | | | 11.5 |
| .80 | 3.4 | 7.2 | 4.3 | | | | | | | | | | | | | | | | | | 10.5 |
| 103.30 | | | | | 5.3 | 2.6 | 2.6 | | | | | | | | | | | | | | 11.2 |
| .35 | | | | | | | | | | | | | | | | | | | | | 36.9 |
| .40 | | | | | 5.6 | | 2.8 | | 2.8 | | | | | | | | | | | | 2.7 |
| .45 | | 3.1 | | 9.2 | 12.3 | 9.2 | 3.1 | | | | | | | | | | | | | | 11.2 |
| .50 | | 5.6 | | | 5.6 | | 2.7 | | | | | | | | | | | | | | 2.7 |
| .55 | | | | | | | | | | | | | | | | | | | | | 11.2 |
| .60 | | 3.0 | 3.0 | 10.2 | | | | | | | | | | | | | | | | | 13.6 |
| .80 | | 3.0 | 3.0 | 3.0 | | | | | | | | | | | | | | | | | 9.0 |
| .90 | 3.1 | | | | | | | | | | | | | | | | | | | | 3.1 |
| 107.35 | | 3.2 | | | | | | | | | | | | | | | | | | | 3.2 |
| .40 | | | | | 6.7 | 3.3 | | | | | | | | | | | | | | | 13.3 |
| .45 | | | | | 2.8 | | | | | | | | | | | | | | | | 8.4 |
| .55 | | | | | 6.4 | | | | | | | | | | | | | | | | 6.4 |
| .60 | | | | | | | | | | | | | | | | | | | | | 3.5 |
| .70 | | | | | | | | | | | | | | | | | | | | | 4.4 |
| 110.40 | | | | | | | | | | | | | | | | | | | | | 7.1 |
| .60 | | | | | 7.7 | 3.9 | | | | | | | | | | | | | | | 11.6 |
| .70 | | | | | 13.2 | 26.4 | 19.8 | 16.5 | 6.6 | | | | | | | | | | | | 82.5 |
| .90 | | | | | 3.4 | 6.7 | | | | | | | | | | | | | | | 10.1 |
| 113.35 | | 15.8 | 4.0 | 4.0 | | | | | | | | | | | | | | | | | 23.8 |
| .45 | | 5.7 | | | | | | | | | | | | | | | | | | | 8.5 |
| .80 | | | | | 2.9 | 2.9 | | | | | | | | | | | | | | | 5.8 |
| 117.40 | | | | | | | | | | | | | | | | | | | | | 10.4 |
| .70 | | 2.7 | 8.0 | | | 10.4 | | | | | | | | | | | | | | | 10.7 |
| .80 | | 7.3 | 2.4 | 2.4 | 4.8 | | | | | | | | | | | | | | | | 16.9 |
| 123.55 | | | | | | 2.6 | | | | | | | | | | | | | | | 2.6 |
| Total | 32.3 | 133.3 | 228.9 | 454.5 | 271.0 | 106.4 | 39.8 | 22.8 | 5.6 | 4.9 | 2.8 | | | | | | | | | | 1,302.3 |

Table V (cont'd)
Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | Total | | |
|--------------|---------------------------------|-------|-------|-------|------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|---------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | | 13.75 | 14.75 |
| Cruise 5705: | | | | | | | | | | | | | | | | | | |
| 70.52 | 14.1 | 14.1 | 28.2 | 21.1 | | | | | | | | | | | | | | 77.5 |
| .60 | | 25.8 | 68.8 | 34.4 | | | | | | | | | | | | | | 129.0 |
| .70 | 8.9 | 53.4 | 93.4 | 71.2 | | | | | | | | | | | | | | 226.9 |
| .80 | 3.5 | | | | | | | | | | | | | | | | | 3.5 |
| .90 | | 3.3 | | | | | | | | | | | | | | | | 3.3 |
| 73.60 | 36.5 | 77.5 | 145.9 | 95.8 | | | | | | | | | | | | | | 355.7 |
| .70 | 18.5 | 33.0 | 49.8 | 33.0 | | | | | | | | | | | | | | 132.1 |
| 77.70 | 29.7 | 59.4 | 89.0 | 89.0 | | | | | | | | | | | | | | 267.1 |
| 80.70 | | 14.3 | 14.3 | 71.4 | | | | | | | | | | | | | | 100.0 |
| .90 | | | | 3.6 | 11.0 | 7.3 | | | | | | | | | | | | 21.9 |
| 83.60 | | | | | | 2.9 | | | | | | | | | | | | 2.9 |
| .80 | 2.8 | | | | | | | | | | | | | | | | | 2.8 |
| .90 | | 3.3 | | | | | | | | | | | | | | | | 42.7 |
| 87.55 | 26.0 | 6.5 | | | | | | | | | | | | | | | | 32.5 |
| .75 | 10.0 | 20.1 | | | | | | | | | | | | | | | | 30.1 |
| .80 | 6.2 | 27.8 | 3.1 | | | | | | | | | | | | | | | 37.1 |
| .85 | 8.3 | 8.3 | | | | | | | | | | | | | | | | 16.6 |
| .90 | | | 3.4 | | | | | | | | | | | | | | | 6.8 |
| 90.55 | 208.1 | 115.6 | | | | | | | | | | | | | | | | 323.7 |
| .60 | 246.2 | 109.4 | 109.4 | | | | | | | | | | | | | | | 465.0 |
| .70 | | | 3.3 | | | | | | | | | | | | | | | 3.3 |
| .75 | 9.8 | 6.5 | | | | | | | | | | | | | | | | 16.3 |
| .80 | | | | | | | | | | | | | | | | | | 5.2 |
| .90 | | | 12.1 | 9.1 | 3.0 | | | | | | | | | | | | | 24.2 |
| 93.40 | 32.3 | 35.0 | | | 2.7 | | | | | | | | | | | | | 70.0 |
| .45 | 68.4 | 42.1 | 15.8 | | | | | | | | | | | | | | | 126.3 |
| .50 | 18.2 | 36.4 | 54.6 | | | | | | | | | | | | | | | 109.2 |
| .55 | 440.8 | 957.0 | 75.4 | | | | | | | | | | | | | | | 1,473.2 |
| .60 | 27.8 | 197.8 | 105.1 | 37.1 | 3.1 | | | | | | | | | | | | | 370.9 |
| .65 | | | 5.1 | 25.6 | 5.1 | | | | | | | | | | | | | 35.8 |
| .70 | 5.4 | 10.8 | 2.7 | 2.7 | | 2.7 | | | | | | | | | | | | 24.3 |
| .75 | | | 10.7 | 5.4 | | | | | | | | | | | | | | 18.8 |
| .85 | 3.4 | | | | | | | | | | | | | | | | | 3.4 |
| 97.30 | | | 8.9 | 4.5 | | | | | | | | | | | | | | 13.4 |
| .55 | 155.5 | 699.8 | 959.0 | 233.3 | | | | | | | | | | | | | | 2,060.6 |

Table V (cont'd)
Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | Total | | | | | | | |
|-----------------------|---------------------------------|------|-------|-------|------|------|------|------|------|------|------|-------|------|-------|-------|-------|-------|-------|-------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | | 9.75 | 10.75 | 11.75 | 12.75 | 13.76 | 14.75 | 15.26 |
| Cruise 5705 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 97.60 | 27.4 | 78.4 | 39.2 | 7.8 | 3.9 | 7.8 | 31.4 | 3.9 | | | | | | | | | | | 199.8 |
| .65 | 8.7 | 26.1 | 110.2 | 29.0 | 2.9 | 8.7 | 5.8 | | | | | | | | | | | | 191.4 |
| .70 | | 21.3 | 71.1 | 37.9 | 9.5 | 7.1 | 4.7 | 7.1 | | | | | | | | | | | 158.7 |
| .75 | 16.8 | 23.4 | 10.0 | 13.4 | 13.4 | 6.7 | 6.7 | | | | | | | | | | | | 97.1 |
| .80 | | | | | 3.5 | | | | | | | | | | | | | | 3.5 |
| .90 | | | | | | | 1.9 | | | | | | | | | | | | 1.9 |
| 100.29 | | | | | | | | | | | | | | | | | | | |
| .33 | | 8.5 | 8.5 | | | | | | | | | | | | | | | | 4.5 |
| .45 | 2.8 | 9.6 | 16.8 | 2.4 | | | | | | | | | | | | | | | 17.0 |
| .50 | | | | | | | | | | | | | | | | | | | 4.5 |
| .60 | | | | | | | | | | | | | | | | | | | 17.0 |
| .65 | 6.0 | 15.0 | 72.2 | 63.2 | 15.0 | 9.0 | 9.0 | | | | | | | | | | | | 5.8 |
| .70 | 13.7 | 82.1 | 99.2 | 27.4 | 13.7 | 20.5 | 13.7 | 6.8 | | | | | | | | | | | 28.8 |
| .75 | 6.8 | 27.4 | 68.4 | 20.5 | 11.4 | 18.2 | 16.0 | 4.6 | | | | | | | | | | | 20.9 |
| .80 | 8.7 | 34.7 | 60.8 | 36.9 | 32.6 | 62.9 | 39.1 | 10.8 | | | | | | | | | | | 198.4 |
| .85 | | 3.4 | 13.7 | 17.1 | 13.7 | 13.7 | 6.8 | 3.4 | | | | | | | | | | | 280.5 |
| .90 | | 6.4 | 32.0 | 38.4 | 9.6 | 3.2 | 3.2 | 3.2 | | | | | | | | | | | 173.3 |
| 103.35 | | | | | | | | | | | | | | | | | | | |
| .45 | 2.6 | 10.2 | 17.9 | 12.8 | 5.1 | | | | | | | | | | | | | | 286.5 |
| .50 | 5.2 | 5.2 | 7.8 | | | | | | | | | | | | | | | | 71.8 |
| .55 | | 3.6 | 14.3 | 21.4 | 10.7 | 3.6 | | | | | | | | | | | | | 96.0 |
| .60 | | 6.9 | 10.4 | 10.4 | 13.9 | 10.4 | 3.5 | | | | | | | | | | | | 13.2 |
| .65 | | 2.6 | 50.4 | 177.6 | 66.2 | 26.5 | 8.0 | 7.9 | 5.2 | 7.9 | 2.6 | | | | | | | | 48.6 |
| .70 | 6.4 | 12.8 | 28.9 | 44.9 | 16.0 | 3.2 | 3.2 | | | | | | | | | | | | 18.2 |
| .75 | | 5.9 | 8.8 | 11.7 | 8.8 | | | | | | | | | | | | | | 53.6 |
| .80 | | 20.6 | 27.4 | 13.7 | | | | | | | | | | | | | | | 55.5 |
| .85 | | 19.4 | 43.6 | 14.5 | | | | | | | | | | | | | | | 354.9 |
| .90 | 5.4 | 10.8 | 81.0 | 89.1 | 29.7 | 13.6 | 5.4 | 2.7 | | | | | | | | | | | 118.8 |
| 107.35 | | | | | | | | | | | | | | | | | | | |
| .40 | | 6.8 | 13.5 | 11.2 | 4.5 | | | | | | | | | | | | | | 41.1 |
| .50 | | | 29.9 | 24.9 | 19.9 | 6.0 | | | | | | | | | | | | | 65.1 |
| .55 | 8.6 | 49.2 | 119.8 | 44.9 | 21.4 | 17.1 | 4.3 | 2.1 | 2.6 | 2.1 | 2.1 | | | | | | | | 77.5 |
| .60 | | | 161.1 | 127.2 | 33.9 | 17.0 | | | | | | | | | | | | | 337.6 |
| .65 | | 3.0 | 36.2 | 42.3 | 21.1 | | | | | | | | | | | | | | 16.8 |
| .70 | | 9.4 | 61.4 | 75.5 | 14.2 | | | | | | | | | | | | | | 36.0 |
| .75 | | | 4.2 | 4.2 | | | | | | | | | | | | | | | 82.2 |
| .80 | | | 3.2 | | | | | | | | | | | | | | | | 269.5 |
| .85 | | 3.6 | | | | | | | | | | | | | | | | | 347.7 |
| .90 | | 2.9 | 5.8 | 2.9 | | | | | | | | | | | | | | | 105.6 |
| | | | | | | | | | | | | | | | | | | | 160.5 |
| | | | | | | | | | | | | | | | | | | | 10.5 |
| | | | | | | | | | | | | | | | | | | | 9.6 |
| | | | | | | | | | | | | | | | | | | | 7.2 |
| | | | | | | | | | | | | | | | | | | | 14.5 |

Table V (cont'd)
Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Total | | | | |
|-----------------------|---------------------------------|--------|--------|--------|-------|-------|-------|-------|------|------|------|------|-------|-------|-------|-------|-------|-------|----------------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | | 12.75 | 13.75 | 14.75 | 15.26 and over |
| Cruise 5705 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 110.55 | 6.3 | 6.3 | | | | | | | | | | | | | | | | | 12.6 |
| .75 | 2.9 | | | | | | | | | | | | | | | | | | 2.9 |
| .80 | 3.0 | 3.0 | 6.0 | 9.1 | | | | | | | | | | | | | | | 21.1 |
| .85 | 2.8 | 2.8 | 2.8 | | | | | | | | | | | | | | | | 8.4 |
| .90 | | | | | 6.1 | | | | | | | | | | | | | | 6.1 |
| 113.35 | | 43.4 | 6.2 | 6.2 | 12.4 | | | | | | | | | | | | | | 68.2 |
| .40 | 6.1 | | | | | | | | | | | | | | | | | | 6.1 |
| .50 | | | | | | 2.6 | | | | | | | | | | | | | 2.6 |
| .70 | | 18.1 | 25.9 | 7.8 | 2.6 | | | | | | | | | | | | | | 54.4 |
| .80 | | | 3.0 | | | | | | | | | | | | | | | | 3.0 |
| 117.45 | 4.6 | | | | | | | | | | | | | | | | | | 4.6 |
| .75 | 2.7 | 5.4 | 5.4 | | | | | | | | | | | | | | | | 16.2 |
| .80 | | 16.1 | 5.4 | | | | | | | | | | | | | | | | 21.5 |
| 119.33 | 5.4 | | | | | | | | | | | | | | | | | | 5.4 |
| 120.75 | | | | | | | | | | | | | | | | | | | 5.0 |
| Total | 1513.5 | 3148.0 | 3277.8 | 1818.2 | 470.9 | 272.5 | 143.4 | 143.7 | 29.4 | 21.1 | 11.4 | | | | | | | | 10,849.9 |
| Cruise 5706: | | | | | | | | | | | | | | | | | | | |
| 63.70 | 6.7 | | | | | | | | | | | | | | | | | | 6.7 |
| .80 | | | 3.6 | 3.6 | 3.6 | 3.6 | | | | | | | | | | | | | 14.4 |
| .90 | | | | | | | | | | | | | | | | | | | 3.6 |
| 67.80 | 6.4 | 3.2 | | | | | | | | | | | | | | | | | 9.6 |
| .90 | 3.2 | 6.3 | 3.2 | | | | | | | | | | | | | | | | 12.7 |
| 70.60 | 6.3 | | | | | | | | | | | | | | | | | | 6.3 |
| .90 | | 16.5 | 22.0 | 27.5 | 22.0 | 11.0 | 5.5 | 11.0 | 5.5 | | | | | | | | | | 121.0 |
| 73.60 | 12.1 | | | | | | | | | | | | | | | | | | 24.2 |
| .70 | | 12.4 | 21.7 | 49.6 | 71.3 | 43.4 | 15.5 | 40.3 | 31.0 | 6.2 | 3.1 | | | | | | | | 294.5 |
| .80 | | 21.6 | 33.9 | 138.6 | 224.8 | 194.0 | 61.6 | 27.8 | 9.3 | 6.2 | 3.1 | | | | | | | | 720.9 |
| .90 | | | | 6.9 | 24.1 | | | | | | | | | | | | | | 31.0 |
| 77.55 | | | 3.0 | 6.0 | | | | | | | | | | | | | | | 24.0 |
| .60 | | 6.4 | | | 6.4 | | | | | | | | | | | | | | 70.4 |
| .70 | | | | | | 11.5 | | | | | | | | | | | | | 69.0 |
| .80 | | | | | | | | | | | | | | | | | | | 2.8 |
| .90 | 3.1 | 18.7 | 6.2 | 9.3 | 3.1 | 3.1 | 6.2 | 3.1 | | | | | | | | | | | 55.9 |

Table V (cont'd)
Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | Total | | | |
|-----------------------|---------------------------------|-------|------|------|------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|----------------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | | 13.75 | 14.75 | 15.26 and over |
| Cruise 5706 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 80.60 | 16.6 | 27.6 | | | | 16.6 | 11.0 | 5.5 | | | | | | | | | | | 77.3 |
| .70 | | | 26.3 | 39.5 | 65.8 | 39.5 | 79.0 | 26.4 | 13.2 | | | | | | | | | | 289.7 |
| .80 | 19.1 | 38.2 | 6.4 | 12.7 | 50.9 | 63.6 | 57.2 | 76.3 | 25.4 | 12.7 | | | | | | | | | 362.5 |
| .90 | 33.2 | 78.5 | 27.2 | 33.2 | 30.2 | 24.2 | 9.1 | 18.2 | 6.0 | | | | | | | | | | 259.8 |
| 83.60 | 3.4 | 6.9 | 27.4 | | 3.4 | 10.3 | 6.9 | 20.6 | 34.3 | 27.5 | 3.4 | | | | | | | | 144.1 |
| .65 | | 6.3 | | | | 3.2 | 6.3 | 3.2 | | | | | | | | | | | 19.0 |
| .70 | | | 9.0 | 24.0 | 36.0 | 33.0 | 9.0 | 21.0 | 3.0 | | | | | | | | | | 135.0 |
| .75 | | | | 10.4 | 20.8 | | 10.4 | | 20.8 | | | | | | | | | | 62.4 |
| .80 | | | | 32.3 | 19.4 | 64.6 | 32.3 | 13.0 | | | | | | | | | | | 161.6 |
| .85 | 5.9 | 17.6 | 5.9 | 29.3 | 20.5 | 17.6 | 8.8 | | | | | | | | | | | | 120.2 |
| .90 | | 3.0 | 11.8 | 8.8 | 3.0 | | | 3.0 | | | | | | | | | | | 29.6 |
| 87.65 | | | | | | | | 11.0 | | | | | | | | | | | 11.0 |
| .85 | | 2.7 | 2.7 | | | | | | | | | | | | | | | | 5.4 |
| .90 | | | | 18.1 | 12.0 | 9.0 | 9.0 | 6.0 | 6.0 | | | | | | | | | | 60.1 |
| 90.70 | | | 3.1 | 3.1 | | | | | 3.1 | | | 3.1 | | | | | | | 12.4 |
| .75 | | | | | | | | | 15.0 | | | | | | | | | | 15.0 |
| .90 | | | | 3.4 | | | | 3.4 | 10.1 | | | 3.4 | 3.4 | | | | | | 23.7 |
| 93.30 | 2.9 | | | | | | | | | | | | | | | | | | 2.9 |
| .40 | | | | | | | | | | | | 3.1 | | | | | | | 3.1 |
| .55 | 32.8 | 45.9 | 39.4 | 3.3 | | 3.3 | 23.0 | 9.9 | 3.3 | 3.3 | | | | | | | | | 164.2 |
| .60 | | 14.8 | 37.0 | 37.0 | 7.4 | | | 14.8 | | | | | | | | | | | 111.0 |
| .65 | 157.8 | 161.5 | 33.0 | 18.4 | 40.4 | 36.7 | 25.7 | 14.7 | 11.0 | 7.3 | | | | | | | | | 506.5 |
| .70 | | 4.0 | 4.0 | | | | | | | | | | | | | | | | 8.0 |
| .75 | 13.9 | 20.9 | 7.0 | | | | | | | | | | | | | | | | 41.8 |
| .80 | 35.2 | 35.2 | 42.2 | 28.2 | | | | | | | | | | | | | | | 140.8 |
| .85 | | | | 7.1 | | | | | | | | | | | | | | | 7.1 |
| .90 | | | | | 3.8 | | | | | | | | | | | | | | 3.8 |
| 97.50 | | | | | | | | | | | | | | | | | | | 3.8 |
| .55 | 6.6 | 6.6 | 6.6 | | | | 6.6 | 19.7 | | | | | | | | | | | 3.8 |
| .60 | 112.8 | 128.1 | 64.0 | 9.2 | 15.2 | 33.6 | 36.6 | 42.7 | 15.2 | 3.0 | 3.0 | | | | | | | | 46.1 |
| .65 | 14.2 | 49.6 | 35.4 | 17.7 | 7.1 | 7.1 | 21.2 | 3.5 | | 3.5 | | | | | | | | | 463.4 |
| .70 | | | 9.7 | 22.5 | 3.2 | 3.2 | | | | | | | | | | | | | 159.3 |
| .75 | 9.3 | 6.2 | 28.0 | 96.4 | 21.8 | 12.4 | 6.2 | | | | | | | | | | | | 38.6 |
| .80 | | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | | | | | | | | | | | | | 180.3 |
| .85 | 3.0 | | 3.0 | | | | | | | | | | | | | | | | 11.2 |
| .90 | 6.1 | 6.1 | 3.0 | 6.1 | 3.0 | 6.1 | | | | | | | | | | | | | 8.0 |
| | | | | | | | | | | | | | | | | | | | 21.3 |

Table V (cont'd)
Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Total | | | |
|-----------------------|---------------------------------|------|------|------|------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------------------|-------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | | 14.75 | 15.26 and over | Dis. |
| Cruise 5706 (cont'd): | | | | | | | | | | | | | | | | | | | | |
| 100.40 | | | 3.9 | | | | | | | | | | | | | | | | | 3.9 |
| .45 | | 3.0 | | | | | | | | | | | | | | | | | | 3.0 |
| .50 | 3.0 | 3.0 | 9.0 | | | | | | | | | | | | | | | | | 18.0 |
| .55 | | 6.6 | 9.9 | 23.2 | 6.6 | | | 19.8 | 23.1 | 9.9 | | | | | | | | | | 99.1 |
| .60 | 3.1 | 3.1 | | 6.2 | 3.1 | 9.3 | 9.3 | 15.5 | 12.4 | | | | | | | | | | | 62.0 |
| .65 | | 8.1 | 27.0 | 16.2 | 16.2 | 5.4 | | | | | | | | | | | | | | 72.9 |
| .70 | | | | 2.8 | 2.8 | | 5.6 | | | | | | | | | | | | | 11.2 |
| .75 | | 3.0 | 15.2 | 15.2 | 3.0 | | | | | | | | | | | | | | | 36.4 |
| .80 | | | | | | | | | | | | | 3.5 | | | | | | | 3.5 |
| .85 | | | | | | 3.1 | | | | | | | | | | | | | | 3.1 |
| .90 | | | | 11.2 | | 5.6 | 2.8 | | | 2.8 | | | | | | | | | | 25.2 |
| 103.30 | 2.8 | | 2.8 | | | | | | | | | | | | | | | | | 2.8 |
| .35 | 5.9 | 2.9 | 2.9 | 5.9 | 5.9 | 8.8 | 2.9 | 2.9 | 2.9 | 2.9 | | | | | | | | | | 41.0 |
| .40 | | | 3.8 | | | 7.6 | 15.3 | 7.6 | | | | | | | | | | | | 34.3 |
| .45 | | | | 3.9 | 7.7 | 15.5 | 15.5 | 3.9 | 3.9 | 3.9 | | | | | | | | | | 65.9 |
| .50 | | | 3.4 | 6.7 | | | 6.8 | | | | | | 3.4 | | | | | | 3.4 | 23.7 |
| .55 | | | 3.3 | 9.8 | 3.3 | | 3.3 | | | | | | | | | | | | | 23.0 |
| .60 | | | | 10.1 | | | 10.1 | 13.4 | 3.4 | 3.4 | 3.4 | 3.4 | | | | | | | | 43.8 |
| .65 | | 3.7 | 25.8 | 48.0 | 66.4 | 59.0 | 18.4 | 7.4 | 3.7 | | | | 3.7 | | | | | | | 239.8 |
| .70 | | | 3.3 | 16.4 | 13.2 | | | | | | | | | | | | | | | 32.9 |
| .75 | 3.4 | | 6.8 | 3.4 | | | | | | | | | | | | | | | | 13.6 |
| 107.40 | | | 3.4 | | | | 3.4 | 3.4 | | | | | | | | | | | | 13.6 |
| .50 | | | | | | | | | | | | | | | | | | | | 8.7 |
| .55 | | | | | | | | | | | | | | | | | | | | 6.2 |
| .60 | | | | | | | | | | | | | | | | | | | | 3.4 |
| .65 | | | | 3.3 | | | | 3.4 | | | | | | | | | | | | 6.6 |
| .70 | 3.2 | | 3.2 | | 3.2 | | 3.2 | 3.2 | 3.2 | | | | | | | | | | | 16.0 |
| .75 | 3.1 | | | 3.1 | | | 3.1 | 6.2 | | | | | | | | | | | | 15.5 |
| .80 | | | 3.6 | | | | | | | | | | | | | | | | | 3.6 |
| 110.45 | | 3.0 | 6.1 | | | | | | | | | | | | | | | | | 9.1 |
| .65 | | | | | 2.5 | | | | | | | | | | | | | | | 2.5 |
| .75 | | | | | 3.2 | | | | | | | | | | | | | | | 3.2 |
| .80 | | | 5.9 | | | | | | | | | | | | | | | | | 5.9 |

Table V (cont'd)
Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | | | Dis. and over | Total |
|----------------------|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|------|-------|-------|-------|-------|------------------|-------------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | 12.75 | 13.75 | | |
| Cruise 5706 (cont'd) | | | | | | | | | | | | | | | | | | |
| 113.75 | | | | 2.5 | 2.5 | 5.1 | 2.5 | | | | | | | | | | | 12.6 |
| .80 | | 15.1 | 10.1 | 10.1 | 10.1 | | | | | | | | | | | | | 35.3 |
| 117.35 | | | 2.4 | | | | | | | | | | | | | | | 2.4 |
| 120.55 | | 2.3 | | | | | | | | | | | | | | | | 2.3 |
| .75 | | | 3.5 | 3.5 | | | 3.5 | | | | | | | | | | | 7.0 |
| Total | 518.5 | 775.3 | 681.4 | 907.8 | 826.4 | 792.6 | 454.4 | 590.3 | 370.4 | 169.9 | 36.0 | 25.8 | 17.0 | 3.4 | 2.9 | 6.3 | 3.7 | 5.9 6,188.0 |
| Cruise 5707: | | | | | | | | | | | | | | | | | | |
| 63.55 | | | | | | | | | | | | | | | | | | 20.2 |
| .60 | | 14.3 | | | | | | | | | | | | | | | | 14.3 |
| 67.55 | 6.4 | 6.4 | 12.7 | 6.4 | | | | | | | | | | | | | | 31.9 |
| .70 | | | | | | 8.0 | | | | | | | | | | | | 8.0 |
| 77.80 | | 3.6 | | | | | | | | | | | | | | | | 3.6 |
| 80.60 | | | 6.4 | | 6.4 | | | | | | | | | | | | | 3.6 |
| .70 | | | | | 14.9 | | | | | | | | | | | | | 12.8 |
| .90 | | | | | 2.7 | | | | | | | | | | | | | 14.9 |
| 83.40 | | 1.2 | | | | | | | | | | | | | | | | 2.7 |
| .51 | 2.4 | 2.4 | | 2.4 | | | 2.4 | | | | | | | | | | | 1.2 |
| .70 | | | | | | | | | | | | | | | | | | 9.6 |
| .85 | | | | | | | | | | | | | | | | | | 5.8 |
| 87.75 | | | | | | | | | | | | | | | | | | 2.8 |
| 90.75 | | | | | | | | | | | | | | | | | | 3.8 |
| .80 | 2.6 | | | | | | | | | | | | | | | | | 4.8 |
| .90 | 10.6 | 10.6 | 5.3 | 15.8 | | 10.6 | | | | | | | | | | | | 5.2 |
| 93.40 | | | | | | | | | | | | | | | | | | 63.5 |
| .45 | | | | | | | | | | | | | | | | | | 2.9 |
| .50 | 2.8 | | 4.4 | 4.4 | 2.2 | | | | | | | | | | | | | 11.0 |
| .55 | | | | | 2.8 | | | | | | | | | | | | | 8.4 |
| .60 | 2.3 | 12.0 | 15.0 | 18.0 | 15.0 | 6.0 | | | | | | | | | | | | 66.0 |
| .65 | 2.2 | 8.9 | 2.2 | 2.2 | 4.7 | | | | | | | | | | | | | 20.9 |
| .70 | | 3.8 | 19.2 | 26.9 | | | | | | | | | | | | | | 17.7 |
| .75 | 3.6 | 10.7 | 3.6 | | 3.6 | | 3.8 | 3.8 | 3.8 | 3.8 | | | | | | | | 65.1 |
| .80 | 5.2 | 5.2 | 2.6 | | 2.6 | | | | | | | | | | | | | 21.5 |
| .85 | 2.8 | 2.8 | | | | | | | | | | | | | | | | 15.6 |
| 97.55 | | 5.7 | 2.9 | | | | | | | | | | | | | | | 5.6 |
| .60 | 17.0 | 12.2 | 4.9 | 4.9 | 2.4 | | | | | | | | | | | | | 8.6 |
| | | | | | | | | | | | | | | | | | | 43.8 |

Table V (cont'd)
Record of the larvae of jack mackerel (*Trachurus symmetricus*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Dis. and over | Total | | | |
|-----------------------|---------------------------------|-------|-------|-------|-------|------|------|------|------|------|------|------|-------|-------|------------------|-------|-------|-------|-------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | | | 12.75 | 13.75 | 14.75 |
| Cruise 5707 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 97.65 | 5.2 | 2.6 | | | | | 2.6 | | | | | | | | | | | | 10.4 |
| .70 | 3.0 | | 5.9 | 35.4 | 11.8 | 8.8 | | | | | | | | | | | | | 64.9 |
| .75 | | | | 5.7 | 8.6 | 2.8 | | | | | | | | | | | | | 17.1 |
| .90 | | 7.5 | 5.0 | | | | | | | | | | | | | | | | 12.5 |
| 100.50 | | | | | | | 2.8 | | | | | | | | | | | | 2.8 |
| .55 | | | | | | | 2.6 | | | | | | | | | | | | 5.2 |
| .60 | | 2.7 | 5.4 | | | | | 2.7 | | | | | | | | | | | 10.8 |
| 103.50 | | | | | | | | 2.5 | 2.5 | 5.0 | 2.5 | 2.5 | | | | | | | 15.0 |
| .55 | | | 5.1 | 12.8 | 2.6 | | | | | | | | | | | | | | 20.5 |
| .60 | | | 2.7 | | | | | | | | | | | | | | | | 2.7 |
| 107.32 | | | | | | | | | | | | | | | | | | | 3.7 |
| .65 | 3.4 | 3.4 | 3.4 | 3.4 | 6.7 | 3.4 | | | | | | | | | | | | | 23.7 |
| 110.65 | | 13.0 | 4.3 | 4.3 | 8.7 | | | | | | | | | | | | | | 30.3 |
| .70 | | | | | 3.4 | | | | | | | | | | | | | | 3.4 |
| .85 | | 3.3 | | | | | | | | | | | | | | | | | 3.3 |
| 117.26 | | | | | 6.8 | | | | | | | | | | | | | | 6.8 |
| .40 | | | | | 3.4 | | | | | | | | | | | | | | 3.4 |
| 120.25 | 11.9 | 3.0 | | | | | | | | | | | | | | | | | 14.9 |
| .35 | | | | | | 2.9 | 5.8 | | | | | | | | | | | | 8.7 |
| Total | 64.7 | 153.7 | 120.5 | 119.0 | 118.6 | 58.4 | 25.6 | 35.9 | 29.2 | 5.1 | 5.0 | 2.4 | 9.1 | 2.5 | | | 2.6 | | 752.3 |
| Cruise 5708: | | | | | | | | | | | | | | | | | | | |
| 110.35 | | | | | | 2.9 | | | | | | | | | | | | | 2.9 |
| 113.35 | | | | | | 2.7 | | | | | | | | | | | | | 5.4 |
| Total | | | | | | 5.6 | | | | | | | | | | | | | 8.3 |
| Cruise 5709: | | | | | | | | | | | | | | | | | | | |
| Cruise 5710: | | | | | | | | | | | | | | | | | | | |
| 103.60 | | | | | | | | | | | | | | | | | | | 3.2 |
| 117.60 | | | | | | | | 2.0 | | | | | | | | | | | 2.0 |
| 120.35 | | 2.3 | | | | | | | | | | | | | | | | | 2.3 |
| Total | | 2.3 | | | | | | 2.0 | | | | | | | | | | | 7.5 |
| Cruise 5711: | | | | | | | | | | | | | | | | | | | |
| Cruise 5712: | | | | | | | | | | | | | | | | | | | |
| 0 | | | | | | | | | | | | | | | | | | | |
| 0 | | | | | | | | | | | | | | | | | | | |




Figure 6.--Pacific mackerel larvae, 1957: Distribution and relative abundance.

RECORD OF THE LARVAE OF THE PACIFIC MACKEREL
(PNEUMATOPHORUS DIEGO), 1957

The distribution and abundance of Pacific mackerel larvae in 1957 are illustrated in figure 6. For uniformity, the same categories of abundance are used in the distribution of Pacific mackerel larvae as in the other larval charts included in this report. The values at an individual station represent the cumulative standard haul total for all occupancies during the year.

The larvae of Pacific mackerel are reported by size in table VI; the size categories are identical to those used for jack mackerel (see p. 59 for midpoints and ranges). The data are summarized by month and area in text table 10.

Pacific mackerel larvae ranked eighteenth in abundance during 1957; thus it is one of the less abundant kinds in the CCOFI area. The number of larvae taken during a season has been fairly constant during the past three years, as is shown in the following tabulation:

| Lines | 1955 | | | 1956 | | | 1957 | | |
|---------|------------------|-------------|---------|------------------|-------------|---------|------------------|-------------|---------|
| | Occur- rences | Num- ber | Percent | Occur- rences | Num- ber | Percent | Occur- rences | Num- ber | Percent |
| 60-77 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 80-93 | 7 | 136 | 7.0 | 3 | 23 | 1.5 | 2 | 19 | 1.0 |
| 97-107 | 20 | 152 | 7.8 | 11 | 365 | 24.0 | 16 | 102 | 5.5 |
| 110-120 | 40 | 1,218 | 62.4 | 21 | 1,090 | 71.8 | 49 | 1,608 | 86.2 |
| 123-137 | 19 | 289 | 14.8 | 4 | 38 | 2.5 | 8 | 108 | 5.8 |
| 140-157 | 6 | 155 | 7.9 | 1 | 3 | 0.2 | 1 | 28 | 1.5 |
| Total | 92 | 1,950 | 99.9 | 40 | 1,519 | 100.0 | 76 | 1,865 | 100.0 |

In the regular CCOFI survey area most Pacific mackerel larvae are taken off central Baja California, especially between lines 110 and 120. During 1957, less than 7 percent of the larvae were taken to the north of this area, a lesser percentage than in 1955 or 1956. Over three-fourths of the larvae were obtained in Sebastian Viscaïno Bay, mainly during July through October.

Four cruises were made into the Gulf of California during 1957. Pacific mackerel larvae taken on these cruises will be dealt with in a separate report, but it should be noted that larvae of this species were much more abundant in the Gulf than on the outer coast.

Text table 10. --Occurrence and abundance (standard haul totals) of Pacific mackerel larvae (Pneumatophorus diego), by month and area, in hauls made during 1957

| Cruise | Central California 60-77 | | Southern California 80-93 | | Northern Baja California 97-107 | | Upper central Baja California 110-120 | | Lower central Baja California 123-137 | | Southern Baja California 140-157 | | Total | |
|---------|--------------------------|------|---------------------------|------|---------------------------------|------|---------------------------------------|--------|---------------------------------------|------|----------------------------------|------|--------|-------|
| | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- |
| reces | ber | ber | reces | ber | reces | ber | reces | ber | reces | ber | reces | ber | reces | ber |
| 5701 | - | - | - | 0 | 0 | 0 | 0 | 0 | 1 | 6 | - | - | 1 | 6 |
| 5702 | - | - | 0 | 0 | 0 | 0 | 1 | 6 | 0 | 0 | 1 | 28 | 2 | 34 |
| 5703 | - | - | 0 | 0 | 1 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 10 |
| 5704 | - | - | 0 | 0 | 4 | 41. | 1 | 3 | 0 | 0 | 0 | 0 | 5 | 44 |
| 5705 | 0 | 0 | 0 | 0 | 8 | 43 | 3 | 32 | 0 | 0 | - | - | 11 | 75 |
| 5706 | 0 | 0 | 1 | 16 | 2 | 6 | 7 | 188 | 0 | 0 | 0 | 0 | 10 | 210 |
| 5707 | 0 | 0 | 1 | 3 | 1 | 2 | 7 | 170 | 3 | 16 | - | - | 12 | 191 |
| 5708 | - | - | - | - | - | - | 18 | 654 | 2 | 8 | - | - | 20 | 662 |
| 5709 | - | - | - | - | - | - | 8 | 257 | 2 | 78 | - | - | 10 | 335 |
| 5710 | - | - | 0 | 0 | 0 | 0 | 4 | 298 | 0 | 0 | - | - | 4 | 298 |
| 5711 | 0 | 0 | 0 | 0 | 0 | 0 | - | - | - | - | - | - | 0 | 0 |
| 5712 | - | - | 0 | 0 | 0 | 0 | 0 | 0 | - | - | - | - | 0 | 0 |
| Total | 0 | 0 | 2 | 19 | 16 | 102 | 49 | 1,608 | 8 | 108 | 1 | 28 | 76 | 1,865 |
| Percent | 0 | 0 | 1.02 | 5.47 | 86.22 | 5.79 | 1.50 | 100.00 | | | | | | |

Table VI
Record of the larvae of Pacific mackerel (*Pneumatophorus diego*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Total | | | | | |
|--------------|---------------------------------|------|------|------|------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|----------------|------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | | 12.75 | 13.75 | 14.75 | 15.26 and over | |
| Cruise 5701: | | | | | | | | | | | | | | | | | | | | |
| 127.40 | | | | | | | 5.5 | | | | | | | | | | | | 5.5 | |
| Total | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | 5.5 | | | | | | | | | | | | 5.5 | |
| Cruise 5702: | | | | | | | | | | | | | | | | | | | | |
| 120.25 | 6.0 | | | | | | | | | | | | | | | | | | | 6.0 |
| 150.25 | | | | | 9.2 | 9.2 | 6.2 | 3.1 | | | | | | | | | | | | 27.7 |
| Total | | | | | | | | | | | | | | | | | | | | |
| 6.0 | | | | | 9.2 | 9.2 | 6.2 | 3.1 | | | | | | | | | | | | 33.7 |
| Cruise 5703: | | | | | | | | | | | | | | | | | | | | |
| 103.35 | | | | | | | | | | | | | | | | | | | | 5.8 |
| .40 | | | | | 4.0 | | | | | | | | | | | | | | | 4.0 |
| Total | | | | | | | | | | | | | | | | | | | | |
| 5.8 | | | | | 4.0 | | | | | | | | | | | | | | | 9.8 |
| Cruise 5704: | | | | | | | | | | | | | | | | | | | | |
| 97.32 | | | | | | | | | 17.0 | | | | | | | | | | | 17.0 |
| 100.33 | | | | | | | 10.4 | | | | | | | | | | | | | 15.6 |
| .40 | | | | | | | | 4.9 | | | | | | | | | | | | 4.9 |
| 107.40 | | | | | | | | 3.3 | | | | | | | | | | | | 3.3 |
| 110.70 | | | | | | | 3.3 | | | | | | | | | | | | | 3.3 |
| Total | | | | | | | | | | | | | | | | | | | | |
| 22.2 | 13.7 | 8.2 | | | | | | | | | | | | | | | | | | 44.1 |
| Cruise 5705: | | | | | | | | | | | | | | | | | | | | |
| 97.30 | | | | | | | | | 4.5 | | | | | | | | | | | 4.5 |
| .60 | | | | | | | | | 3.9 | | | | | | | | | | | 3.9 |
| .65 | | | | | | | | | 2.9 | | | | | | | | | | | 2.9 |
| 100.65 | | | | | | | 3.0 | 3.0 | | | | | | | | | | | | 12.0 |
| .70 | | | | | | | 3.4 | | | | | | | | | | | | | 3.4 |
| 103.45 | | | | | | | | | 2.6 | | | | | | | | | | | 2.6 |
| .60 | | | | | | | | | 3.5 | | | | | | | | | | | 3.5 |
| .65 | | | | | | | | | 5.3 | 2.6 | 2.6 | | | | | | | | | 10.5 |
| 117.45 | | | | | | | 2.3 | 2.3 | | | | | | | | | | | | 4.6 |
| .50 | | | | | | | 2.8 | 2.8 | | | | | | | | | | | | 5.6 |
| 119.33 | | | | | | | 21.4 | | | | | | | | | | | | | 21.4 |
| Total | | | | | | | | | | | | | | | | | | | | |
| 21.4 | 11.5 | 25.0 | 11.8 | 2.6 | 2.6 | | | | | | | | | | | | | | | 74.9 |

Table VI (cont'd)
Record of the larvae of Pacific mackerel (*Pneumatophorus diego*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Total | | | | | |
|--------------|---------------------------------|------|------|------|------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|----------------|-------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | | 12.75 | 13.75 | 14.75 | 15.26 and over | Dis. |
| Cruise 5706: | | | | | | | | | | | | | | | | | | | | |
| 93.55 | | | 13.1 | | | | | | | | | | | | | | | 3.3 | | 16.4 |
| 97.60 | | | | | | | | | | | | | | | | | | 3.0 | | 3.0 |
| .65 | | | | 3.5 | | | | | | | | | | | | | | | | 3.5 |
| 117.35 | | 24.5 | 68.6 | 29.4 | 4.9 | | 2.4 | | | | | | | | | | | | | 129.8 |
| .40 | | | | | 1.4 | 2.7 | | | | | | | | | | | | | | 4.1 |
| .45 | 17.9 | 23.0 | 2.6 | | | | | | | | | | | | | | | | | 43.5 |
| .70 | | | | | | | | | | | | | | | | | | | | 2.5 |
| 119.33 | | 2.7 | | | | | | | | | | | | | | | | | | 2.7 |
| 120.25 | | | | 2.5 | | | | | | | | | | | | | | | | 2.5 |
| .45 | | | | 3.2 | | | | | | | | | | | | | | | | 2.5 |
| Total | 17.9 | 50.2 | 84.3 | 38.6 | 6.3 | 2.7 | 2.4 | 2.4 | 5.5 | 3.3 | | | | | | | | | | 211.2 |
| Cruise 5707: | | | | | | | | | | | | | | | | | | | | |
| 87.55 | | | | | | | | | | | | | | | | | | | | 2.9 |
| 97.32 | | | | 2.2 | | | | | | | | | | | | | | | | 2.9 |
| 113.35 | | | | | 7.9 | | | | | | | | | | | | | | | 7.9 |
| .40 | | 6.4 | | | | | | | | | | | | | | | | | | 6.4 |
| 117.30 | | | 3.6 | | | | | | | | | | | | | | | | | 3.6 |
| 120.25 | 3.0 | 3.0 | 11.9 | 5.9 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 35.8 |
| .30 | 9.7 | 19.4 | 9.7 | 6.5 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 3.2 | 51.7 |
| .35 | | | 2.9 | 5.7 | 8.6 | 5.7 | 5.7 | 5.7 | 8.6 | 11.5 | 2.9 | | | | | | | | | 51.6 |
| .40 | 1.9 | 5.7 | 1.9 | | | 1.9 | | | 1.9 | | | | | | | | | | | 13.3 |
| 123.37 | | | | | | | | | | | | | | | | | | | | 2.9 |
| 130.45 | | 3.0 | 3.0 | | | | | | | | | | | | | | | | | 2.9 |
| 133.40 | | | 3.4 | 3.4 | | | | | | | | | | | | | | | | 6.0 |
| Total | 14.6 | 37.5 | 36.4 | 23.7 | 19.7 | 10.6 | 11.9 | 13.5 | 14.5 | 5.8 | 2.9 | | | | | | | | | 191.1 |
| Cruise 5708: | | | | | | | | | | | | | | | | | | | | |
| 110.35 | | | | | | | | | | | | | | | | | | | | 2.9 |
| 113.30 | | | 5.4 | | 2.7 | 5.4 | 5.4 | | 2.9 | | | | | | | | | | | 18.9 |
| 115.27 | | | | | 2.9 | 2.9 | 11.6 | 11.6 | 14.4 | 2.9 | | | | | | | | | | 46.3 |
| .30 | | | | | | | 3.4 | | | | | | | | | | | | | 3.4 |
| .35 | | | | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 2.8 | 11.2 |
| 117.26 | | | | | 2.9 | 8.8 | 5.8 | 38.0 | 55.4 | 17.5 | 2.9 | 5.8 | | | | | | | | 137.1 |
| .30 | | 2.9 | 2.9 | 2.9 | 2.9 | 5.9 | 14.7 | 14.7 | 14.7 | 8.8 | | | | | | | | | | 52.8 |

Table VI (cont'd)
Record of the larvae of Pacific mackerel (*Pneumatophorus diego*), 1957

| Station | Midpoint of size class (in mm.) | | | | | | | | | | | | | | Total | | | | |
|-----------------------|---------------------------------|------|------|------|------|------|------|-------|-------|------|------|------|-------|-------|-------|-------|-------|-------|----------------|
| | 2.00 | 2.50 | 3.00 | 3.50 | 4.00 | 4.50 | 5.00 | 5.75 | 6.75 | 7.75 | 8.75 | 9.75 | 10.75 | 11.75 | | 12.75 | 13.75 | 14.75 | 15.26 and over |
| Cruise 5708 (cont'd): | | | | | | | | | | | | | | | | | | | |
| 117.35 | | | | | 6.4 | 6.4 | 22.4 | 9.6 | 6.4 | | | | | | | | | | 51.2 |
| .40 | | | | | | | 2.7 | | | | | | | | | | | | 2.7 |
| 118525 | 9.1 | 6.0 | 6.0 | 3.0 | | | | | | | | | | | | | | | 24.1 |
| .30 | | | | 3.2 | | | | | | | | | | | | | | | 9.6 |
| .35 | 6.2 | 25.0 | | | | | | | 3.2 | | | | | | | | | | 31.2 |
| 119.33 | | 5.8 | 5.8 | 2.9 | 8.7 | 14.5 | 11.6 | 23.2 | 2.9 | 2.9 | | | | | | | | | 78.3 |
| 120.25 | 3.0 | | 5.9 | 11.8 | 8.9 | 3.0 | 3.0 | | 3.0 | | | | | | | | | | 41.6 |
| .30 | | | | | 3.3 | 13.4 | | 6.7 | | | | | | | | | | | 26.7 |
| .35 | | | | 2.1 | 4.3 | 2.1 | 6.4 | 2.1 | | | | | | | | | | | 19.1 |
| .40 | | | 2.0 | 6.1 | 4.1 | 14.2 | 12.2 | 30.4 | 16.3 | 4.0 | 2.0 | | | | | | | | 91.3 |
| .45 | | | | | 2.9 | 2.9 | | | | | | | | | | | | | 5.8 |
| 127.34 | | | | | | | 2.9 | 2.9 | | | | | | | | | | | 5.8 |
| 137.23 | | | | | 2.7 | | | | | | | | | | | | | | 2.7 |
| Total | 18.3 | 36.8 | 28.0 | 34.8 | 43.4 | 79.3 | 77.8 | 160.4 | 116.3 | 48.5 | 13.3 | 5.8 | | | | | | | 662.7 |
| Cruise 5709: | | | | | | | | | | | | | | | | | | | |
| 113.30 | | | | | | | | | 4.7 | 2.4 | | | | | | | | | 7.1 |
| 117.26 | | | | | | | 2.6 | 2.6 | 2.6 | | | | | | | | | | 7.8 |
| 118525 | | | | | | | | 2.5 | | | | | | | | | | | 2.5 |
| .30 | | | | | 22.5 | 3.8 | 18.8 | 22.5 | 37.5 | 22.5 | 26.3 | 3.8 | | | | | | | 161.5 |
| 119.33 | 6.0 | 3.0 | 3.0 | 14.9 | 8.9 | 3.0 | | 3.0 | | | | | | | | | | | 3.8 |
| 120.30 | | 3.1 | | | | | | 3.0 | | | | | 3.0 | | | | | | 44.8 |
| .35 | 3.0 | 3.0 | 3.0 | 6.0 | | | 3.1 | 3.1 | | | | | | | | | | | 9.3 |
| .40 | | | | | | | | 3.0 | | | | | 3.0 | | | | | | 18.0 |
| 123.37 | 3.4 | 3.4 | 10.1 | 20.2 | 23.6 | 13.5 | | | | | | | | | | | | | 6.0 |
| 127.34 | | | | | | | | | | | | | | | | | | | 74.2 |
| | | | | | | | | | | | | | | | | | | | 3.4 |
| Total | 12.4 | 35.0 | 13.9 | 59.9 | 55.0 | 57.4 | 28.2 | 40.5 | 14.1 | 5.4 | 3.0 | 3.0 | 3.0 | | | | | | 334.6 |
| Cruise 5710: | | | | | | | | | | | | | | | | | | | |
| 117.26 | | | 3.0 | 3.0 | | | | | 3.0 | | | | | | | | | | 9.0 |
| 119.33 | | 2.6 | 7.8 | 7.8 | 10.4 | 7.8 | 2.6 | 2.6 | 2.6 | | | | | | | | | | 44.2 |
| 120.30 | 8.4 | 21.1 | 29.5 | 33.8 | 25.3 | 8.4 | 2.1 | | | | | | | | | | | | 128.6 |
| .35 | 2.3 | 13.6 | 34.0 | 38.6 | 20.4 | 6.8 | | | | | | | | | | | | | 115.7 |
| Total | 10.7 | 37.3 | 74.3 | 83.2 | 56.1 | 23.0 | 7.7 | 2.6 | 2.6 | | | | | | | | | | 297.5 |
| Cruise 5711: | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | 0 |
| Cruise 5712: | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | 0 |




Figure 7. --Hake larvae, 1957: Distribution and relative abundance.

RECORD OF THE LARVAE OF HAKE (MERLUCCIIUS PRODUCTUS), 1957

The records of hake larvae contained in table VII are the numbers of larvae (standard haul totals) taken in each plankton collection made in 1957. Length measurements were not made on hake larvae. The distribution and relative abundance of hake larvae in 1957 are shown in figure 7. The basic data contained in table VII are summarized by month and area in text table 11.

Hake larvae constituted 16 percent of the fish larvae taken on CCOFI cruises during 1957. They were exceeded in abundance only by anchovy larvae, as was also the situation during each of the preceding five years (1952-1956). Monthly abundance and occurrences of hake larvae in 1957 are compared with similar data for 1956 in the following summary:

Monthly abundance of hake larvae in 1956 and 1957

| Months | 1956 | | | 1957 | | |
|-----------|------------------|----------------------------|---------------------|------------------|----------------------------|---------------------|
| | Occur- rences | Standard haul totals | Percent of total | Occur- rences | Standard haul totals | Percent of total |
| January | 47 | 33,376 | 37.14 | 19 | 1,278 | 1.63 |
| February | 74 | 39,746 | 44.23 | 67 | 37,998 | 48.54 |
| March | 111 | 15,010 | 16.70 | 98 | 34,589 | 44.18 |
| April | 64 | 1,047 | 1.17 | 81 | 4,047 | 5.17 |
| May | 32 | 301 | 0.33 | 62 | 260 | 0.33 |
| June | 16 | 195 | 0.22 | 16 | 88 | 0.11 |
| July | 7 | 90 | 0.10 | 4 | 16 | 0.02 |
| August | 3 | 47 | 0.05 | 1 | 3 | 0.01 |
| September | 0 | 0 | 0 | 0 | 0 | 0 |
| October | 1 | 6 | 0.01 | 0 | 0 | 0 |
| November | 0 | 0 | 0 | 1 | 4 | 0.01 |
| December | 5 | 39 | 0.04 | 0 | 0 | 0 |
| Total | 360 | 89,857 | 99.99 | 349 | 78,283 | 100.00 |

Most hake larvae were taken during a two-month period, February and March. Due to the abbreviated coverage in January 1957 (only 57 stations occupied, all off Baja California), hake abundance must be underestimated for this month. Over 40 percent of the larvae were taken off California (station lines 60-93) in 1957, as compared to less than 3 percent in this area in 1956 and approximately 8 percent in 1955.

Text table 11.--Occurrence and abundance (standard haul totals), of hake larvae (Merluccius productus), by month and area, in hauls made during 1957

| Cruise | Central California 60-77 | | Southern California 80-93 | | Northern Baja California 97-107 | | Upper central Baja California 110-120 | | Lower central Baja California 123-137 | | Southern Baja California 140-157 | | Total | |
|---------|--------------------------|------|---------------------------|--------|---------------------------------|--------|---------------------------------------|--------|---------------------------------------|-------|----------------------------------|-------|--------|--------|
| | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- |
| | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber | rences | ber |
| 5701 | - | - | - | - | 0 | 0 | 8 | 965 | 11 | 313 | - | - | 19 | 1,278 |
| 5702 | - | 8 | 377 | | 18 | 24,517 | 16 | 10,233 | 16 | 1,385 | 9 | 1,486 | 67 | 37,998 |
| 5703 | - | 31 | 28,712 | | 14 | 773 | 20 | 3,482 | 22 | 1,254 | 11 | 368 | 98 | 34,589 |
| 5704 | - | 35 | 2,731 | | 21 | 806 | 15 | 431 | 7 | 58 | 3 | 21 | 81 | 4,047 |
| 5705 | 2 | 11 | 23 | 128 | 14 | 93 | 4 | 21 | 2 | 7 | - | - | 45 | 260 |
| 5706 | 2 | 6 | 3 | 16 | 9 | 58 | 2 | 8 | 0 | 0 | 0 | 0 | 16 | 88 |
| 5707 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 13 | 1 | 3 | - | - | 4 | 16 |
| 5708 | - | - | - | - | - | - | 1 | 3 | 0 | 0 | - | - | 1 | 3 |
| 5709 | - | - | - | - | - | - | 0 | 0 | 0 | 0 | - | - | 0 | 0 |
| 5710 | - | - | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | - | - | 0 | 0 |
| 5711 | 0 | 0 | 1 | 4 | 0 | 0 | - | - | - | - | - | - | 1 | 4 |
| 5712 | - | - | 0 | 0 | 0 | 0 | 0 | 0 | - | - | - | - | 0 | 0 |
| Total | 4 | 17 | 101 | 31,968 | 76 | 26,247 | 69 | 15,156 | 59 | 3,020 | 23 | 1,875 | 332 | 78,283 |
| Percent | 0.02 | | 40.84 | | 33.53 | | 19.36 | | 3.86 | | 2.40 | | 100.01 | |

Table VII
Record of the larvae of hake (Merluccius productus), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 60.50 | - | - | - | - | | | NQ | - | - | - | - | - |
| .55 | - | - | - | - | | | | - | - | - | - | - |
| .60 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | | | | - | - | - | - | - |
| 63.52 | - | - | - | - | | | | - | - | - | - | - |
| .55 | - | - | - | - | | | | - | - | - | - | - |
| .60 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | - | | | - | - | - | - | - |
| 67.50 | - | - | - | - | | | NS | - | - | - | - | - |
| .55 | - | - | - | - | | | | - | - | - | - | - |
| .60 | - | - | - | - | 4 | | | - | - | - | - | - |
| .70 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | - | 3 | | - | - | - | - | - |
| 70.52 | - | - | - | - | 7 | | | - | - | - | - | - |
| .55 | - | - | - | - | | | | - | - | - | - | - |
| .60 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | | | | - | - | - | - | - |
| 73.50 | - | - | - | - | | | | - | - | - | - | - |
| .55 | - | - | - | - | | | | - | - | - | - | - |
| .60 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | - | | | - | - | - | - | - |
| 77.50 | - | - | - | - | | | | - | - | - | - | - |
| .55 | - | - | - | - | | | 3 | - | - | - | - | - |
| .60 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | - | | | - | - | - | - | - |

Table VII (cont'd)
Record of the larvae of hake (Merluccius productus), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 80.51 | - | | 6 | | | | | - | - | | | - |
| .55 | - | | 212 | 15 | | | | - | - | | | |
| .60 | - | | 133 | | | 6 | | - | - | | | |
| .70 | - | | | 26 | | | | - | - | | | |
| .80 | - | | 168 | 9 | | | | - | - | NS | | - |
| .90 | - | | 59 | 6 | 4 | | | - | - | | | - |
| 82.47 | - | | 11 | 5 | | | | - | - | | | - |
| 83.40 | - | | | 16 | | | | - | - | | | - |
| .43 | - | | 3 | | | | | - | - | | | - |
| .48 | - | - | 20 | - | - | - | - | - | - | - | - | - |
| .51 | - | | - | 5 | | | NS | - | - | | | - |
| .52 | - | | 130 | - | - | - | - | - | - | - | - | - |
| .55 | - | - | - | - | 7 | 3 | | - | - | - | - | - |
| .60 | - | | 706 | 52 | 6 | | | - | - | | | - |
| .65 | - | - | - | - | - | | | - | - | - | - | - |
| .70 | - | - | 57 | 30 | | | | - | - | - | - | - |
| .75 | - | - | - | - | - | | | - | - | - | - | - |
| .80 | - | - | - | 48 | 6 | | | - | - | - | - | - |
| .85 | - | - | - | - | - | | | - | - | - | - | - |
| .90 | - | - | - | 9 | 3 | | NS | - | - | - | - | - |
| 86.46 | - | | - | 15 | - | - | - | - | - | - | - | - |
| 87.35 | - | 16 | 6 | 8 | | | | - | - | - | - | - |
| .38 | - | | - | - | - | - | - | - | - | - | - | - |
| .40 | - | 98 | 11 | 49 | | | NS | - | - | - | - | - |
| .45 | - | - | 44 | - | | | | - | - | - | - | - |
| .50 | - | | 23 | 193 | | | | - | - | - | - | - |
| .55 | - | | - | - | | | | - | - | - | - | - |
| .60 | - | | | 12 | | | | - | - | | 4 | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | | 341 | 284 | 3 | | | - | - | - | - | - |
| .75 | - | - | - | - | 7 | | | - | - | - | - | - |
| .80 | - | - | - | 13 | 12 | | NS | - | - | - | - | - |
| .85 | - | - | - | - | 11 | | | - | - | - | - | - |
| .90 | - | - | - | 101 | 3 | | | - | - | - | - | - |
| 90.28 | - | 6 | 16 | 28 | | | | - | - | - | - | - |
| .30 | - | | 60 | 8 | | | | - | - | - | - | - |
| .37 | - | | 72 | | | | | - | - | - | - | - |
| .45 | - | | 12 | 53 | | 7 | | - | - | - | - | - |

Table VII (cont'd)
Record of the larvae of hake (Merluccius productus), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 90.50 | - | - | - | - | 3 | | | - | - | | | |
| .55 | - | 36 | 49 | 217 | | | | - | - | | | |
| .60 | - | 18 | 490 | 88 | | | | - | - | | | |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | | 24718 | 336 | 3 | | | - | - | - | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | | 160 | 12 | | | | - | - | - | - | - |
| .85 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | | 976 | 99 | 6 | | | - | - | - | - | - |
| 93.27 | - | | 15 | | | | | - | - | - | - | - |
| .30 | - | 43 | 24 | 150 | 2 | | | - | - | - | - | - |
| .35 | - | - | - | 62 | 2 | | | - | - | - | - | - |
| .40 | - | 111 | 45 | 60 | 11 | | | - | - | - | - | - |
| .45 | - | - | - | 62 | 3 | | | - | - | - | - | - |
| .50 | - | 49 | 18 | 358 | 9 | | | - | - | - | - | - |
| .55 | - | - | - | 197 | 6 | | | - | - | - | - | - |
| .60 | - | | 91 | 80 | | | | - | - | - | - | - |
| .65 | - | - | - | - | 10 | | | - | - | - | - | - |
| .70 | - | | 36 | | 5 | | | - | - | - | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | NS | | | | - | - | - | - | - |
| .85 | - | - | - | - | 3 | | | - | - | - | - | - |
| .90 | - | - | - | 25 | 3 | | | - | - | - | - | - |
| 97.30 | - | 11 | 5 | 3 | | | | - | - | - | - | - |
| .32 | - | | - | 28 | | | | - | - | - | - | - |
| .35 | - | - | - | - | - | - | - | - | - | - | - | - |
| .40 | - | 80 | 9 | | | | | - | - | - | - | - |
| .45 | - | - | - | 12 | | | | - | - | - | - | - |
| .50 | - | 1508 | 393 | NS | 3 | | | - | - | - | - | - |
| .55 | - | - | - | NS | | | | - | - | - | - | - |
| .60 | - | 1529 | - | 77 | | | | - | - | - | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | 76 | - | 3 | | | | - | - | - | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | 7 | | | - | - | - | - | - |
| .85 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | 61 | | | | - | - | - | - | - |

Table VII (cont'd)
Record of the larvae of hake (Merluccius productus), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 100.29 | - | 328 | 10 | 52 | 4 | | | - | - | | - | - |
| .30 | - | - | - | - | - | - | - | - | - | | - | - |
| .33 | - | 907 | 14 | 52 | | | | - | - | - | - | - |
| .35 | - | - | - | - | - | - | - | - | - | | - | - |
| .40 | - | 941 | 21 | 74 | | | | - | - | | - | - |
| .45 | - | - | - | 25 | | 12 | | - | - | - | - | - |
| .50 | - | | | | 5 | | | - | - | | - | - |
| .55 | - | - | - | 17 | | | | - | - | - | - | - |
| .60 | | 24 | 3 | 39 | | | | - | - | | - | - |
| .65 | - | - | - | - | 3 | | | - | - | - | - | - |
| .70 | - | | - | 5 | | | - | - | - | | - | - |
| .75 | - | - | - | - | | | | - | - | | - | - |
| .80 | - | | - | | | | - | - | - | | - | - |
| .85 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | | - | 16 | | | | - | - | - | - | - |
| 103.30 | | 884 | 88 | 76 | | 3 | | - | - | | - | - |
| .35 | | 12096 | 72 | | 10 | 6 | | - | - | | - | - |
| .38 | | 157 | - | - | - | - | - | - | - | - | - | - |
| .40 | | 232 | 8 | | 17 | 12 | | - | - | | - | - |
| .45 | - | - | - | | 5 | 8 | | - | - | - | - | - |
| .50 | | 948 | 9 | 3 | | | | - | - | | - | - |
| .55 | - | - | - | | | | | - | - | - | - | - |
| .60 | | | NS | | 4 | 3 | | - | - | | - | - |
| .65 | - | - | - | - | 11 | 4 | | - | - | - | - | - |
| .70 | - | | | | | | | - | - | | - | - |
| .75 | - | - | - | - | | 3 | | - | - | - | - | - |
| .80 | - | - | - | | | | - | - | - | | - | - |
| .85 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | | | | - | - | - | - | - | - |
| 107.32 | | 4196* | 123 | 38 | | - | | - | - | NQ | - | - |
| .35 | | | | 32 | 8 | | | - | - | | - | - |
| .40 | | 289 | 15 | 110 | | | | - | - | | - | - |
| .45 | - | - | - | 80 | 3 | | | - | - | - | - | - |
| .50 | | 294 | 3 | | 2 | | | - | - | | - | - |
| .55 | - | - | - | 3 | 11 | | | - | - | - | - | - |
| .60 | | 17 | | | | 7 | | - | - | | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | | | | | | | - | - | | - | - |

* - Samples combined. Values adjusted accordingly.

Table VII (cont'd)
 Record of the larvae of hake (Merluccius productus), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 107.75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .85 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | | | | - | - | - | - | - |
| 110.33 | - | 3633 | 14 | 21 | | | | | | | - | - |
| .35 | - | 1545 | 260 | 13 | 6 | | | | | | - | - |
| .40 | | 162 | 7 | 56 | | | | | | | - | - |
| .45 | - | - | - | 27 | | | | - | - | - | - | - |
| .50 | | 3 | 15 | 9 | 3 | | | - | - | - | - | - |
| .55 | - | - | - | 3 | | | | - | - | - | - | - |
| .60 | | | 8 | 16 | | | | - | - | - | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | | | 241 | | | | - | - | - | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | | | | | | | - | - | - | - | - |
| .85 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | 12 | - | 3 | | | | - | - | - | - | - |
| 113.30 | 40 | 97 | 23 | 9 | 9 | | | | | | - | - |
| .35 | 16 | 370 | 571 | 4 | | | | | | | - | - |
| .40 | | 20 | 602 | 6 | | | | | | | - | - |
| .45 | - | - | - | - | | | | - | - | - | - | - |
| .50 | | | 30 | | | | | - | - | - | - | - |
| .55 | - | - | - | - | | | | - | - | - | - | - |
| .60 | | | 3 | | | | | - | - | - | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | | | | | | | | - | - | - | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .85 | - | - | - | - | - | | | - | - | - | - | - |
| .90 | - | - | - | - | - | | | - | - | - | - | - |
| 115.27 | - | - | - | - | - | - | - | | | - | - | - |
| .30 | - | - | - | - | - | - | - | | | - | - | - |
| .35 | - | - | - | - | - | - | - | | | - | - | - |
| .40 | - | - | - | - | - | - | - | | | - | - | - |
| 117.26 | | 396 | 3 | | | 6 | | | | | - | - |
| .30 | 148 | 2441 | 425 | 19 | | | 4 | | | | - | - |
| .35 | 619 | 961 | 607 | | | | | 3 | | | - | - |
| .40 | | 53 | 33 | | | | | | | | - | - |

Table VII (cont'd)
 Record of the larvae of hake (Merluccius productus), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 117.45 | - | - | - | | | | | - | - | - | - | - |
| .50 | 3 | 11 | 237 | | | | | - | - | - | - | - |
| .55 | - | - | - | | | | | - | - | - | - | - |
| .60 | | | 15 | | | | | - | - | - | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | | | | | | | | - | - | - | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | | | | | - | - | - | - | - |
| .85 | - | - | - | - | - | | - | - | - | - | - | - |
| .90 | - | - | - | - | - | | - | - | - | - | - | - |
| 118.39 | - | - | 115 | | 3 | | | - | - | | - | - |
| 118.525 | - | - | - | - | - | - | - | | | - | - | - |
| .30 | - | - | - | - | - | - | - | | | - | - | - |
| .35 | - | - | - | - | - | - | - | | | - | - | - |
| 119.33 | - | - | 376 | | | | | | | | | |
| 120.25 | 9 | 161 | 30 | 2 | | 2 | | | | | - | - |
| .30 | 94 | 363 | 108 | | | | | 6 | | | - | - |
| .35 | | 5 | | | | | | 3 | | | - | - |
| .40 | | | | - | | | | | | | - | - |
| .45 | 36 | | - | - | | | | | | | - | - |
| .50 | | | - | 2 | | | | - | - | | - | - |
| .55 | | | - | | | | | - | - | - | - | - |
| .60 | | | | | | | | - | - | | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | | | | | | | | - | - | | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | | | | | | | - | - | | - | - |
| .85 | - | - | - | - | - | | - | - | - | - | - | - |
| .90 | - | - | - | - | - | | - | - | - | | - | - |
| 123.37 | 7 | 236 | 143 | | | | | | | | - | - |
| .42 | 3 | 3 | 126 | | 3 | | | | | | - | - |
| .45 | - | - | - | - | - | - | - | | | - | - | - |
| .50 | | | | | - | | | - | - | | - | - |
| .55 | - | | | | - | | | - | - | - | - | - |
| .60 | | | | | - | | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | | - | - |

Table VII (cont'd)
Record of the larvae of hake (Merluccius productus), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 127.34 | | 21 | 186 | | | | | | | | - | - |
| .40 | 11 | 84 | 8 | | | | | | | | - | - |
| .45 | - | 60 | 22 | 9 | 4 | | | | | | - | - |
| .50 | | 5 | 6 | | - | | | - | - | | - | - |
| .55 | - | | 7 | | - | | | - | - | - | - | - |
| .60 | | | 38 | | - | | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | | - | - |
| 130.30 | | 6 | 64 | | | | | | | | - | - |
| .35 | 53 | 558 | 12 | | | | | | | | - | - |
| .40 | 116 | 190 | 16 | | | | | | | | - | - |
| .45 | - | - | - | | - | | | | | - | - | - |
| .50 | 3 | | 3 | | | | | - | - | | - | - |
| .55 | - | - | - | | - | | 3 | - | - | - | - | - |
| .60 | | | 3 | | | | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | | - | - |
| 133.25 | 50 | | 92 | | | | | | | | - | - |
| .30 | 11 | 38 | 98 | 19 | | | | | | | - | - |
| .35 | 6 | - | 35 | | - | | | - | - | | - | - |
| .40 | 5 | 8 | 40 | 12 | | | | - | - | | - | - |
| .45 | - | - | - | | - | | | - | - | - | - | - |
| .50 | - | | | 3 | | | | - | - | | - | - |
| .55 | - | - | - | | - | - | - | - | - | - | - | - |
| .60 | - | - | - | | - | - | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | | - | - |
| 137.23 | | 28 | 29 | 7 | | | | | | | - | - |
| .30 | | 10 | 8 | 3 | | | | | | | - | - |
| .35 | 48 | 113 | 134 | - | - | | | - | - | | - | - |
| .40 | - | 22 | 178 | | NS | | | - | - | | - | - |
| .45 | - | - | - | 5 | - | | | - | - | - | - | - |
| .50 | - | 5 | 6 | | | | | - | - | | - | - |
| .55 | - | - | - | | - | - | - | - | - | - | - | - |
| .60 | - | - | - | | - | - | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | | - | - |

Table VII (cont'd)
 Record of the larvae of hake (Merluccius productus), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 140.30 | - | 432 | 8 | | - | | - | - | - | - | - | - |
| .35 | - | 355 | 11 | | - | | - | - | - | - | - | - |
| .40 | - | 8 | 16 | | - | | - | - | - | - | - | - |
| .45 | - | - | - | | - | - | - | - | - | - | - | - |
| .50 | - | 20 | 3 | | - | | - | - | - | - | - | - |
| .55 | - | - | - | | - | - | - | - | - | - | - | - |
| .60 | - | - | - | | - | - | - | - | - | - | - | - |
| .70 | - | - | - | | - | - | - | - | - | - | - | - |
| .80 | - | - | - | | - | - | - | - | - | - | - | - |
| 143.26 | - | 8 | | 3 | - | - | - | - | - | - | - | - |
| .30 | - | 571 | | | - | | - | - | - | - | - | - |
| .35 | - | 33 | 20 | | - | | - | - | - | - | - | - |
| .40 | - | | | | - | | - | - | - | - | - | - |
| .50 | - | | | - | - | | - | - | - | - | - | - |
| 147.20 | - | 12 | 3 | 6 | - | - | - | - | - | - | - | - |
| .25 | - | 47 | 72 | 12 | - | - | - | - | - | - | - | - |
| .30 | - | | 200 | | - | - | - | - | - | - | - | - |
| .35 | - | - | - | | - | - | - | - | - | - | - | - |
| .40 | - | | 3 | | - | - | - | - | - | - | - | - |
| .45 | - | - | - | | - | - | - | - | - | - | - | - |
| .50 | - | - | - | | - | - | - | - | - | - | - | - |
| .55 | - | - | - | | - | - | - | - | - | - | - | - |
| .60 | - | - | - | | - | - | - | - | - | - | - | - |
| .70 | - | - | - | | - | - | - | - | - | - | - | - |
| .80 | - | - | - | | - | - | - | - | - | - | - | - |
| 148.20 | - | - | - | - | - | - | - | - | - | - | - | - |
| .25 | - | - | - | - | - | - | - | - | - | - | - | - |
| .30 | - | - | - | - | - | - | - | - | - | - | - | - |
| .40 | - | - | - | - | - | - | - | - | - | - | - | - |
| .50 | - | - | - | - | - | - | - | - | - | - | - | - |
| 150.16 | - | - | - | - | - | - | - | - | - | - | - | - |
| .19 | - | | | - | - | - | - | - | - | - | - | - |
| .20 | - | - | - | | - | - | - | - | - | - | - | - |
| .25 | - | | | | - | - | - | - | - | - | - | - |
| .30 | - | - | 30 | | - | - | - | - | - | - | - | - |
| .35 | - | | - | | - | - | - | - | - | - | - | - |
| .40 | - | | 2 | | - | - | - | - | - | - | - | - |
| Total | 1278 | 37998 | 34589 | 4047 | 260 | 88 | 16 | 3 | 0 | 0 | 4 | 0 |

Note: No hake found on lines 153 and 157.

RECORD OF THE LARVAE OF ROCKFISH (SEBASTODES SPP.), 1957

Larvae of the genus Sebastes are grouped together as rockfish larvae. Forty-nine species of rockfish are known to occur off California.

As in previous seasons, rockfish larvae occurred in greatest abundance off southern California (station lines 80-93), as is shown in text table 12, and the following tabulation:

| Lines | Total samples taken | Occurrences of rockfish larvae | Percent occurrence | Total number of larvae taken | Percent taken in each area | Average number per haul |
|---------|---------------------|--------------------------------|--------------------|------------------------------|----------------------------|-------------------------|
| 60-77 | 101 | 68 | 67.3 | 3,170 | 8.7 | 31.3 |
| 80-93 | 364 | 228 | 62.7 | 23,163 | 63.5 | 63.6 |
| 97-107 | 287 | 90 | 31.4 | 3,779 | 10.4 | 13.2 |
| 110-120 | 360 | 109 | 30.3 | 4,878 | 13.4 | 13.5 |
| 123-137 | 267 | 63 | 23.6 | 1,466 | 4.0 | 5.5 |
| 140-157 | 102 | 3 | 2.9 | 17 | <0.1 | 0.2 |
| Total | 1,481 | 561 | 37.9 | 36,473 | 100.0 | 24.6 |

Rockfish larvae were more abundant off southern California (station lines 80-93) in 1957 than during the previous two seasons, with respect to both the percentage of the total larvae taken and the absolute number of larvae taken.

| Lines | 1955 | | 1956 | | 1957 | |
|---------|--------|---------|--------|---------|--------|---------|
| | Number | Percent | Number | Percent | Number | Percent |
| 60-77 | 2,893 | 10.2 | 1,981 | 6.8 | 3,170 | 8.7 |
| 80-93 | 13,503 | 47.8 | 14,674 | 50.4 | 23,163 | 63.5 |
| 97-107 | 3,721 | 13.2 | 4,703 | 16.1 | 3,779 | 10.4 |
| 110-120 | 6,336 | 22.4 | 6,306 | 21.6 | 4,878 | 13.4 |
| 123-137 | 1,796 | 6.3 | 1,424 | 4.9 | 1,466 | 4.0 |
| 140-157 | 24 | 0.1 | 56 | 0.2 | 17 | <0.1 |
| Total | 28,273 | 100.0 | 29,144 | 100.0 | 36,473 | 100.0 |

A record of the rockfish larvae taken in each plankton collection made in 1957 is given in table VIII.

Text table 12.--Occurrence and abundance (standard haul totals) of rockfish larvae (*Sebastes* spp.), by month and area, in hauls made during 1957

| Cruise | Central California 60-77 | | Southern California 80-93 | | Northern Baja California 97-107 | | Upper central Baja California 110-120 | | Lower central Baja California 123-137 | | Southern Baja California 140-157 | | Total | |
|---------|--------------------------|-------|---------------------------|--------|---------------------------------|-------|---------------------------------------|-------|---------------------------------------|-------|----------------------------------|------|--------|--------|
| | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- | occur- | num- |
| ences | ber | ber | ences | ber | ences | ber | ences | ber | ences | ber | ences | ber | ences | ber |
| 5701 | - | - | - | - | 4 | 213 | 8 | 693 | 6 | 104 | - | - | 18 | 1,010 |
| 5702 | - | - | 26 | 8,372 | 10 | 855 | 16 | 1,636 | 6 | 138 | 0 | 0 | 58 | 11,001 |
| 5703 | - | - | 24 | 3,166 | 8 | 841 | 15 | 1,778 | 14 | 789 | 1 | 11 | 62 | 6,585 |
| 5704 | - | - | 31 | 5,760 | 17 | 970 | 17 | 255 | 8 | 234 | 2 | 6 | 75 | 7,225 |
| 5705 | 20 | 995 | 36 | 2,894 | 22 | 579 | 20 | 267 | 8 | 84 | - | - | 106 | 4,819 |
| 5706 | 23 | 1,293 | 38 | 1,282 | 13 | 91 | 11 | 52 | 4 | 24 | 0 | 0 | 89 | 2,742 |
| 5707 | 23 | 836 | 26 | 385 | 7 | 39 | 8 | 62 | 9 | 61 | - | - | 73 | 1,383 |
| 5708 | - | - | - | - | - | - | 5 | 34 | 3 | 13 | - | - | 8 | 47 |
| 5709 | - | - | - | - | - | - | 4 | 16 | 3 | 12 | - | - | 7 | 28 |
| 5710 | - | - | 15 | 184 | 4 | 22 | 4 | 18 | 2 | 7 | - | - | 25 | 231 |
| 5711 | 2 | 46 | 20 | 483 | 3 | 57 | - | - | - | - | - | - | 25 | 586 |
| 5712 | - | - | 12 | 637 | 2 | 112 | 1 | 67 | - | - | - | - | 15 | 816 |
| Total | 68 | 3,170 | 228 | 23,163 | 90 | 3,779 | 109 | 4,878 | 63 | 1,466 | 3 | 17 | 561 | 36,473 |
| Percent | 8.69 | | 63.51 | | 10.36 | | 13.37 | | 4.02 | | 0.05 | | 100.00 | |

Table VIII
Record of the larvae of rockfish (Sebastes spp.), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 60.50 | - | - | - | - | 18 | | NQ | - | - | - | - | - |
| .55 | - | - | - | - | 34 | 12 | 81 | - | - | - | - | - |
| .60 | - | - | - | - | 71 | 3 | 46 | - | - | - | - | - |
| .70 | - | - | - | - | 14 | 45 | 72 | - | - | - | - | - |
| .80 | - | - | - | - | | 47 | - | - | - | - | - | - |
| .90 | - | - | - | - | | | - | - | - | - | - | - |
| 63.52 | - | - | - | - | 78 | 152 | 21 | - | - | - | - | - |
| .55 | - | - | - | - | 110 | 255 | 101 | - | - | - | - | - |
| .60 | - | - | - | - | 33 | 143 | 43 | - | - | - | - | - |
| .70 | - | - | - | - | | 27 | 49 | - | - | - | - | - |
| .80 | - | - | - | - | | 4 | 4 | - | - | - | - | - |
| .90 | - | - | - | - | - | | - | - | - | - | - | - |
| 67.50 | - | - | - | - | 23 | 105 | NS | - | - | - | - | - |
| .55 | - | - | - | - | | 45 | 140 | - | - | - | - | - |
| .60 | - | - | - | - | 46 | 42 | 17 | - | - | - | - | - |
| .70 | - | - | - | - | 12 | | 8 | - | - | - | - | - |
| .80 | - | - | - | - | | 3 | 6 | - | - | - | - | - |
| .90 | - | - | - | - | - | | - | - | - | - | - | - |
| 70.52 | - | - | - | - | 70 | | 87 | - | - | - | - | - |
| .55 | - | - | - | - | | | | - | - | - | - | - |
| .60 | - | - | - | - | 17 | 19 | 7 | - | - | - | - | - |
| .70 | - | - | - | - | 67 | 36 | 4 | - | - | - | - | - |
| .80 | - | - | - | - | | | - | - | - | - | - | - |
| .90 | - | - | - | - | 3 | 6 | - | - | - | - | - | - |
| 73.50 | - | - | - | - | 18 | 21 | 5 | - | - | - | 27 | - |
| .55 | - | - | - | - | 37 | 124 | 41 | - | - | - | 19 | - |
| .60 | - | - | - | - | 123 | 60 | 6 | - | - | - | - | - |
| .70 | - | - | - | - | | 19 | 30 | - | - | - | - | - |
| .80 | - | - | - | - | | 22 | - | - | - | - | - | - |
| .90 | - | - | - | - | - | | - | - | - | - | - | - |
| 77.50 | - | - | - | - | 33 | | 3 | - | - | - | - | - |
| .55 | - | - | - | - | 75 | 84 | 3 | - | - | - | - | - |
| .60 | - | - | - | - | 113 | 19 | 43 | - | - | - | - | - |
| .70 | - | - | - | - | | | 19 | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | - | | | - | - | - | - | - |

Table VIII (cont'd)
Record of the larvae of rockfish (Sebastes spp.), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 80.51 | - | 178 | 9 | 34 | | | | - | - | | - | 18 |
| .55 | - | 3659 | 94 | 268 | 11 | | 34 | - | - | 12 | 34 | 273 |
| .60 | - | 639 | 24 | 68 | | 22 | 32 | - | - | 3 | 26 | 33 |
| .70 | - | 22 | | | 57 | 13 | | - | - | | 3 | 27 |
| .80 | - | | | | | 6 | | - | - | NS | - | |
| .90 | - | 12 | | | | | | - | - | | - | 3 |
| 82.47 | - | 32 | 52 | 113 | 252 | 7 | 40 | - | - | | 19 | - |
| 83.40 | - | 67 | 6 | 8 | | | 2 | - | - | | 4 | - |
| .43 | - | 144 | 8 | 66 | 20 | 5 | 6 | - | - | 11 | 28 | - |
| .48 | - | - | 217 | - | - | - | - | - | - | - | - | - |
| .51 | - | 1483 | - | 207 | 66 | 39 | 17 | - | - | 6 | 33 | - |
| .52 | - | 401 | 321 | - | - | - | - | - | - | - | - | - |
| .55 | - | - | - | - | 109 | | 11 | - | - | - | - | - |
| .60 | - | 14 | | 60 | 6 | 24 | 27 | - | - | | | - |
| .65 | - | - | - | - | - | 3 | | - | - | - | - | - |
| .70 | - | - | | | | | | - | - | | | - |
| .75 | - | - | - | - | - | 42 | | - | - | - | - | - |
| .80 | - | - | - | | 14 | 13 | | - | - | | | - |
| .85 | - | - | - | - | - | | | - | - | - | - | - |
| .90 | - | - | - | | | | NS | - | - | | - | 3 |
| 86.46 | - | 49 | - | 192 | - | - | - | - | - | - | - | - |
| 87.35 | - | 69 | 156 | 141 | 41 | | | - | - | 3 | 4 | - |
| .38 | - | 61 | - | - | - | - | - | - | - | - | - | - |
| .40 | - | 147 | 287 | 49 | 70 | 24 | NS | - | - | | 4 | - |
| .45 | - | - | 69 | - | 306 | 29 | | - | - | 3 | 4 | - |
| .50 | - | 270 | 502 | 1596 | 26 | 121 | 7 | - | - | 22 | 191 | - |
| .55 | - | 20 | - | - | 6 | 57 | 6 | - | - | 5 | 23 | - |
| .60 | - | 226 | | 48 | 131 | 54 | 5 | - | - | 3 | 34 | - |
| .65 | - | - | - | - | | 83 | 10 | - | - | - | - | - |
| .70 | - | | 12 | 23 | | 143 | 56 | - | - | | | - |
| .75 | - | - | - | - | | 24 | | - | - | - | - | - |
| .80 | - | - | - | 26 | | | NS | - | - | | | - |
| .85 | - | - | - | - | 14 | | | - | - | - | - | - |
| .90 | - | - | - | | 3 | 9 | | - | - | | - | - |
| 90.28 | - | 99 | 241 | 62 | 3 | 8 | 28 | - | - | | | 36 |
| .30 | - | 50 | 433 | 110 | 4 | 16 | | - | - | | 3 | 4 |
| .37 | - | | 72 | 136 | 13 | 13 | | - | - | 7 | | |
| .45 | - | 437 | 18 | 171 | 5 | 29 | 9 | - | - | | 9 | 22 |

Table VIII (cont'd)
Record of the larvae of rockfish (Sebastes spp.), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 90.50 | - | - | - | - | 177 | 7 | - | - | - | 56 | 14 | 212 |
| .55 | - | - | - | 298 | 1017 | 56 | 9 | - | - | 33 | 30 | - |
| .60 | - | 18 | 17 | 12 | - | 94 | 19 | - | - | 10 | - | - |
| .65 | - | - | - | - | 9 | 84 | 5 | - | - | - | - | - |
| .70 | - | - | - | 64 | - | 116 | 6 | - | - | - | - | - |
| .75 | - | - | - | - | - | 45 | 5 | - | - | - | - | - |
| .80 | - | - | 57 | - | - | 3 | 31 | - | - | - | - | 3 |
| .85 | - | - | - | - | - | - | 3 | - | - | - | - | - |
| .90 | - | - | - | 20 | 12 | 3 | - | - | - | - | - | 3 |
| 93.27 | - | 149 | 192 | 19 | 18 | - | - | - | - | - | 11 | - |
| .30 | - | 27 | 77 | 68 | 8 | - | 8 | - | - | - | - | - |
| .35 | - | - | - | 51 | 27 | 18 | - | - | - | 4 | - | - |
| .40 | - | - | 6 | 200 | 19 | 6 | 3 | - | - | 6 | 3 | - |
| .45 | - | - | - | 408 | 3 | - | - | - | - | - | - | - |
| .50 | - | 86 | 277 | 671 | 18 | 3 | 3 | - | - | - | 6 | - |
| .55 | - | - | - | 514 | 180 | 3 | 3 | - | - | - | - | - |
| .60 | - | 13 | 19 | 57 | 15 | 15 | - | - | - | - | - | - |
| .65 | - | - | - | - | 67 | 37 | - | - | - | - | - | - |
| .70 | - | - | - | - | 152 | 8 | - | - | - | - | - | - |
| .75 | - | - | - | - | - | - | - | - | - | - | - | - |
| .80 | - | - | - | NS | - | - | - | - | - | - | - | - |
| .85 | - | - | - | - | 3 | - | - | - | - | - | - | - |
| .90 | - | - | - | - | 12 | - | - | - | - | - | - | - |
| 97.30 | - | 61 | 486 | 135 | 103 | 6 | 8 | - | - | 4 | 13 | - |
| .32 | - | - | - | 57 | 22 | - | 2 | - | - | - | 39 | 85 |
| .35 | - | - | - | - | - | - | - | - | - | 4 | - | - |
| .40 | - | - | 90 | - | 4 | 3 | - | - | - | - | 5 | - |
| .45 | - | - | - | 61 | 3 | - | 8 | - | - | - | - | - |
| .50 | - | - | - | NS | 6 | - | 13 | - | - | - | - | - |
| .55 | - | - | - | NS | 143 | - | - | - | - | - | - | - |
| .60 | - | 26 | - | 13 | - | 9 | - | - | - | - | - | - |
| .65 | - | - | - | - | - | 4 | - | - | - | - | - | - |
| .70 | - | - | - | 8 | - | - | - | - | - | - | - | - |
| .75 | - | - | - | - | - | - | - | - | - | - | - | - |
| .80 | - | - | - | - | - | - | - | - | - | - | - | - |
| .85 | - | - | - | - | - | - | - | - | - | - | - | - |
| .90 | - | - | - | - | 8 | - | - | - | - | - | - | - |

Table VIII (cont'd)
Record of the larvae of rockfish (Sebastes spp.), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 100.29 | - | 271 | 35 | 52 | 9 | 12 | | - | - | | - | - |
| .30 | - | - | - | - | - | - | - | - | - | 6 | - | - |
| .33 | - | 90 | 3 | 84 | 26 | 6 | | - | - | - | - | - |
| .35 | - | - | - | - | - | - | - | - | - | - | - | - |
| .40 | - | 67 | | 44 | 89 | | | - | - | | - | 27 |
| .45 | - | - | - | 6 | 6 | | | - | - | - | - | - |
| .50 | - | | | | 12 | 12 | | - | - | | - | - |
| .55 | - | - | - | 17 | 14 | 7 | | - | - | - | - | - |
| .60 | | | | 101 | 33 | 3 | | - | - | | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | | | | | | - | - | - | | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | | | | | | | - | - | | - | - |
| .85 | - | - | - | - | | 6 | | - | - | - | - | - |
| .90 | - | | | | | | | - | - | | - | - |
| 103.30 | 103 | 190 | 38 | 83 | 14 | 17 | 2 | - | - | 8 | - | - |
| .35 | 61 | 111 | 9 | | 7 | | 3 | - | - | | - | - |
| .38 | | | - | - | - | - | - | - | - | - | - | - |
| .40 | | | | | 3 | | | - | - | | - | - |
| .45 | - | - | - | | 20 | | | - | - | - | - | - |
| .50 | | | | | 18 | 3 | | - | - | | - | - |
| .55 | - | - | - | 3 | | | | - | - | - | - | - |
| .60 | | | NS | | | | 3 | - | - | | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | | | | | | | - | - | | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | | - | - |
| .85 | - | - | - | - | | | | - | - | | - | - |
| .90 | - | - | - | - | | | | - | - | | - | - |
| 107.32 | 46 | 24* | 133 | 32 | 28 | - | | - | - | NQ | - | - |
| .35 | | | | 244 | 4 | 3 | | - | - | | - | - |
| .40 | 3 | | | 27 | 7 | | | - | - | | - | - |
| .45 | - | - | - | 3 | | | | - | - | - | - | - |
| .50 | | | | | | | | - | - | | - | - |
| .55 | - | - | - | | | | | - | - | - | - | - |
| .60 | | 6 | | | | | | - | - | | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | 9 | 47 | | | | | - | - | | - | - |

* - Samples combined. Values adjusted accordingly.

Table VIII (cont'd)
Record of the larvae of rockfish (Sebastes spp.), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 107.75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | - | - | - |
| .85 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | - | - | - | | | | - | - | - | - | - |
| 110.33 | - | 63 | 58 | | 8 | 6 | 14 | 11 | 2 | 9 | - | - |
| .35 | - | 12 | 432 | 6 | 12 | | | | | | - | - |
| .40 | | 3 | | 49 | | | | | | | - | - |
| .45 | - | - | - | 33 | 15 | | | - | - | - | - | - |
| .50 | | | | 3 | 9 | | 3 | - | - | | - | - |
| .55 | - | - | - | | 3 | 10 | | - | - | - | - | - |
| .60 | | | | 3 | | | | - | - | | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | - | | | 10 | | | | - | - | | - | 67 |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | | | | | | | - | - | | - | - |
| .85 | - | - | - | - | | | | - | - | - | - | - |
| .90 | - | | | | | | | - | - | - | - | - |
| 113.30 | 70 | 317 | 163 | 26 | 9 | 2 | 8 | | | | - | - |
| .35 | 8 | 70 | 228 | | | 3 | 8 | 3 | 3 | | - | - |
| .40 | | 8 | 14 | 9 | | | | | | | - | - |
| .45 | - | - | - | | | 3 | | - | - | - | - | - |
| .50 | | | 3 | | | | | - | - | | - | - |
| .55 | - | - | - | | | | | - | - | | - | - |
| .60 | | | | | | | | - | - | 3 | - | - |
| .65 | - | - | - | - | 5 | | | - | - | - | - | - |
| .70 | | | | | | | | - | - | | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | 3 | | | | - | - | | - | - |
| .85 | - | - | - | - | - | | | - | - | - | - | - |
| .90 | - | - | - | - | - | | | - | - | - | - | - |
| 115.27 | - | - | - | - | - | - | | | | | - | - |
| .30 | - | - | - | - | - | - | | | | | - | - |
| .35 | - | - | - | - | - | - | | | | | - | - |
| .40 | - | - | - | - | - | - | | | 3 | | - | - |
| 117.26 | 426 | 132 | | 8 | 7 | | | | 8 | 3 | - | - |
| .30 | 127 | 568 | 111 | 27 | 23 | | | | | | - | - |
| .35 | | 246 | 69 | 11 | 8 | 7 | | | | | - | - |
| .40 | 43 | 7 | 30 | 10 | 44 | | | 3 | | | - | - |

Table VIII (cont'd)
Record of the larvae of rockfish (Sebastes spp.), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 107.45 | - | - | - | | 32 | 1 | | - | - | - | - | - |
| .50 | 8 | 6 | 343 | | 8 | | | - | - | 3 | - | - |
| .55 | - | - | - | | 24 | | | - | - | - | - | - |
| .60 | | | 25 | | 3 | | | - | - | | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | | | | | | | | - | - | | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | | - | - |
| .85 | - | - | - | - | - | | | - | - | - | - | - |
| .90 | - | - | - | - | - | | | - | - | - | - | - |
| 118.39 | - | - | 76 | 11 | 27 | | | - | - | | - | - |
| 118.25 | - | - | - | - | - | - | | | | - | - | - |
| .30 | - | - | - | - | - | - | | 10 | | - | - | - |
| .35 | - | - | - | - | - | - | | | | - | - | - |
| 119.33 | - | - | 136 | | 5 | 11 | 10 | | | | - | - |
| 120.25 | | 30 | 6 | 2 | | 2 | | | | | - | - |
| .30 | 5 | 117 | 84 | | | | 13 | 7 | | | - | - |
| .35 | | 44 | | | | | 3 | | | | - | - |
| .40 | | 7 | | - | 2 | 4 | | | | | - | - |
| .45 | 6 | 6 | - | - | 10 | 3 | | | | | - | - |
| .50 | | | - | 23 | 13 | | | - | - | | - | - |
| .55 | | | - | 21 | | | | - | - | - | - | - |
| .60 | | | | | | | | - | - | | - | - |
| .65 | - | - | - | - | | | | - | - | - | - | - |
| .70 | | | | | | | 3 | | | | - | - |
| .75 | - | - | - | - | | | | - | - | - | - | - |
| .80 | - | - | - | - | | | | - | - | | - | - |
| .85 | - | - | - | - | - | | | - | - | - | - | - |
| .90 | - | - | - | - | - | | | - | - | | - | - |
| 123.37 | 12 | 32 | 418 | | 33 | 6 | 3 | | 3 | | - | - |
| .42 | 11 | | 48 | | 3 | | | 3 | | | - | - |
| .45 | - | - | - | - | - | - | | | | - | - | - |
| .50 | | | | 20 | - | | | - | - | | - | - |
| .55 | - | | | 8 | - | | | - | - | - | - | - |
| .60 | | 4 | | | - | | | - | - | | - | - |
| .70 | - | - | - | - | - | - | | - | - | | - | - |
| .80 | - | - | - | - | - | - | | - | - | | - | - |

Table VIII (cont'd)
 Record of the larvae of rockfish (Sebastes spp.), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 127.34 | 12 | 8 | 92 | 5 | 3 | | | | | 3 | - | - |
| .40 | 60 | | 8 | | | | 6 | 7 | | | - | - |
| .45 | - | | 22 | | | | | | | | - | - |
| .50 | | | 6 | | - | | | - | - | | - | - |
| .55 | - | | | | - | | | - | - | - | - | - |
| .60 | | | 3 | | - | | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | | - | - |
| 130.30 | | | 32 | 5 | 3 | | | | 5 | | - | - |
| .35 | | 65 | | | 24 | | | | | | - | - |
| .40 | 6 | 10 | | | | | | 3 | | | - | - |
| .45 | - | - | - | | - | | 3 | | | | - | - |
| .50 | | | | | | | | - | - | | - | - |
| .55 | - | - | - | | - | | | - | - | - | - | - |
| .60 | | | | | | | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | 4 | - | - |
| 133.25 | 3 | | 10 | | 9 | | 11 | | | | - | - |
| .30 | | 19 | 33 | | | | 20 | | 4 | | - | - |
| .35 | | - | | 170 | - | 10 | 3 | - | - | | - | - |
| .40 | | | | | | 5 | | - | - | | - | - |
| .45 | - | - | - | | - | | | - | - | - | - | - |
| .50 | - | | | | 6 | 3 | | - | - | | - | - |
| .55 | - | - | - | | - | - | - | - | - | - | - | - |
| .60 | - | - | - | | - | - | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | | - | - |
| 137.23 | | | | | | | 11 | | | | - | - |
| .30 | | | 6 | 13 | 3 | | | | | | - | - |
| .35 | | | 88 | - | - | | 4 | - | - | | - | - |
| .40 | - | | 20 | 10 | NS | | | - | - | | - | - |
| .45 | - | - | - | | - | | | - | - | - | - | - |
| .50 | - | | 3 | 3 | | | | - | - | | - | - |
| .55 | - | - | - | | - | - | - | - | - | - | - | - |
| .60 | - | - | - | | - | - | | - | - | | - | - |
| .70 | - | - | - | | - | - | | - | - | | - | - |
| .80 | - | - | - | | - | - | | - | - | | - | - |

Table VIII (cont'd)
Record of the larvae of rockfish (Sebastodes spp.), 1957

| Station | Cruise and month | | | | | | | | | | | |
|---------|------------------|--------------|--------------|--------------|-------------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|
| | 5701 Jan. | 5702 Feb. | 5703 Mar. | 5704 Apr. | 5705 May | 5706 June | 5707 July | 5708 Aug. | 5709 Sept. | 5710 Oct. | 5711 Nov. | 5712 Dec. |
| 140.30 | - | | | | - | | - | - | - | - | - | - |
| .35 | - | | 11 | | - | | - | - | - | - | - | - |
| .40 | - | | | 4 | | | - | - | - | - | - | - |
| .45 | - | - | - | | - | - | - | - | - | - | - | - |
| .50 | - | | | | - | | - | - | - | - | - | - |
| .55 | - | - | - | | - | - | - | - | - | - | - | - |
| .60 | - | - | - | | - | - | - | - | - | - | - | - |
| .70 | - | - | - | | - | - | - | - | - | - | - | - |
| .80 | - | - | - | | - | - | - | - | - | - | - | - |
| 143.26 | - | | | | - | - | - | - | - | - | - | - |
| .30 | - | | | | - | | - | - | - | - | - | - |
| .35 | - | | | 2 | | | - | - | - | - | - | - |
| .40 | - | | | | - | | - | - | - | - | - | - |
| .50 | - | | | | - | | - | - | - | - | - | - |
| 147.20 | - | | | | - | - | - | - | - | - | - | - |
| .25 | - | | | | - | - | - | - | - | - | - | - |
| .30 | - | | | | - | - | - | - | - | - | - | - |
| .35 | - | - | - | | - | - | - | - | - | - | - | - |
| .40 | - | | | | - | - | - | - | - | - | - | - |
| .45 | - | - | - | | - | - | - | - | - | - | - | - |
| .50 | - | - | - | | - | - | - | - | - | - | - | - |
| .55 | - | - | - | | - | - | - | - | - | - | - | - |
| .60 | - | - | - | | - | - | - | - | - | - | - | - |
| .70 | - | - | - | | - | - | - | - | - | - | - | - |
| .80 | - | - | - | | - | - | - | - | - | - | - | - |
| 148.20 | - | - | - | - | - | - | - | - | - | - | - | - |
| .25 | - | - | - | - | - | - | - | - | - | - | - | - |
| .30 | - | - | - | - | - | - | - | - | - | - | - | - |
| .40 | - | - | - | - | - | - | - | - | - | - | - | - |
| .50 | - | - | - | - | - | - | - | - | - | - | - | - |
| 150.16 | - | - | - | | - | - | - | - | - | - | - | - |
| .19 | - | | | | - | - | - | - | - | - | - | - |
| .20 | - | - | - | | - | - | - | - | - | - | - | - |
| .25 | - | | | | - | - | - | - | - | - | - | - |
| .30 | - | - | | | - | - | - | - | - | - | - | - |
| .35 | - | | | | - | - | - | - | - | - | - | - |
| .40 | - | | | | - | - | - | - | - | - | - | - |
| Total | 1010 | 11001 | 6585 | 7225 | 4819 | 2742 | 1383 | 47 | 28 | 231 | 586 | 816 |

Note: No rockfish found on lines 153 and 157.

LITERATURE CITED

AHLSTROM, ELBERT H.

1948. A record of the pilchard eggs and larvae collected during surveys made in 1939 to 1941. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. No. 54, 76 pp.
1952. Pilchard eggs and larvae and other fish larvae, Pacific coast, 1950. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. : Fisheries No. 80, 58 pp.
1953. Pilchard eggs and larvae and other fish larvae, Pacific coast, 1951. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. : Fisheries No. 102, 55 pp.
- 1954a. Pacific sardine (pilchard) eggs and larvae and other fish larvae, Pacific coast - 1952. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. : Fisheries No. 123, 76 pp.
- 1954b. Distribution and abundance of egg and larval populations of the Pacific sardine. U. S. Dept. Interior, Fish and Wildlife Service, Fish. Bull. 93, vol. 56, pp. 83-140.
1958. Sardine eggs and larvae and other fish larvae, Pacific coast, 1956. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. : Fisheries No. 251, 84 pp.

AHLSTROM, ELBERT H., and D. KRAMER

1955. Pacific sardine (pilchard) eggs and larvae and other fish larvae, Pacific coast, 1953. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. : Fisheries No. 155, 74 pp.
1956. Sardine eggs and larvae and other fish larvae, Pacific coast, 1954. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. : Fisheries No. 186, 79 pp.
1957. Sardine eggs and larvae and other fish larvae, Pacific coast, 1955. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. : Fisheries No. 224, 90 pp.

THRAILKILL, JAMES R.

- In Press. Zooplankton volumes off the Pacific coast, 1957. U. S. Dept. Interior, Fish and Wildlife Service, Spec. Sci. Rept. : Fisheries.

MBL WHOI Library - Serials



5 WHSE 01456

