

LIST OF PLATES

COD GILL-NETS.

	Page.
PLATE I.—Method of hanging cod gill-nets in Norway	16
PLATE II.—Manner in which the glass floats are attached to the tops of the nets	16
PLATE III.—Way in which the sinkers are fastened to the bottom of the nets	16
PLATE IV.—Size of twine of which the American nets are made	16
PLATE V.—Norwegian method of setting the nets at the bottom	16
PLATE VI.—Way in which nets are set at different depths to ascertain the position of the fish ..	16
PLATE VII.—Manner in which the ends of a gang of nets are attached to the stone anchors and buoy-line in Norway; also showing the position of the glass floats and sinkers	16
PLATE VIII.—Way in which cod gill-nets are set at the bottom on the east coast of Newfoundland.	16
PLATE IX.—The ordinary way in which cod gill-nets are set floating at Newfoundland	16
PLATE X.—Norwegian net and trawl buoy made of glass floats	16
PLATE XI.—Way in which cod gill-nets are set for underrunning in Ipswich Bay	16
PLATE XII.—Manner in which the nets are underrun	16

SPANISH MACKEREL.

PLATE XIII.—(I.) Fig.	1. Micropyle and micropylar area of egg	166
	2. Unimpregnated egg	166
	3. Morula stage of cleavage of the germinal disk	166
	4. Germinal disk of egg	166
	5. Incipient blastoderm of the egg	166
	6. Developing blastoderm of the egg	166
	7. Blastoderm of the egg 7 hours after impregnation	166
	8. Embryo 11 hours after impregnation	166
PLATE XIV.—(II.) Fig.	9. Caudal view of embryo	168
	10. Embryo 14 hours after impregnation	168
	11. Transverse section of caudal view of embryo	168
	12. Embryo 18 hours after impregnation	168
	13. Young Spanish mackerel 24 hours from impregnation	168
PLATE XV.—(III.) Fig.	14. Young Spanish mackerel 36 hours from impregnation	170
	15. Young Spanish mackerel 45 hours from impregnation	170
PLATE XVI.—(IV.) Fig.	16. Spanish mackerel three days after it left the egg	172
	17. Head of Spanish mackerel six days after hatching	172

SEA-HORSE.

PLATE XVII. Fig. 1. Young sea-horse (<i>Hippocampus antiquorum</i>)	199
---	-----

SILVER GAR.

PLATE XVIII. Fig. 1.	Egg of the silver gar (<i>Tylosurus longirostris</i>)	300
	Germinal disk 4½ hours after impregnation	300
	Germinal disk 10 hours after impregnation	300
	Blastoderm of silver gar 24 hours after impregnation	300
	Blastoderm of silver gar 31½ hours after impregnation	300
	Blastoderm of silver gar 43½ hours after impregnation	300
	Head of embryo of silver gar 51 hours after impregnation	300
	Tail of embryo of silver gar 51 hours after impregnation	300
PLATE XIX. Fig.	9. Embryo silver gar 51 hours after impregnation	300
	10. Embryo silver gar 52 hours after impregnation	300
	11. Embryo silver gar 70 hours after impregnation	300
	12. Embryo silver gar 94½ hours after impregnation	300
	13. Origin of the mode of anastomosis of larger vessels	300
PLATE XX. Fig.	14. Embryo silver gar 116½ hours after impregnation	300
	15. Heart of silver gar 140 hours after impregnation	300
	16. Embryo silver gar 165½ hours after impregnation	300

SALMON.

PLATE XXI. Fig. 1-23. Nuclear cleavage figures of salmon eggs	338
---	-----

ERRATA.

[Fish Com. edition U. S. F. C. Bull., 1881.]

- On page 166, and on the plate facing it, for Plate I read Plate XIII.
- On page 167, and on the plate facing it, for Plate II read Plate XIV.
- On page 168, and on the plate facing it, for Plate III read Plate XV.
- On page 169, and on the plate facing it, for Plate IV read Plate XVI.
- On page 300, and on the plate facing it, for Plate XIX read Plate XVIII.
- On page 300, and on the plate facing it, for Plate XX read Plate XIX.
- On page 301, and on the plate facing it, for Plate XXI read Plate XX.
- On page 338, and on the plate facing it, for Plate XVIII read Plate XXI.