LIST OF ILLUSTRATIONS.

Plate		Dago
1 /124	in 1) The Belfast Tokston, down of Male Tokston, living mainly a little over 99 nounde	- ago.
1. (E) 9 (TN	\mathbf{R}_{i}) The Denast Loopster; (orsai view of mais Looster; firing weight a fitting over 25 pounds	202
2. (FI	with Fig. 1 Ergs howing Fermala	959
3 / FY	which Decirity and Bernard Conductor Framala	252
J (F)	ig. =) i tomo valv o inving iou i iousovi, i omnovi i o invingi i o i a do invingi i o i a do i i o i o i o i o	959
5 (F)	15. 0) Matu Mato Dobta, and a constant shown in Plate 4	252
6 (TF)	2, of Idult Rample Lobstan with external Ears. Aarsal view	252
7 (17)	16. 7) Hault Fondel, ventrel view of Lobster shown in Plete 6	252
8. (Fi	12, 9) Inumature Kennala Labatar, dorsal view. (Fig. 10) Immature Male Lobatar. (Fig. 11) Immature Fennal	202
~,	Lobster. (Fig. 12) Immature Male Lobster.	252
9. (Fi	ig. 13) Immature Female Lobster. (Fig. 14) Immature Male Lobster	252
10. (F	12, 15) Immature Female Lobster: dorsal view	252
11. (Fi	ig. 16) Ventral view of immature Female Lobster shown in Plate 10	252
12. (Fi	ig. 17) Immature Male Lobster	252
13. (Fi	ig. 18) Immature Male Lobster	252
14. (Fi	ig. 19) Male Lobster, showing abnormal symmetrical development in large Claws	252
15. (Fi	ig. 20) Right crushing-claw of Lobster, probably a Male; estimated weight about 25 pounds. (Fig. 20a) Right	i
	. crushing-claw of Female Lobster of about average size	252
16. (Fi	ig. 21) Red Female Lobster, colored from life. (Fig. 22) Adult Male Lobster, colored from life	252
17. (Fi	ig. 23) Eggs of Lobster, showing an unusual color variation. (Fig. 24) Cluster of fresh Eggs of Lobster, colored	
	from life. (Fig. 25) Cluster of Egg Embryos from swimmerets of Female, shown in Plate 7. (Fig. 26)	
	Profile view of Embryo released from eggshell. (Fig. 27) Side view of Embryo, as seen through the	
	transparent Shell. (Fig. 28) Dorsal view of Embryo, shown in Fig. 27	252
18. (Fi	ig. 29) Lobster Hatching. (Fig. 30) Lateral view of a Lobster teased from an Egg about to Hatch. (Fig. 31)	
	Profile view of Lobster in Fifth Stage	252
19. (Fi	ig. 32) First swimming stage of the Lobster	252
20. (Fi	ig. 38) Profile view of the first Larva of the Lobster	252
21. (Fi	ig. 34) Second Larva; profile view	252
22. (Fi	ig. 35) 'Third Larva; lateral view	252
23. (Fi	ig. 36) Fourth Larva; dorsal view	252
24. (Fi	ig. 37) Sixth stage; dorsal view	252
20. (F1	1g. 38) Young Lobster in sixth stage; profile view	252
20. (F)	1g. 39) Young, immature Lobster; Male.	252
27. (F1	19. 40) Right first antenna of first Larva, from below. (Fig. 41) Left first antenna of second Larva, from below	
	(Fig. 42) Lott first antenna of Larva, irom below. (Fig. 43) Left first autonna of fourth Larva, from	
	above. (Fig. 44) Leit first antenna of nith stage, from below. (Fig. 45) Kight first and second antenna	
	of first Larva, from above. (Fig. 40) Right second antenna of second Larva, from above. (Fig. 47) Left	
	second antenna of third Larva, from above. (Fig. 48) Left second antenna of fourth Larva, from below.	010
28 (17)	(Fig. 49) Frozman portion of left first and second antennae of Lobster in inth stage, seen from below	202
-0. (1.1	ig. of lest instand second antennas of infut Larva, as seen from above. (Fig. of) Angle hist inagina of infut	
	Larva, from anterior face. (Fig. 52) reminal joint of 1611 first perception of first Larva, from anterior side (Fig. 53) The of endoudite of face maxilla of fare Larva (Fig. 54) Front view of month and spr	
	rounding norts of first Lows (Fig 55) Right mandials of Auth Laws from babind showing groups	
	and outfing plates of first Sal Laftmandilla of fourth Larva from outar side (First S7) Mondibles of fourth	
	Lava, from antarior side.	252
29, (Fi	iz, 58) Left first maxillined of first Larva, from the inner side. (Fig. 59) Left first maxillined of fourth Larva,	
,- ·	from outer side. (Fig. 60) Right second maxilla of first Larva, from outer side. (Fig. 61) Right first	
	maxilla of fourth Larva, from inner side. (Fig. 62) Right first maxilla of fifth Larva, from outer side	252
30. (Fi	ig. 63) Left second maxilliped of first Larva, from anterior face. (Fig. 64) Left second maxilliped of fourth	
	Larva, from anterior face. (Fig. 65) Right third maxilliped of fourth Larva, from dorsal surface, natural	

VII

Plate	Page
 position. (Fig. 66) Left first pereiopod of fourth Larva, from below. (Fig. 67) Left first pereiopod of fourth Larva, from below. (Fig. 68) Part of left third maxilliped of fourth Larva, from below, showing serrated inner margin of third segment. (Fig. 69) Left third maxilliped of first Larva, from above 31. (Fig. 70) Left fourth pereiopod of first Larva, from above. (Fig. 71) Serrated spine from propodus of left second pereiopod of fourth Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 74) Left second pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Larva, from the side. (Fig. 76) Left fourth pereiopod of first Lar	1 ago. f 2 252 1 1 1
 above. (Fig. 77) Antennæ of Embryo 32. (Fig. 78) Bud of first left abdominal appendage of fifth Larva. (Fig. 79) Seminal receptacle of Female. (Fig. 80) Left first abdominal appendage of Lobster. (Fig. 81) Ventral view of young Female Lobster. (Fig. 82) Left first abdominal appendage of the sixth stage of development. (Fig. 83) Left first abdominal appendage of the sixth stage of development. (Fig. 83) Left first abdominal appendage of Female in eighth stage. (Fig. 85) Left first abdominal appendage of Female in eighth stage. (Fig. 86) Left first abdominal appendage of Female in eighth stage. (Fig. 86) Left first abdominal appendage of Female. (Fig. 87) Left first abdominal appendage of Male. (Fig. 88) Left first abdominal appendage of Female. (Fig. 80) Ventral view of Female Lobster in eighth stage. (Fig. 86) Left first abdominal appendage of Female. (Fig. 80) Ventral view of Female Lobster in eighth stage. (Fig. 81) Left first abdominal appendage of Male. (Fig. 83) Left first abdominal appendage of Female. (Fig. 80) Ventral view of Female Lobster in eighth stage. (Fig. 81) Left first abdominal appendage of Male. (Fig. 80) Ventral view of Female Lobster in eighth stage. (Fig. 80) Left first abdominal appendage of Female. (Fig. 80) Ventral view of Female Lobster in eighth stage. (Fig. 80) Ventral view of Female Lobster in eighth stage. 	. 252) f f
 age of young Male. (Fig. 9) Vontral view of young Male. 33. (Fig. 92) Left cheliped of fourth Larva in process of regeneration from stump. (Fig. 93) Left fourth pleopod of second Larva, from outer face. (Fig. 94) Left second pleopod of third Larva, from outer face. (Fig. 95) First abdominal segment of shell of Lobster in sixth stage, seen from behind. (Fig. 96) Left cheliped of molted shell of fifth Larva, seen from above. (Fig. 97) Left second pleopod of fourth Larva, from anterior molted shell of fifth Larva, the larva is the stage. (Fig. 97) Left second pleopod of fourth Larva, from anterior molted shell of fifth Larva, been from above. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth Larva, from anterior form. (Fig. 97) Left second pleopod of fourth second pleopod fourth seco	. 252 f) f
 14ce. (Fig. 98) Sterna of the last three thoracic somities of fifth Larva, (Fig. 99) Left fourth perception of fourth Larva. (Fig. 100) Right second antenna of Lobster in seventh stage, seen from above	f 252 1 252
35. (Fig. 107) Left first antenna of the Embryo, frontal view. (Fig. 108) Right second antenna of same Embryo, from below. (Fig. 109) Rostrum of second Larva, from above. (Fig. 110) Profile view of carapace of first Larva. (Fig. 111) Profile view of carapace of second Larva. (Fig. 112) Profile view of carapace of third Larva. (Fig. 113) Profile view of carapace of fourth Larva. (Fig. 114) Profile view of carapace of fifth Larva. (Fig. 113) Profile view of carapace of fight below.	050
 86. (Fig. 116) Section of reproductive organ of Embryo near time of hatching. (Fig. 117) Section of right reproductive organ of first Larva. (Fig. 118) Right second antenna of an adult fomale Lobster, overgrown with algre. (Fig. 119) Oviduct and part of ovarian lobe from left side, showing a row of unextruded Eggs in duct. (Fig. 120) Reproductive organs of adult Male Lobster from above. (Fig. 121) Disk-shaped concretion, probably containing glycogen, from maxilla. (Fig. 122) Large granular cell from first maxilliped. (Fig. 	
 123) Reproductive organs of adult Female dissected out, viewed from above	252
 section of vas deforens through glandular portion. (Fig. 129) Ripe sperm cells	252
recently hatched a brood	252
 gland-like organs. 40. (Fig. 144) Right pleopod of adult Female, seen from posterior surface. (Fig. 145) Fold of glandular epithelium from transverse section of ovary of Lobster. (Fig. 146) Transverse section of lobe of ovary. (Fig. 147) Part of transverse section of ovarian lobe from a Lobster with external eggs near the point of hatching. 	252
 (Fig. 148) Part of transverse section of ovary. (Fig. 149) Part of transverse section of ovary	252
 Glandular epithelium from transverse section of ovary of Lobster	252
bifurcation in left anterior lobe. (Fig. 165) Part of gastrolith, separated into its constituent spicules	252

ILLUSTRATIONS.

Plate No.	age.
 43. (Fig. 166) Bud of right fourth pereiopod in process of regeneration from young Lobster. (Fig. 167) Part of transverse section of oviduct of Lobster, with ovary nearly ripe. (Fig. 168) Part of transverse section of oviduct of Lobster, with external Eggs in early yolk segmentation. (Fig. 160) Longitudinal section of first, second, and third segments of right first pereiopod of young Lobster in sixth stage. (Fig. 170) Internal surface of cuticle of second joint of first maxilla. (Fig. 171) Part of section of gastrolithic plate from female Lobster with hard shell. (Fig. 172) Section of left first pereiopod of Lobster 9 inches long, in process of regeneration. (Fig. 173) Section of right first (crushing) cheliped of adult male Lobster. (Fig. 174) Part of longitudinal section of first Larva. 44. (Fig. 175) Right fourth pereiopod of adult Lobster in process of regeneration, from below. (Fig. 176) Stump of right 	252
first pereiopod of adult Lobster in course of regeneration, from below. (Fig. 177) Surface view of mem- brane between old and new shells of molting Lobster. (Fig. 178) Right second pereiopod of adult Male in process of regeneration, from below. (Fig. 179) Left second antenna of adult Lobster in process of regener- ation, from above. (Fig. 180) Antennæ of the Isopod, <i>Ligea oceanica</i> , from above; that of the left side in course of regeneration. (Fig. 181) Regenerating left antenna of the same, showing the new flagellum inclosed in the exoskeleton of the joint. (Fig. 182) Left first cheliped of adult Lobster in process of regen- eration, seen from the inner and anterior side. (Fig. 183) Profile view of masticatory stomach of Lobster, showing gastrolithic plate. (Fig. 184) Profile view of masticatory stomach of Male Lobster 7.5 inchess long. nearly ready to molt	252
45a. (Fig. 185) Molted shell of Lobster shown in Fig. 186	252 252
 (Fig. 189) Part of right crushing-chela of Female Lobster, showing budding of dactyl. (Fig. 190) Propodus of left crushing-claw, from below. (Fig. 191) Left crushing-claw, seen from above; outgrowth from dactyl in horizontal plane. (Fig. 192) Left crushing-chela, from above; secondary dactyl bent downward slightly. (Fig. 193) Right cutting-chela, from below; fingers bent up; dactyls articulate at joint with propodus. (Fig. 194) Dactyl of left cutting-chew, seen from below. (Fig. 195) Chela of second and third percisiond. 	252
from below, showing two supernumerary dactyls. (Fig. 196) Right dactyl of cutting-chela, seen from outer	959
48. (Fig. 197) Deformed right cutting-claw. (Fig. 198) Right cutting-claw. (Fig. 199) Double monster of first	959
49. (Fig. 201) Gland-cell from togumental gland of second maxilla. (Fig. 202) Gland-cell from same preparation as Fig. 201. (Fig. 203) Part of macerated tegumental gland from metastoma. (Fig. 204) Cell from macerated tegumental gland of first maxilla. (Fig. 205) Gland-cell from same preparation as Fig. 204. (Fig. 204) Cell from macerated tegumental gland of abdominal appendage of Female before egg extrusion. (Fig. 207) Same preparation as ast, rolled under cover slip and seen from opposite side. (Fig. 208) Tegumental gland from metastoma of Female with ripe ovaries. (Fig. 209) Gland-cell from tegumental gland of abdominal appendage of Female algorithm to abdominal appendage of Female Lobster preparing to molt. (Fig. 211) Tegumental gland from abdominal appendage of Female Lobster with mature ovaries. (Fig. 213) Gland-cells from abdominal appendage of Female labster with mature ovaries. (Fig. 213) Gland-cells from tegumental gland from abdominal appendage of Female Lobster preparing to molt. (Fig. 211) Tegumental gland from abdominal appendage of Female after ovulation. (Fig. 213) Gland-cells from same preparation as Figs. 204, 205. (Fig. 214) Macerated tegumental gland	
 from metastoma of Female	252
 Surface view of Egg in abnormal yolk segmentation (Fig. 227) Surface view of Egg in invagination stage. (Fig. 228) Surface view of abnormal embryo in egg- nauplius stage. (Fig. 220) Surface view of abnormal Embryo. (Fig. 230) Surface view of abnormal Embryo in egg-nauplius stage. (Fig. 231) Surface view of abnormal Embryo in egg-nauplius stage. (Fig. 232) Lateral view of Embryo, showing large white patch behind abdomen. (Fig. 233) Lateral view of Embryo about 5 weeks old, showing lateral fold of carapace covering the antenne. (Fig. 234) Surface 	252
 view of Embryo about 25 days old, showing the large optic lobes of cephalo-thoracic appendages. (Fig. 235) Lateral view of double monster in Egg-nauplius stage	252
Egg in late segmentation of yolk. (Fig. 245) Part of transverse section, showing multiple karyokinesis and formation of nests of nuclei. (Fig. 246) Part of transverse section through Embryo in invagination stage. (Fig. 247) Part of section of Egg to show nest of nuclei at surface. (Fig. 248) From section through Embryo in invagination stage. (Fig. 249) Fertilized Egg nucleus	252

ILLUSTRATIONS.	

Plate	
No.	Page.
 (Fig. 250) Surface view of Embryo in invagination stage. (Fig. 251) Part of transverse section through area of invagination. (Fig. 252) Surface view of Embryo in region of invaginate area. (Fig. 253) Part of longitudinal section of intestine of Embryo in a late stage of development. 	252
opment. (Fig. 254) Fart of transverse section through invaginate area of an earlier stage than last.	
(Fig. 255) Part of longitudinal section through area of invagination. (Fig. 256) Concretion from intes-	
tine of an Embryo which was nearly ready to hatch	252
Cut 1. Curve of fecundity of the Lobster	52 52
Cut 3. Curve showing the relative feeundity of 352 Lobsters, each 10 inches long	54
Cut 4 (Plate A). Part of transverse section of exopodite of pleopod of Female Lobster (the cuticle removed),	• •
showing the distribution of the cement glands Cut 5 (Plate A). Diagram of vertical section through skin, showing a tegumental gland in sections, and its	77
duct opening to the exterior	77
Cut 6 (Plate B). Left cheliped of Lobster, seen from the dorsal side	86 86
Cut 8 (Plate C). The gastrolith of a Lobster nearly ready to molt, removed from the wall of the stomach	89
Cut 9 (Plate C). Diagrammatic section through the wall of the stomach of a molting Lobster, cutting gastrolith	89
Cut 10 (Plate C). Section of the decidnous portion of old cuticular lining of stomach overlying gastrolith	89
Cut 11 (Plate C). Section of gastrolithic sac from wall of stomach underlying gastrolith, as it appears while the latter is still in place in the stomach	89
Cut 12 (Plate D). First left perciopod of adult Lobster, seen from in front, showing anterior border at base of limb.	100
Cut 13 (Plate D). Basal portion of first left perciopod of adult Lobster, from under side	100
Cut 14 (Plate D). Second left pereiopod of Female, 10 inches long, seen from under side	100
Cut 15. Part of first cheliped of fourth Larva, showing the base of the limb and distinct articulation between	100
the second and third joints.	102
Cut 17 (Plate E). Double right cutting-claw of the same Lobster, seen from above	147
Cut 18. From transverse section of a part of ovary of Lobster, hardened with ventral side uppermost, to show the effect of gravity upon the nucleolus	154
Cut 19. From transverse section of a part of same ovary, hardened with dorsal side uppermost, to show effect of gravity upon the nucleolus	154
Cnt 20 (Plate F). Egg Embryo, showing membranes abnormally distended after prolonged immersion in picro- nitric acid.	205
Cut 21 (Plate F). Projection of an Egg with 15 yolk-cells, all near the surface or approaching it	205
Cut 22 (Plate F). Projection of an Egg with 28 yolk cells, 3 in karyokinesis, mostly near the surface	205
Cut 23 (Plate G). Egg with about 32 yolk segments present, seen from vegetative pole.	206
Cut 25 (Plate G). Surface view of Embryo 8 days old in invagination stage, showing pit at surface, embryonic	200
area, and mass of in-wandering cells which penetrate deeply into the yolk	206
Cut 26 (Plate G). Surface view of Egg in invagination stage, pit transversely elongated, showing tendency to become horseshoe shaped	206
Cut 27 (Plate H). Surface view of Embryo, showing buds of first pair of antennæ and clouds of in-wandering cells	208
Cut 28 (Plate H). Surface view of Embryo, showing buds of first pair of antennæ and of mandibles	208
Cut 29 (Plate H). Surface view of early Egg-nauplius Embryo, showing buds of the first and second antennæ and the mandibles	208
Cut 30 (Plate H). Surface view of Egg-nauplius, second antennæ bifid; labrum and thoracic abdominal fold present	208
Cut 31 (Plate I). Surface view of Egg-nauplius, showing thoracic abdominal fold	209
Cut 32 (Plate I), Surface view of Egg-nauplius, showing parts much more concentrated than in earlier stages.	209
Cut 33 (Flate 1). Surface view of Embryo who into into intaning budged.	209 209
Cut 35 (Plate J). Surface view of Embryo with eye plgment in form of crescent, as seen from the surface	210
Cut 36 (Plate J). Embryo 61 days old; area of eye-pigment semicircular; telson behind brain	210
Cut 37 (Plate J). Embryo 122 days old; area of eye-pigment rounded or irregularly oval in outline	210
Out os (riate J). Empryo 211 days old; area or eye-pigment irregular, somewhat oval or rounded in outline	210 215
Cut 40. Sagittal section through abnormal Embryo	215
55. Fish-cultural Apparatus used at Freising	320
56. Chart of Mobile Bay, Alabama, showing location of Natural Oyster Beds	825
57. Shells of Specimens of cultivated Uysters from Plant Beds at mouth of Fowl River, Alabama	340
59. Bunch of Ovsters from Great Point Clear Reef, showing growth of Mussels and Barnacles	340
60. Shells of Oysters from Fish River Reef	340

x

ILLUSTRATIONS.

No.		Pag
61. S	pecimen Shells of young Oysters from Bon Secours Reef.	3
62. O	ysters from vicinity of Little Dauphin Island	3
63. S	hells of Oysters from Little Dauphin Bay.	3
64. M	lap of Buzzards Bay, showing positions of Observation Stations relative to Sources of Marine Food	3
65. O	rganisms found abundantly in common sea water that has stood a few days in an open, shallow dish	3
66. O	rganisms common in the Plankton of Buzzards Bay	3
67. O	rganisms common in Plankton of Buzzards Bay, continued.	3
68. P	latting of Organisms of the longitudinal section designated "Letter M"	3
69. P	latting of Organisms as obtained from longitudinal section called "Letter K"	3
70. P	latting of the same group of Organisms as obtained from the cross section "Letter N"	3
71. P	latting of the same groups as collected from cross section "Letter H" (low water)	3
72. F	igs, 1 to 8, outlining plan for culture of Salmonoids and Crawfish in the smaller water-courses	3
73. P	ike Perch or Wall-eyed Pike (Stizostedion vitreum). Pike or Pickerel (Lucius lucius)	3
74. W	Vhite Catfish or Schurlkill Catfish (Ameiurus catus). Yellow Catfish or Bullhead (Ameiurus nebulosus). Spot-	
	ted Cattish (Ictaturus punctatus).	3
75. A	siatic Carp; Scale Carp (Oyprinus carpio). German Carp; Leather Carp (Oyprinus carpio coriaceus). Tench	
-	(Irnea linea).	33
76. S	had (Uupea sapidissina).	4
77. V	V hitchish (Coregonus clupeiformis). A tiantic saimon (saino sain). Eastern brook Irolt (saiveinus fontmains).	4
78. V	on Behr Trout or European Brown Frout (Salmo Jario). Loch Leven Frout (Salmo tratta tevenensis). Lake Frout	
70 T	or Mackinaw Trout (Satzetinus namajeash).	4
79. E	(Anguina chrystyph). Grappy; Strawberry Bass; Canco Bass (Fomous sparoaes). Grappy; Sac-a-iai; Bach-	
00 1	610T [Pomozis annuaris]. Warmouth Deca (Chemohenting colored)	4
80. IS	Sock Bass (Amotopites rupeerru). Warmouth bass (Omenorgens guissis)	4
81. S	mail-mouth Black Dass (<i>micropierus acionieu</i>). Large-Botth Black Bass (<i>micropierus sainoides</i>). Yellow Perch or Ringed Perch (<i>Perca flavescens</i>)	4
82. S	triped Bass (Roccus lineatus)	4
83. V	Vhite Bass (Roccus chrysons). Tautog (Tautoga onitis)	4

хı