

SOME STATISTICAL TECHNIQUES FOR ESTIMATING ABUNDANCE INDICES FROM TRAWL SURVEYS

MICHAEL PENNINGTON¹

ABSTRACT

Methods are presented for estimating an index of relative abundance from trawl survey catch per tow data. The estimated variance of the index takes into account the within survey variability in catch and possible yearly changes in catchability. Applying the techniques to a series of surveys for yellowtail flounder, *Limanda ferruginea*, off the northeast coast of the United States yields an abundance index with a variance which is 40% lower than the variance of the original survey index for the current value and 57% lower for values not near the ends of the survey series.

The average number of fish caught per tow during a trawl survey is often used as an index of a species's relative abundance (Grosslein 1969; Clark 1979). Catch per tow data are usually quite variable because of the heterogeneous distribution of many fish stocks (Byrne et al. 1981). A further source of variability for survey indices of abundance is that the catchability of a particular species with respect to the survey trawl may change from year to year (Byrne et al. 1981; Collie and Sissenwine 1983). As a result, the observed time series of abundance indices reflects changes in the population, within survey sampling variability, and varying catchability over time.

This paper uses various statistical methods to construct from the catch per tow data an index of abundance which more closely tracks the population than does the original (average catch per tow) series. Specifically, since the distribution of catch per tow data is often highly skewed and contains a proportion of zeros, estimates of the mean catch per tow for each survey are made based on the Δ -distribution (Aitchison and Brown 1957). Next, time series techniques are used to estimate the component of the series generated by the actual changes in the population.

The methods are applied to data for yellowtail flounder, *Limanda ferruginea*, from a series of groundfish trawl surveys conducted off the northeast coast of the United States as part of the National Marine Fisheries Service's MARMAP program. The resulting index of abundance is substantially more precise than the original index.

STATISTICAL METHODS

Sources of Variability

Let y_t denote the observed average catch per tow for the survey conducted in year t and $z'_t = E[y_t]$, the expected value of y_t . Since a species catchability may change from year to year with respect to the survey trawl, let $z = E[z'|p]$ denote the expected value of z' given a population level p . Then

$$y_t = z_t + e_t.$$

The error term, e_t , can be expressed as

$$e_t = (y_t - z'_t) + (z'_t - z_t),$$

where the first error component is due to the within survey variability and the second is due to changes in catchability.

In order to construct an index of abundance, it is necessary to assume a functional relationship between z_t and p_t . A reasonable assumption made in practice (and in this paper) is that

$$z_t = ap_t.$$

If the relationship is not linear, then the unadjusted catch per tow index will be a biased measure of relative abundance.

Estimating the Mean Catch per Tow

The distribution of marine survey data often can be described by what is called a Δ -distribution (Aitchison and Brown 1957). That is, the data contain

¹Northeast Fisheries Center Woods Hole Laboratory, National Marine Fisheries Service, NOAA, Woods Hole, MA 02543.

a proportion of zeros and the nonzero values are distributed lognormally. The minimum variance unbiased estimates of the mean, c , and its variance, $\text{var}(c)$, for the Δ -distribution are given by (Pennington 1983),

$$c = \begin{cases} \frac{m}{n} \exp(\bar{y}) G_m(s^2/2), & m > 1, \\ \frac{x_1}{n}, & m = 1, \\ 0, & m = 0, \end{cases} \quad (1)$$

and

$$\text{var}(c) = \begin{cases} \frac{m}{n} \exp(2\bar{y}) \left[\frac{m}{n} G_m^2(s^2/2) - \left(\frac{m-1}{n-1} \right) \times G_m \left(\frac{m-2}{m-1} s^2 \right) \right], & m > 1, \\ \left(\frac{x_1}{n} \right)^2, & m = 1, \\ 0, & m = 0, \end{cases} \quad (2)$$

where n is the number of tows, m is the number of nonzero values, \bar{y} and s^2 are the sample mean and variance respectively of the nonzero \log_e values, x_1 is the single (untransformed) nonzero value when $m = 1$, and

$$G_m(x) = 1 + \frac{m-1}{m} x + \sum_{j=2}^{\infty} \frac{(m-1)^{2j-1} x^j}{m^j (m+1)(m+3)\dots(m+2j-3)j!}$$

The series defining $G_m(x)$ is a function of x [e.g., $x = s^2/2$ in Equation (1)] and m which is easily evaluated for particular values of x and m using a computer.

Figure 1, which is an extension of a graph in Aitchison and Brown (1957, p. 98), shows the large sample efficiency of the ordinary sample statistics as compared with their most efficient estimates for the Δ -distribution with 50% zeros. Estimates of σ^2 , the variance of the nonzero \log_e values, are often between 1 and 2 for trawl surveys. Thus (Fig. 1) the




FIGURE 1.—The efficiency of \bar{x} and s^2 (the sample mean and variance, respectively) for the Δ -distribution with 50% zeros.

sample mean is a fairly efficient estimator of the mean for trawl surveys, but the sample variance is highly inefficient. Though for larger values of σ^2 , which, for example, are common for egg surveys (Pennington and Berrien 1984), the sample mean is also very inefficient. It does not follow that the variance of c is necessarily small, but it is smaller, and as σ^2 increases, much smaller than the variance of the sample mean. However, it should be noted that if the sample variance is used to estimate the variance of the sample mean for moderate sample sizes because of the inefficiency of the sample variance, the estimated variance of c will often be greater than the estimated variance of the sample mean.

Estimating the Index of Abundance

As an index of abundance, the series of yearly catch per tow estimates, y_t , (based, e.g., on the Δ -distribution theory if appropriate) has two drawbacks. First, its estimated variance when derived from the within survey variance can be an underestimate since catchability may vary from year to year. The second and more serious deficiency is that the index for a particular year is based only on that year's survey which disregards relevant information contained in the surveys for other years.

One method to construct an abundance index based on the entire survey series is briefly as follows (more details can be found in Pennington (1985)).

Suppose the population (or z_t) can be represented by the autoregressive integrated moving average process (Box and Jenkins 1976, Chap. 4)

$$\Phi(B) z_t = \theta(B) a_t.$$

where the a_t 's are independently identically distributed (*iid*) and normally distributed (N) with mean zero and variance σ_a^2 [*iid* $N(0, \sigma_a^2)$]. If $y_t = z_t + e_t$, and the e_t 's are assumed *iid* $N(0, \sigma_e^2)$, then y_t will follow the model

$$\Phi(B) y_t = \eta(B) c_t, \quad (3)$$

where the c_t 's are *iid* $N(0, \sigma_c^2)$. Now if model (3) and the ratio σ_e^2/σ_c^2 are known, then the maximum likelihood estimate of z_t is given by

$$\hat{z}_t = y_t - \frac{\sigma_e^2}{\sigma_c^2} (\hat{e}_t - \pi_1 \hat{e}_{t+1} - \pi_2 \hat{e}_{t+2} - \dots - \pi_{T-t} \hat{e}_T), \quad (4)$$

where T denotes the last year of the series, the \hat{e}_t 's are the estimated residuals generated by model (3), and the π values are calculated using the identity

$$\Phi(B) = (1 - \pi_1 B - \pi_2 B^2 - \dots) \eta(B). \quad (5)$$

The variance of \hat{z}_t is given approximately by

$$\text{var}(\hat{z}_t) \doteq \sigma_e^2 \left[1 - (\pi_0^2 + \pi_1^2 + \dots + \pi_{T-t}^2) \frac{\sigma_e^2}{\sigma_c^2} \right], \quad (6)$$

where $\pi_0 = 1$.

The model for y_t [Equation (3)] is usually obtained in practice by fitting a model to the observed series using procedures described in Box and Jenkins (1976). If catchability is constant over time, the within survey sampling variance provides an estimate of σ_e^2 . But if catchability varies, another approach is necessary.

Toward this end, consider the expression

$$z_t = z_{t-1} e^{a_t}$$

or

$$(1 - B) \ln z_t = a_t. \quad (7)$$

Suppose the factors causing the change in population from year $t - 1$ to year t (such as recruitment, fishing mortality, natural mortality, and migrations) produce a_t 's which are approximately *iid* $N(0, \sigma_a^2)$. If the measurement errors are multiplicative, then

$$\ln y_t = \ln z_t + e_t. \quad (8)$$

Assuming the e_t 's are *iid* $N(0, \sigma_e^2)$ and independent of the a_t 's, then it follows as above that y_t can be represented by the model

$$(1 - B) \ln y_t = (1 - \theta B) c_t. \quad (9)$$

where the c_t 's are *iid* $N(0, \sigma_c^2)$

For model (9) [generated by Equations (7) and (8)]

$$\theta = \sigma_e^2/\sigma_c^2 \quad (10)$$

and

$$(1 - \theta)^2 = \sigma_e^2/\sigma_c^2.$$

Therefore, assuming the above approximations to the population dynamics, fitting model (9) to the observed survey series provides an estimate, $\hat{\theta}$, of σ_e^2/σ_c^2 and an estimate of σ_c^2 . The π -weights for the model are from Equation (5) given by

$$\pi_i = (1 - \hat{\theta}) \hat{\theta}^{i-1}, \quad i \geq 1. \quad (11)$$

It may be noted that if model (9) is valid and catchability is constant over time then the estimate of σ_e^2 given by $\hat{\theta} \hat{\sigma}_c^2$ [from Equation (10)] would approximately equal the estimate of σ_e^2 based on the within survey sampling variance.

AN APPLICATION

The Northeast Fisheries Center conducts an extensive groundfish trawl survey as part of its MARMAP program two times a year: in the fall since 1963 and in the spring since 1968 (Grosslein 1969). The survey region is divided into sampling strata based on geographic boundaries and depth contours (Fig. 2). For each survey, trawl stations are chosen randomly within each stratum. One of the objectives of the surveys is to provide indices of abundance for the many species of commercial value in the region.

Yellowtail flounder is an important New England fishery resource whose population has fluctuated considerably over the survey period (Clark et al. 1984). Commercial catch statistics exist for yellowtail flounder, but age data suitable for a VPA (Virtual Population Analysis) are unavailable. Major




FIGURE 2.—The National Marine Fisheries Service's MARMAP survey strata.

yellowtail flounder fisheries are off southern New England (strata 5, 6, 8, 9) and on Georges Bank (strata 13-21). The two stocks are fairly distinct but with some intermixing (Clark et al. 1984).

The nonzero catch per tow survey data for yellowtail flounder are approximately lognormally distributed within a stratum. Therefore, the estimators based on the Δ -distribution [Equations (1) and (2)] were used to estimate the mean catch per tow and its variance in each stratum. The regional estimates for southern New England and Georges Bank were then calculated in the usual manner for each survey (see, e.g., Pennington and Brown 1981).

Model (9) was fit to each series (spring 1968-84 and fall 1963-84 in both regions) and the model's adequacy checked (Box and Jenkins 1976, Chap. 8). Table 1 contains summary statistics and parameter

estimates for the four series. Since the series are relatively short, the averages of the areal and seasonal estimates are used as the final estimates of θ and σ_c^2 (last line in Table 1).

Abundance indices for the two regions and seasons were calculated by applying to each series Equation (4) with $\hat{\theta} = 0.4$, the π -weights given by Equation (11), and the $\hat{\epsilon}_t$'s (for each series) generated by model (9). An estimate of σ_c^2 equal to 0.20 and of σ_n^2 equal to 0.18 were obtained from Equation (10). The estimated variance of the index equals, from Equation (6), 0.12 for the current value and declines to 0.09 for values not near the series' end points. This compares with a variance of 0.20 ($= \sigma_c^2$) for the original index. Figures 3 (log scale) and 4 (linear scale) show plots of the estimated index and the observed catch per tow series for the fall surveys off southern New England.

DISCUSSION

The major advantage of estimating an index of abundance from the entire survey series is that it can produce an index with a variance considerably smaller than the variance of the observed series. But the application also demonstrates that estimates of the accuracy of an index based only on the within survey sampling variance can be misleading. For example, the 1972 survey value for yellowtail flounder off southern New England is considered an anom-

TABLE 1.—Summary statistics and parameter estimates for the yellowtail flounder survey series. The first three sample autocorrelations (r_1 , r_2 , and r_3) are for the first differenced logged series.

Survey	No. of years	r_1	r_2	r_3	$\hat{\theta}$	SE($\hat{\theta}$)	σ_c^2
Southern New England							
Spring	17	-0.23	0.12	-0.18	0.21	0.28	0.57
Fall	22	-0.26	0.07	-0.31	0.40	0.22	0.71
Georges Bank							
Spring	17	-0.32	0.00	-0.09	0.61	0.23	0.36
Fall	22	-0.30	-0.06	0.18	0.36	0.23	0.33
Average		-0.28	0.03	-0.10	0.40	0.12	0.50

¹Assuming the estimates of θ are independent.




FIGURE 3.—Logged average catch per tow and the estimated index of abundance for southern New England yellowtail flounder.




FIGURE 4.—Average catch per tow and the estimated index of abundance for southern New England yellowtail flounder.

aly (Collie and Sissenwine 1983). It does appear anomalous if comparisons are made using 0.11, the estimated variance based on the within survey variance, but not if the estimate of 0.20 ($= \hat{\sigma}_a^2$) is considered (Fig. 3).

Assessing the accuracy of an index of abundance for marine stocks is difficult since the true levels are never known with certainty. But they can be compared with other indicators of abundance. The methods were applied to the haddock stock on Georges Bank (Pennington 1985) for which a VPA exists. It was found that model (7) adequately describes the dynamics of the VPA series, and the survey series follows model (9). The resulting index of abundance is quite similar to the VPA estimates.

Collie and Sissenwine (1983) give a method for estimating the relative abundance of a fish stock using survey data and commercial catch statistics. They observe that their method produces estimates which compare favorably with VPA estimates. Figure 5 shows plots of Collie and Sissenwine's estimate of the relative abundance of southern New England yellowtail flounder and the index based only on the survey data.

Finally, it should be noted that the purpose of the modeling stage in the estimation procedure is not necessarily to develop a realistic model for the population, but to describe the important stochastic properties of the series. As the observed series becomes longer, more precise estimates can be made. For shorter series, given the large variabil-

ity inherent in marine trawl surveys, a preliminary estimate of between 0.3 and 0.4 for the smoothing parameter θ appears to be an appropriate initial value to use for estimating an abundance index until more information becomes available.

LITERATURE CITED

- AITCHISON, J., AND J. A. C. BROWN.
1957. The lognormal distribution. Cambridge Univ. Press, Lond., 176 p.
- BOX, G. E. P., AND G. M. JENKINS.
1976. Time series analysis: forecasting and control. Rev. ed. Holden-Day, San Franc., 575 p.
- BYRNE, C. J., T. R. AZAROVITZ, AND M. P. SISENWINNE.
1981. Factors affecting variability of research trawl surveys. Can. Spec. Publ. Aquat. Sci. 58:238-273.
- CLARK, S. H.
1979. Application of bottom trawl survey data to fish stock assessment. Fisheries 4:9-15.
- CLARK, S. H., M. M. MCBRIDE, AND B. WELLS.
1984. Yellowtail flounder assessment update. U.S. Dep. Commer., NOAA, Natl. Mar. Fish. Serv., Woods Hole Lab. Ref. Doc. No. 84-39, 29 p.
- COLLIE, J. S., AND M. P. SISENWINNE.
1983. Estimating population size from relative abundance data measured with error. Can. J. Fish. Aquat. Sci. 40: 1871-1879.
- GROSSLEIN, M. D.
1969. Groundfish survey program of BCF Woods Hole. Comm. Fish. Rev. 31(8-9):22-30.
- PENNINGTON, M.
1983. Efficient estimators of abundance, for fish and plankton surveys. Biometrics 39:281-286.
1985. Estimating the relative abundance of fish from a series




FIGURE 5.—Survey index of abundance (solid line) and Collie and Sissenwine's index (broken line) for southern New England yellowtail flounder.

of trawl surveys. *Biometrics* 41:197-202.

PENNINGTON, M., AND P. BERRIEN.

1984. Measuring the precision of estimates of total egg production based on plankton surveys. *J. Plankton. Res.* 6:

869-879.

PENNINGTON, M., AND B. E. BROWN.

1981. Abundance estimators based on stratified random trawl surveys. *Can. Spec. Publ. Aquat. Sci.* 58:149-153.